

**UP AND UP DAYTIME NIGHTTIME CHILDRENS MULTI SYMPTOM COLD-
acetaminophen, dextromethorphan hydrobromide, diphenhydramine
hydrochloride, guaifenesin, and phenylephrine hydrochloride
Target Corporation**

Up&up Daytime Nighttime Children's Multi Symptom Cold Drug Facts

Day Time Multi-Symptom Cold

DO NOT TAKE BOTH PRODUCTS WITHIN 4 HOURS OF EACH OTHER

Drug Facts

<i>Active ingredients (in each 5 mL)</i>	<i>Purposes</i>
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help your child get to sleep
 - nasal congestion due to a cold
 - stuffy nose

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- **each 5 mL contains:**sodium 2 mg
- store at room temperature
- do not refrigerate
- dosing cup provided

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue No.1, FD&C Red #40, flavors, potassium citrate, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Nighttime Children's Multi-Symptom Cold

DO NOT TAKE BOTH PRODUCTS WITHIN 4 HOURS OF EACH OTHER

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5	Antihistamine/cough

mg	suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose
 - sneezing
 - cough
- controls cough to help your child get to sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert:Acetaminophen may cause severe skin reactions.

Symptoms may include:

- Skin reddening
- Blisters
- Rash

Sore throat warning:if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other drug containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if your child has

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed (**see Overdose warning**)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- **children 6 to under 12 years of age:** 10 mL in dosing cup provided every 4 hours; do not give more than 5 doses in any 24-hour period
- **children under 6 years of age:** do not use

Other information

- **each 10 mL contains:**sodium 4 mg
- store at room temperature
- do not refrigerate

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, potassium citrate, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions?

1-866-467-2748.

Dist. by: Target Corporation

Minneapolis, MN 55403

Made In the U.S.A.

With domestic and imported ingredients.

TM & © 2024 Target Brands, Inc.

PRINCIPAL DISPLAY PANEL - Kit Carton**VALUE
PACK**

Compare to active ingredients in Children's Mucinex® Multi-Symptom Cold* and Children's Mucinex® Nighttime Multi-Symptom Cold**

NDC 82442-476-08

Daytime Nighttime Children's Multi-Symptom Cold

Dextromethorphan HBr 5 mg (Cough Suppressant)

Guaifenesin 100 mg (Expectorant)

Phenylephrine HCl 2.5 mg (Nasal Decongestant)

- **Breaks up Mucus**
- **Provides relief from stuffy nose, cough and chest congestion**

Age 4 to 11 Years**Dosage Cup Included**

Berry Flavor

NATURALLY AND ARTIFICIALLY FLAVORED

4 FL OZ (118 mL)

NIGHTTIME

Multi-Symptom Cold

Acetaminophen 325 mg(Pain Reliever/Fever Reducer)

Diphenhydramine HCl 12.5 mg (Antihistamine/Cough Suppressant)

Phenylephrine HCl 5 mg (Nasal Decongestant)

- Provides relief from sneezing, stuffy or runny nose, cough, fever and sore throat

Very Berry Flavor

NATURALLY AND ARTIFICIALLY FLAVORED

Ages 6 to 11 years

Dosage Cup Included

4 FL OZ (118 mL)

8 FL OZ (236 mL) TOTAL

*** This product is not manufactured or distributed by RB Health (US), distributor of Children's Mucinex® Multi-Symptom Cold.**

****This product is not manufactured or distributed by RB Health (US), distributor of Children's Mucinex® Nighttime Multi-Symptom Cold.**

TAMPER EVIDENT: DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS BROKEN OR MISSING.

Important: Keep this carton for future reference for full labeling.



UP AND UP DAYTIME NIGHTTIME CHILDRENS MULTI SYMPTOM COLD

acetaminophen, dextromethorphan hydrobromide, diphenhydramine hydrochloride, guaifenesin, and phenylephrine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82442-476
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-476-08	1 in 1 CARTON; Type 0: Not a Combination Product	07/19/2024	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL
Part 2	1 BOTTLE	118 mL

Part 1 of 2

DAYTIME CHILDRENS MULTI SYMPTOM COLD

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:82442-469
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:82442-469-04	118 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
---	------------------	---	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/19/2024	

Part 2 of 2

UP AND UP NIGHTTIME CHILDRENS MULTI SYMPTOM COLD

acetaminophen, diphenhydramine hydrochloride, and phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:82442-377
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	BERRY (Very)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82442-377-04	118 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/19/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/19/2024	

Labeler - Target Corporation (006961700)