

**IBUPROFEN- ibuprofen tablet  
WALGREENS CO.**

-----  
**Ibuprofen Tablets USP, 200mg  
Pain Reliever/ Fever Reducer (NSAID)**

**Active ingredient  
(in each tablet)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever / Fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Allergy Alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of the reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

### **Other information**

- store between 20-25°C (68-77°F). [See USP Controlled Temperature]
- avoid excessive heat above 40°C (104°F)

### **Inactive ingredients**

colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide.

### **Questions or comments?**

call 1-877-770-3183 weekdays 8:00 AM EST to 5:00PM PST.

### **Principal Display Panel**



COATING FREE AREA

700000004608  
200000003489

PAPER BOX  
PLASTIC BOTTLE

**Ibuprofen**  
TABLETS USP, 200 mg  
PAIN RELIEVER / FEVER REDUCER (NSAID)

**WALGREENS**

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - stomach pain
  - black stools
  - blood in stool
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the pain relief area
- any new symptoms appear

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years

- ask a doctor

**Other information**

- store between 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature].
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, povidone K-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?**  
call 1-877-770-5183 weekdays 8:00 AM EST to 5:00 PM PST

**Ibuprofen**  
TABLETS USP, 200 mg  
PAIN RELIEVER / FEVER REDUCER (NSAID)

**15 COATED TABLETS**

**WALGREENS**

Compare to the active ingredient in Advil® Tablets

NDC 0383-9594-07

ACTUAL SIZE

**\$2**

Contains no ingredient made from a gluten-containing grain (wheat, barley or rye). Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Wyeth Consumer Healthcare, distributor of Advil® Tablets.

REGISTERED BY:  
WALGREEN CO.  
DEERFIELD, IL 60015  
LAWN SUBSCRIPTION  
GUARANTEED

walgreens.com  
©2003 Walgreen Co.  
MADE IN INDIA  
Lot  
Exp.

ITEM 751722 W10383-0823-L

3 11917 01961 1

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THE BOX FOR IMPORTANT INFORMATION. DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING.**

**Drug Facts**

**Active ingredient (in each tablet) Purpose**  
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
  - temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

COATING FREE AREA

200000004608  
700000003489

W3ORG0523-F

COATING FREE AREA

1 3/4      1 3/4

Principal Display Panel



## IBUPROFEN

ibuprofen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-9594
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

## Inactive Ingredients


Ingredient Name	Strength
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>POLYETHYLENE GLYCOL 4000</b> (UNII: 4R4HFI6D95)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

## Product Characteristics

<b>Color</b>	brown ((Reddish Brown color Tablets))	<b>Score</b>	score with uneven pieces
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9594-10	100 in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	10/15/2021	
2	NDC:0363-9594-07	15 in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2023	



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	10/15/2021	

**Labeler** - WALGREENS CO. (008965063)

Revised: 12/2025

WALGREENS CO.