

**ACETAMINOPHEN- acetaminophen tablet
WALGREEN CO.**

Acetaminophen 500mg

Active ingredient

(in each Caplet)

Acetaminophen, USP 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Liver warning

This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days

- new symptoms occur

- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Keep out of the reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

ask a doctor

Other information

- store between 20-25°C (68-77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for lot number and expiration date

Inactive ingredients

hydroxypropyl methyl cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?

call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST.



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9595
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z 65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ 989GH94E)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics

Color	white	Score	score with uneven pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	G;551
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9595-10	100 in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	10/15/2021	

Walgreens
Pain Reliever
ACETAMINOPHEN USP, 500 mg
PAIN RELIEVER / FEVER REDUCER

COMBO PACK
Walgreens
Pain Reliever
ACETAMINOPHEN USP, 500 mg
PAIN RELIEVER / FEVER REDUCER

EXTRA STRENGTH CAPLETS

100 CAPLETS**
(**CAPSULE-SHAPED TABLETS)

ACTUAL SIZE

Drug Facts (continued)
 ■ redness or swelling is present
 These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of the reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ do not take more than directed (see overdose warning)
 adults and children 12 years and over ■ take 2 caplets every 6 hours while symptoms last
 ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
 ■ do not use for more than 10 days unless directed by a doctor
 children under 12 years ask a doctor

Other information
 ■ store between 20-25°C (68-77°F). See USP Controlled Room Temperature
 ■ avoid high humidity
 ■ See end panel for lot number and expiration date

Inactive ingredients
 hydroxypropyl methyl cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?
 call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

CONTAINS NO ASPIRIN

Pharmacist Recommended
 Health expertise you rely on.™

100% SATISFACTION GUARANTEED
 walgreens.com ©2021 Walgreen Co. MADE IN INDIA

ORG0321-F

ITEM 323865 W10383-0321-L
 3 11917 03237 5

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/15/2021	

Labeler - WALGREEN CO. (008965063)

Revised: 12/2025

WALGREEN CO.