

CHILDRENS BENADRYL ALLERGY PLUS CONGESTION- diphenhydramine hydrochloride and phenylephrine hydrochloride solution
Johnson & Johnson Consumer Inc.

Childrens Benadryl Allergy Plus Congestion

Drug Facts

<i>Active ingredients (in each 5 mL)</i>	<i>Purposes</i>
Diphenhydramine HCl 12.5 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
 - nasal congestion
 - stuffy nose
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion
 - stuffy nose
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes

- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dose**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- find right dose on chart below
- mL = milliliter
- take every 4 hours
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 4 years	do not use
children 4 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL
adults and children 12 years and over	10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: **sodium 10 mg**
- store between 20-25°C (68-77°F)
- **do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing**

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

call **1-877-717-2824** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-341-01

Children's
Benadryl[®]

ALLERGY PLUS
CONGESTION

Diphenhydramine HCl / antihistamine
Phenylephrine HCl / nasal decongestant
oral solution

4 Hours/Dose

RELIEF OF:

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Sinus Congestion
- Stuffy Nose

Alcohol Free
Sugar Free

ALLERGY + CONGESTION RELIEF

Grape!
Flavored Liquid

4 fl oz (118 mL)

Children's
Benadryl[®]

ALLERGY PLUS
CONGESTION

How can we help?
☎ 1-877-717-2824
benadryl.com

NDC 50580-341-01

Children's
Benadryl[®]

ALLERGY PLUS
CONGESTION

Diphenhydramine HCl / antihistamine
Phenylephrine HCl / nasal decongestant
oral solution

4 Hours/Dose

RELIEF OF:

- ✔ Runny Nose
- ✔ Sneezing
- ✔ Itchy, Watery Eyes
- ✔ Itchy Throat or Nose
- ✔ Sinus Congestion
- ✔ Stuffy Nose

Alcohol Free
Sugar Free



Actual Size

Tips for Outdoor Allergens:



- Avoid being outside when the pollen count is high, typically in the morning
- Shower and change clothing after spending time outdoors

Tips for Indoor Allergens:



- Wash sheets and blankets in hot water and use the dryer weekly



• Minimize contact and keep pets out of the bedroom to help with pet allergies

4 fl oz (118 mL)



Drug Facts (continued)

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help

Children's
Benadryl[®]

ALLERGY PLUS

CONGESTION

Drug Facts

Active ingredients (in each 5 mL)	Purposes
Diphenhydramine HCl 12.5 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
 - nasal congestion
 - stuffy nose
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion
 - stuffy nose
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose
- marked drowsiness may occur
- avoid alcoholic drinks

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- find right dose on chart below
- mL = milliliter
- take every 4 hours
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 4 years	do not use
children 4 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL
adults and children 12 years and over	10 mL

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F)
- do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

call 1-877-717-2824 (toll-free) or 215-273-8755 (collect)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

©J&JCI 2019

30046178

For more information, visit:
www.benadryl.com

CHILDRENS BENADRYL ALLERGY PLUS CONGESTION

diphenhydramine hydrochloride and phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-341
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-341-01	1 in 1 CARTON	09/30/2014	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2014	

Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 1/2024

Johnson & Johnson Consumer Inc.