STOMACH RELIEF ULTRA STRENGTH- bismuth subsalicylate liquid QUALITY CHOICE (Chain Drug Marketing Association)

Drug Facts

Active ingredient (in each 15 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of childen.

In case of overdose, get medical help or contact a Posion Control Center (1-800-222-1222) right away.

Directions

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
 - 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 - 15 mL (1 dose) every 1/2 as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

Other information

each 15 mL contains: sodium 5 mg
each 15 mL contains: salicylate 206 mg

- low sodium
- keep tightly closed

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

*Compare to the active ingredient in **Pepto-Bismol® Ultra**

Ultra Stomach Relief

Stomach Relief

Bismuth subsalicylate 525 mg | Upset Stomach Reliever,

Antidiarrheal

For Relief of::

- heartburn
- indigestion
- nausea
- upset stomach
- diarrhea

2x strength per ouncet

alcohol free

sugar free

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Distributed by C.D.M.A., Inc.©

43157 W 9 Mile Rd

Novi, MI 48375

www.qualitychoice.com

Package Label

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

se of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions on to do not take more than 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea mL = millilite meshake well before using memeasure only with dosing cup provided. Do not use any other dosing device. Reep dosing cup with product devicts and children 12 years and over. The first multiple for first multi every hour as needed for diarrhea/traveler's diarrhea ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) - children under 12 years of age: ask a doctor

Other information = each 15 mL contains: sodium 5 mg ■ each 15 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call 1-800-935-2362 Monday-Friday 9AM-5PM EST



This product is not manufactured or distributed by The Procter & Gamble Company. Pepto Bismol[®] is a registered trademark of The Procter & Gamble Company. *vs. Original Pepto-Bismot*

PLD-B494A LB008591





NDC 63868-813-08

*Compare to the Active Ingredient in Pepto Bismol® Ultra

Ultra Stomach Relief

Stomach Relief

Bismuth Subsalicylate | Upset Stomach Reliever, Antidiarrheal

For Relief of:

Heartburn Indigestion

Diarrhea

Nausea Upset Stomach

2x strength per ounce

Alcohol Free Sugar Free

8 fl oz (237 mL)

Drug Facts
TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR N

Active ingredient (in each 15 mL dose) Bismuth subsalicylate 525 mg. **Purpose**

..Upset stomach reliever/Antidiarrheal

Uses relieves

■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including metartburn indigestion meausea gas fullness belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

- Allergy alert: Contains salicylate. Do not take if you are
- allergic to salicylates (including aspirin)
 taking other salicylate products

Do not use if you have
■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have m fever m mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ diabetes ■ gout ■ arthritis anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

QUALITY CHOICE Stomach Relief

STOMACH RELIEF ULTRA STRENGTH

bismuth subsalicylate liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:63868-813

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION -**BISMUTH** 525 ma UNII:ZS9CD1I8YE) **SUBSALICYLATE** in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 85KN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868- 813-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/25/2021	06/25/2026
Marketing Information				
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
01	C Monograph D	rug M008	06/25/2021	06/25/2026

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 10/2024 QUALITY CHOICE (Chain Drug Marketing Association)