

OXYCARE II- hydrogen peroxide, lactic acid, dodecyl benzene sulfonic acid solution
Ecolab Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

FOR INDUSTRIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

WARNING

Causes eye irritation. Wash skin thoroughly after handling. Do not mix with bleach or other chlorinated products – will cause chlorine gas.

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.

Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh OxyCare II, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Using fresh OxyCare II, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, and using fresh OxyCare II, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Using fresh OxyCare II, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: OxyCare II is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

FIRST AID

IN CASE OF EYE CONTACT: Rinse with plenty of water.

IN CASE OF SKIN CONTACT: Rinse with water.

IF SWALLOWED: Rinse mouth. Get medical attention if symptoms occur.

IF INHALED: Get medical attention if symptoms occur.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0026. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

Representative Label and Principal Display

A&L Laboratories

OxyCare™ II

Sanitizing Teat Dip

- An aid in reducing the spread of organisms which may cause mastitis

ACTIVE INGREDIENT:

Hydrogen peroxide.....	0.5%
Lactic acid.....	1.7%
Dodecyl benzene sulfonic acid.....	0.5%

SKIN CONDITIONING SYSTEM:

(contains sorbitol, glycerin).....	8.0%
------------------------------------	------

INERT INGREDIENTS:	89.3%
---------------------------------	-------

TOTAL:	100.0%
---------------------	--------

Manufactured by

Ecolab Inc. · 370 Wabasha Street N · St Paul MN 55102-1390 USA · tel: 1 800 392 3392

© 2015 Ecolab USA Inc. · All rights reserved

Made in United States

761351/5300/0315

NET CONTENTS: 208 L (55 US GAL)

6301555



OxyCare™ II

Sanitizing Teat Dip
Desinfectante de pezones por inmersión

• An aid in reducing the spread of organisms which may cause mastitis

ACTIVE INGREDIENT:	
Hydrogen peroxide.....	0.5%
Lactic acid.....	17%
Dodecyl benzene sulfonic acid.....	0.5%
SKIN CONDITIONING SYSTEM:	
(contains sorbitol, glycerin).....	8.0%
INERT INGREDIENTS:	89.3%
TOTAL:	100.0%

FOR INDUSTRIAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

NOT FOR HUMAN USE

DIRECTIONS:

IMPORTANT: Do not further dilute with water or mix with any other teat dips. If product in dip cup becomes visibly dirty, discard contents and replenish with fresh product. Do not reuse or return any unused product to the original container.
Udder Prep: When using an udder wash step before milking, make sure to wash teats with appropriate udder wash solution using proper cleaning procedures. Teats should then be dried with single-service towels.

Directions for Teat Dipping

Pre-Milk Dipping: Before each cow is milked, and using fresh OxyCare II, dip each teat full-length into the teat dip cup. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Dipping: Using fresh OxyCare II, dip each teat full-length into the teat dip cup. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Directions for Teat Spraying

Pre-Milk Spraying: Before each cow is milked, and using fresh OxyCare II, spray entire teat. Wipe teats after dipping using single-service towels to avoid contamination of milk.

Post-Milk Spraying: Using fresh OxyCare II, spray entire teat immediately after each milking. Do not wipe. Allow to air dry. Do not turn cows out in freezing weather until the product is completely dry.

Expanded Usage: When freshening cows, begin dipping teats twice daily for about 10 days before calving.

PRECAUTION: OxyCare II is not intended to cure or help the healing of chapped or irritated teats. As with any germicide, irritation or sensitization may occur in sensitive animals. In case of teat irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

Consult your Ecolab representative for specific use instructions and recommended dispensing equipment.

WARNING

Causes eye irritation. Wash skin thoroughly after handling. Do not mix with bleach or other chlorinated products - will cause chlorine gas.

FIRST AID

IN CASE OF EYE CONTACT: Rinse with plenty of water.

IN CASE OF SKIN CONTACT: Rinse with water.

IF SWALLOWED: Rinse mouth. Get medical attention if symptoms occur.

IF INHALED: Get medical attention if symptoms occur.

READ SAFETY DATA SHEET (SDS) BEFORE USING THIS PRODUCT

EMERGENCY HEALTH INFORMATION: 1 800 328 0026. If located outside the United States and Canada, call collect 1 651 222 5352 (number is in the US).

• Ayuda a evitar la proliferación de organismos que pueden causar mastitis

PARA USO INDUSTRIAL SOLAMENTE

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

NO ES PARA USA HUMANO

INSTRUCCIONES:

IMPORTANTE: No lo vuelva a diluir con agua ni lo mezcle con otros productos para baño de pezones. Si el producto del recipiente para realizar el baño está visiblemente sucio, deseché el contenido y rellene el recipiente con producto fresco. No vuelva a utilizar el producto ni vuelva a colocar producto sin usar en el contenedor original.

Preparado para ubres: Si se realiza un paso de limpieza de ubres antes del ordeño, asegúrese de limpiar los pezones con una solución para limpieza de ubres adecuada y procedimientos de limpieza apropiados. Los pezones deben secarse con toallas de uso único.

Instrucciones para el sellado de pezones

Sellado del pezón previo al ordeño: Antes de ordeñar cada vaca, bañe toda la superficie de cada pezón en un recipiente que contenga OxyCare II fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Sellado del pezón posterior al ordeño: Bañe toda la superficie de cada pezón en un recipiente que contenga OxyCare II fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo.

Instrucciones para el rociado de pezones

Rociado previo al ordeño: Antes de ordeñar cada vaca, rocíe toda la superficie de cada pezón con OxyCare II fresco. Limpie los pezones con toallas de uso único luego del baño para evitar la contaminación de la leche.

Rociado de pezones posterior al ordeño: Inmediatamente después del ordeño, rocíe toda la superficie de cada pezón con OxyCare II fresco. No utilice paños. Deje secar al aire. No deje las vacas en un clima helado hasta que el producto se haya secado por completo.

Uso Expandido: Al refrescar a las vacas, comience con un baño antiséptico de sus pezones dos veces por día por aproximadamente 10 días antes de parir.

PRECAUCIÓN: OxyCare II no está diseñado para curar o ayudar con la cicatrización de los pezones agrietados o irritados.

Como cualquier otro germicida, puede provocar irritación o sensibilización en animales sensibles. En caso de irritación o agrietamiento de los pezones, haga examinar dichas afecciones y, de ser necesario, hágalas tratar con un veterinario.

Consulte con su Representante de Ecolab para obtener instrucciones específicas sobre el uso y el equipo dispensador recomendado.

ATENCIÓN

Provoca irritación ocular. Lave la piel minuciosamente después del manejo. No se mezcle con el blanqueo o otros productos tratados con cloro - causará el gas de cloro.

PRIMEROS AUXILIOS

EN CASO DE CONTACTO CON LOS OJOS: Enjuague con mucha agua.

EN CASO DE CONTACTO CON LA PIEL: Enjuague con agua.

EN CASO DE INGESTIÓN: Enjuague la boca. Consultar un médico si los síntomas aparecen.

EN CASO DE INHALACIÓN: Consultar un médico si los síntomas aparecen.

LEA LA HOJA DE DATOS DE SEGURIDAD ANTES DE USAR ESTE PRODUCTO

INFORMACIÓN DE EMERGENCIA SOBRE SALUD: 1 800 328 0026. Si está fuera de los Estados Unidos y Canadá, llame a cobro revertido al 1 651 222 5352 (número en los EE.UU.).

Manufactured by
Ecolab Inc. • 370 Wabasha Street N • St Paul MN 55102-1390 USA • tel: 1 800 392 3392
© 2015 Ecolab USA Inc. • All rights reserved
Made in United States
761351/530Q/0315

NET CONTENTS: 208 L (55 US GAL)

6301555



OXYCARE II

hydrogen peroxide, lactic acid, dodecyl benzene sulfonic acid solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:47593-538
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	5 mg in 1 mL
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)	LACTIC ACID	17 mg in 1 mL
DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)	DODECYLBENZENESULFONIC ACID	5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

SORBITOL (UNII: 506T60A25R)				
GLYCERIN (UNII: PDC6A3C0OX)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47593-538-17	56800 mL in 1 DRUM		
2	NDC:47593-538-21	208000 mL in 1 DRUM		
3	NDC:47593-538-44	1150000 mL in 1 PACKAGE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
unapproved drug other			04/27/2015	

Labeler -
Ecolab Inc. (006154611)