SUPHEDRINE PE SINUS CONGESTION NON-DROWSY- phenylephrine hcl tablet, film coated

ARMY AND AIR FORCE EXCHANGE SERVICE

Exchange Select 44-453-Sinus

Active ingredient (in each tablet)

Phenylephrine HCl 10 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over: take 1 tablet every 4 hours. Do not take more than 6 tablets in 24 hours.
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dextrose monohydrate, dibasic calcium phosphate dihydrate, FD&C red #40, lecithin, magnesium stearate, maltodextrin, microcrystalline cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium citrate dihydrate, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

exchange **∕ select**™

Compare To The Active Ingredient of SUDAFED PE® SINUS CONGESTION*

SUPHEDRINE PE SINUS CONGESTION

Phenylephrine HCl 10 mg, Nasal Decongestant

- Maximum Strength
- Sinus Pressure
- Congestion

NON-DROWSY

Actual Size

72 Tablets

quality value

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN

OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark SUDAFED PE® SINUS CONGESTION.

50844 REV0820L45323

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"

Manufactured For Your Military Exchanges Distributed by: LNK International, Inc. Hauppauge, NY 11788 1-800-426-9391

SUPHEDRINE PE

SINUS CONGESTION

SUPHEDRINE SINUS CONGESTION

exchange select

Compare To The Active Ingredient of SUDAFED PE® SINUS CONGESTION*

PHEDRIN CONG

Phenylephrine HCl 10 mg, Nasal Decongestant

- Maximum Strength
- Sinus Pressure
- Congestion

NON-DROWSY

Tablets

quality value



PACTON GOATAN TEACHANG es Manufactured For Your Military Exchange, Distributed by: LNK International, Inc. Hauppauge, NY 1178 1-800-426-9291 "SATISFACTION GUARANTEED OR YOUR MONEY BACK"

50844 REV0820L45323

Johnson Corporation, owner of the registered trademark SUDAFED PE® SINUS CONGESTION. This product is not manufactured or distributed by Johnson &

Questions or comments? 1-800-426-9391

citrate dinydrate, titanium dioxide cellulose, silicon dioxide, sodium carboxymethylcellulose, sodium lecithin, magnesium stearate, maltodextrin, microcrystalline monohydrate, dibasic calcium phosphate dihydrate, FD&C red #40 Inactive ingredients croscarmellose sodium, dextrose

- see end flap for expiration date and lot number
- store at 25°C (7°F); excursions permitted between 15°-30°C
 - TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS Opened or blister is torn or broken Uther information
 - children under 12 years: ask a doctor
- hours. Do not take more than 6 tablets in 24 hours.
- adults and children 12 years and over: take 1 tablet every 4

Directions

help or contact a Poison Control Center right away. Keep out of reach of children. In case of overdose, get medical

If pregnant or breast-feeding, ask a health professional before Drug Facts (continued) ■ symptoms do not improve within 7 days or occur with fever ■ uetvousness, dizziness, or sleeplessness occur Stop use and ask a doctor if

When using this product do not exceed recommended dosage.

- quilicuity in urination due to enlargement of the prostate gland
 - thyroid disease high blood pressure
 - diabetes

Ask a doctor before use if you have

before taking this product prescription drug contains an MAOI, ask a doctor or pharmacist after stopping the MAOI drug. If you do not know if your oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks

Do not use it you are now taking a prescription monoamine

- hay fever or other upper respiratory allergies
- temporarily relieves nasal congestion due to the common cold, ■ temporarily relieves sinus congestion and pressure

sasn

Nasal decongestant Purpose

Phenylephrine HCI 10 mg Active ingredient (in each tablet)

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

44-453

SUPHEDRINE PE SINUS CONGESTION NON-DROWSY

phenylephrine hcl tablet, film coated

TAMPER EVIDENT. DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



Product Information

NDC:55301-753 **Product Type HUMAN OTC DRUG Item Code (Source)**

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -

UNII:1WS297W6MV)

PHENYLEPHRINE **HYDROCHLORIDE**

10 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)

TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	44;453
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301- 753-44	1 in 1 CARTON	01/15/2021	
1		18 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:55301- 753-23	3 in 1 CARTON	01/15/2021	
2		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Application Number or Monograph Marketing Start Marketing End Marketing

Category	Citation	Date	Date
OTC Monograph Drug	M012	01/15/2021	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55301-753) , pack(55301-753)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55301-753)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(55301-753)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		117025878	manufacture(55301-753)	

Revised: 1/2025 ARMY AND AIR FORCE EXCHANGE SERVICE