

**ANTACID PLUS- aluminum hydroxide, magnesium hydroxide,
dimethicone liquid
BETTER LIVING BRANDS, LLC**

SC ANTACID original

Active ingredients (in each 10 mL dose)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)
Magnesium hydroxide 400 mg
Simethicone 40 mg

Purposes

Antacid
Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Do not take more than 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- children under 12 years: ask a doctor
- measure with dosing cup provided

- mL = milliliter

Other information

- **each 10 mL dose contains:** magnesium 165 mg, sodium 6 mg
- store at room temperature tightly closed
- protect from freezing

Inactive ingredients

benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package Label



Compare to
MYLANTA®
REGULAR STRENGTH
active ingredients*

Quality
Guaranteed
NDC 21130-629-12

Antacid Plus Regular Strength

Alumina, Magnesia, and Simethicone
Oral Suspension USP

Fast Acting
Soothing Relief of:

- Acid Indigestion
- Heartburn
- Sour Stomach
- Pressure & Bloating

Contains less than
0.15% Alcohol

Original
Flavored Liquid

12 FL OZ (355 mL)

285-05112-0 REV GC0820

Drug Facts

TAMPER-EVIDENT: Do not use if imprinted neckband is missing or broken.

Active ingredients (in each 10 mL dose)

Aluminum hydroxide (equivalent to dried gel, USP) 400 mg..... Antacid
Magnesium hydroxide 400 mg Antacid
Simethicone 40 mg Antigas

Purposes

Uses relieves: • heartburn • sour stomach
• acid indigestion • symptoms referred to as gas

Warnings

Do not take more than 120 mL (12 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

Ask doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals as needed, at bedtime, or as directed by a doctor
- children under 12 years of age: ask a doctor
- measure with dosing cup provided
- mL = milliliter

Other information

- each 10 mL dose contains: magnesium 165 mg, sodium 6 mg
- store at room temperature tightly closed
- protect from freezing

Inactive ingredients benzyl alcohol, butylparaben, flavor (contains alcohol), hydroxyethylcellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765



Scan here
for more
information

*This product is not manufactured or distributed by the owner of the registered trademark MYLANTA®.

985-05112-1 REV GC0621

RD 20313

S6143

DISTRIBUTED BY
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
1-888-723-3929
www.betterlivingbrandsLLC.com
MADE IN U.S.A.



ANTACID PLUS

aluminum hydroxide, magnesium hydroxide, dimethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-629
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -	ALUMINUM	400 mg

UNII:5QB0T2IUN0)	HYDROXIDE	in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0K00R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-629-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	08/01/2020	

Labeler - BETTER LIVING BRANDS, LLC (009137209)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(21130-629)

Revised: 11/2023

BETTER LIVING BRANDS, LLC