RUGBY NICOTINE POLACRILEX GUM, COATED FRUIT FLAVOR- nicotine polacrilex gum, chewing Rugby Laboratories, Inc.

Nicotine Polacrilex Gum 2 mg and 4 mg Coated Fruit - Rugby

Drug Facts

Active ingredient (in each chewing piece) - 2 mg

Nicotine Polacrilex 2 mg (nicotine)

Active ingredient (in each chewing piece) - 4 mg

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

• reduces withdrawal symptoms, including nicotine craving, associated with guitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- · a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions - 2 mg

if you are under 18 years of age, ask a doctor before use

- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every	1 piece every	1 piece every
1 to 2 hours	2 to 4 hours	4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Directions - 4 mg

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every	1 piece every	1 piece every
1 to 2 hours	2 to 4 hours	4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 2 mg

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20 25°C (68 77°F)
- protect from light

Other information - 4 mg

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20 25°C (68 77°F)

protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide and xylitol.

Inactive ingredients - 4 mg

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 lake, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide and xylitol.

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal display panel 2 mg

Compare to the active ingredient in Nicorette® Gum*

Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (Nicotine)

Stop Smoking Aid

2 mg

Coated Fruit Flavor

Sugar-Free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use nicotine Polacrilex Gum USP, 4 mg

TO INCREASE YOUR SUCCESS IN QUTTING:

- 1. You must be motivated to quit.
- 2. **Use Enough** Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
- 3. Use Long Enough Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. **Use With a Support Program** as directed in the enclosed User's Guide.
- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN

Distributed by:

RUGBY® LABORATORIES

Indianapolis, IN 46268

	Pa	cka	ae	labe	12	ma
--	----	-----	----	------	----	----

	RUGBY Nicotine Gum Coated Fruit Flavor

^{*}This product is not manufacture or distributed by Haleon group of companies, distributor of Nicorette® Gum.

Compare to the active ingredient in Nicorette® Gum*

Nicotine Gum

Nicotine Polacrilex Gum USP, 4 mg (Nicotine)

Stop Smoking Aid

4 mg

Coated Fruit Flavor

Sugar-Free

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP

If you smoke your first cigarette <u>MORE THAN</u> 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. **Use Enough** Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
- 3. **Use Long Enough** Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use With a Support Program as Directed in the enclosed User's Guide.
- not for sale to those under 18 years of age
- proof of age required
- · not for sale in vending machines or from any source where proof of age cannot be verified

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed by:

RUGBY® LABORATORIES

Indianapolis, IN 46268

Package label 4 mg

RUGBY Nicotine Gum Coated Fruit Flavor

RUGBY NICOTINE POLACRILEX GUM, COATED FRUIT FLAVOR

Product Information Product Type HUMAN OTC DRUG Item Code (Source) Route of Administration BUCCAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg		

^{*}This product is not manufactured or distributed by Haleon group of companies, distributor of Nicorette® Gum.

Inactive Ingredients			
	Ctth		
Ingredient Name	Strength		
ACACIA (UNII: 5C5403N260)			
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
SODIUM CARBONATE (UNII: 45P3261C7T)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
XYLITOL (UNII: VCQ006KQ1E)			

Product Characteristics				
Color	brown (light brown)	Score	no score	
Shape	SQUARE	Size	14mm	
Flavor	FRUIT (coated fruit)	Imprint Code		
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0536-3386-37	4 in 1 CARTON	07/08/2009	06/30/2016	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:0536-3386-01	10 in 1 CARTON	07/08/2009		
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA079044	07/08/2009		

RUGBY NICOTINE POLACRILEX GUM, COATED FRUIT FLAVOR

nicotine polacrilex gum, chewing

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-3387		
Route of Administration	BUCCAL				

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients			
Ingredient Name	Strength		
ACACIA (UNII: 5C5403N260)			
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)			
SODIUM BICARBONATE (UNII: 8MDF5V39Q0)			
SODIUM CARBONATE (UNII: 45P3261C7T)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
XYLITOL (UNII: VCQ006KQ1E)			

Product Characteristics						
Color	brown (light tan)	Score	no score			
Shape	SQUARE	Size	14mm			
Flavor	FRUIT (coated fruit)	Imprint Code				
Contains						

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-3387-37	4 in 1 CARTON	07/08/2009	06/30/2014
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0536-3387-01	10 in 1 CARTON	07/08/2009	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA079038	07/08/2009			

Labeler - Rugby Laboratories, Inc. (079246066)

Revised: 2/2024 Rugby Laboratories, Inc.