

**GOOD SENSE PAIN RELIEF PM EXTRA STRENGTH- acetaminophen, diphenhydramine
hcl tablet, film coated**

L. Perrigo Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Perrigo Pain Relief PM Drug Facts

Active ingredients (in each caplet)

Acetaminophen 500 mg

Diphenhydramine HCl 25 mg

Purpose

Pain reliever

Nighttime sleep-aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- in children under 12 years of age
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease

- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- new symptoms occur
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> • take 2 caplets at bedtime • do not take more than 2 caplets of this product in 24 hours
children under 12 years	do not use

Inactive ingredients

carnauba wax, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions?

1-800-719-9260

Principal Display Panel

See New Warnings

Extra Strength

Pain Relief PM CAPLETS

Pain Reliever/Nighttime Sleep-Aid

Acetaminophen

Diphenhydramine HCl

For Adults

Actual Size

Compare to active ingredients of Tylenol® PM Caplets

50 Caplets

Non-Habit Forming



EXP. DATE :

EXP.

LOT NO.



Drug Facts

Store at 20-25°C (68-77°F).

**Active ingredients
(in each caplet)**

Purpose

Acetaminophen 500 mg.....Pain reliever
Diphenhydramine HCl 25 mg.....Nighttime sleep-aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

- Do not use**
- in children under 12 years of age
 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - with any other product containing diphenhydramine, even one used on skin
 - if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland ▶

Drug Facts (continued)

- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- new symptoms occur
 - sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 caplets at bedtime ■ do not take more than 2 caplets of this product in 24 hours
children under 12 years	do not use

Drug Facts (continued)

Inactive ingredients: carnauba wax, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, poly sorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions? 1-800-719-9200

Distributed By

Gluten Free

Perrigo
Illigan, MI 49010

**DO NOT USE IF PRINTED
SEAL UNDER CAP IS
BROKEN OR MISSING**

GOOD SENSE PAIN RELIEF PM EXTRA STRENGTH

acetaminophen, diphenhydramine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0 113-0437
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (Light blue)	Score	no score
--------------	-------------------	--------------	----------

Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	L437;PM
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0437-71	1 in 1 CARTON	07/23/1992	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/23/1992	

Labeler - L. Perrigo Company (006013346)

Revised: 12/2019

L. Perrigo Company