

IBUPROFEN ORAL - ibuprofen oral suspension
Bryant Ranch Prepack

Ibuprofen Oral Suspension USP, 100 mg/5 mL (OTC)

ACTIVE INGREDIENT(S)

(in each 5 mL)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/fever reducer

USE(S)

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

WARNINGS

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:☐

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- takes more or for a longer time than directed
- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

DO NOT USE

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

□

- **each 5 mL contains** : sodium 2 mg
- store between 20-25°C (68-77°F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

INACTIVE INGREDIENT SECTION

Berry flavor (with dye)

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Yellow #10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Berry flavor (dye free)

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Grape flavor

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Blue #1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Bubble Gum flavor

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

QUESTIONS SECTION

Call 1-888-588-1418 from 9 AM to 5 PM EST,

Monday-Friday.



Distributed by:
Camber Consumer Care, Inc.
Piscataway, NJ 08854, USA

HOW SUPPLIED

Ibuprofen Oral Suspension Grape
• NDC: 72162-1910-2: 118 mL in a BOTTLE

Repackaged/Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504

Ibuprofen Oral Suspension Grape



LOT
EXP
GTIN



Package
Insert

Drug Facts	
Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	
Uses	
Temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache -reduces fever	
Warnings	
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives/swelling/asthma/fever/rash/low blood pressure/rapid heartbeat. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: child -has had stomach ulcers or bleeding problems/makes a blood thinning (anticoagulant) or steroid drug/takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others/takes more or for a longer time than directed/heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed Sore Throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, rashes, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor	
Other information	
*Each 5 mL contains: sodium 2 mg *Store between 20-25°C (68-77°F) *Do not use if carton is opened or printed bottle is broken or missing -see bottom panel for lot number and expiration date	
Directions	
*This product does not contain directions or complete warnings for adult use-do not give more than directed-shake well before using/mL = milliliter- find right dose on chart. If possible, use weight to dose; otherwise use age -use only enclosed dosing cup. Do not use any other dosing device -if needed, repeat dose every 6-8 hours-do not use more than 4 times a day- replace original bottle cap to maintain child resistance (Weight)(Age)(yr) Dose(mL) under 24-under 2 years-ask a doctor 24-35 lbs- 2-3 years-5 mL 35-47 lbs-4-5 years-7.5 mL 48-59 lbs-6-8 years-10 mL 60-71 lbs-9-10 years-12.5 mL 72-85 lbs- 11 years-15 mL, or as directed by a doctor	
Inactive Ingredients	
Berry flavor (with age): Acesulfame potassium, anhydrous citric acid, carbonylmethylcellulose sodium, D&C Yellow #10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polyorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum. Berry flavor (pib-free): Acesulfame potassium, anhydrous citric acid, carbonylmethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polyorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum. Grape flavor: Acesulfame potassium, anhydrous citric acid, carbonylmethylcellulose sodium, D&C Red #33, FD&C Blue #1, flavors, glycerin, microcrystalline cellulose, polyorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum. Bubble Gum flavor: Acesulfame potassium, anhydrous citric acid, carbonylmethylcellulose sodium, D&C Red #33, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polyorbate 80, propylene	

NDC 72162-1910-2

**Children's Ibuprofen
Oral Suspension (NSAID)
Alcohol Free**

100 mg per 5 mL



**4 FL OZ (118 mL) Grape
Flavor**

Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504 USA

Manufactured by:
AptaPharma Inc.



7216219102

IBUPROFEN ORAL
ibuprofen oral suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72162-1910(NDC:69230-309)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72162-1910-2	1 in 1 CARTON	12/24/2024	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210602	09/18/2019	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(72162-1910) , RELABEL(72162-1910)

Revised: 1/2025

Bryant Ranch Prepack