

**REVOLT- selamectin solution**  
**Aurora Pharmaceutical, Inc.**

-----  
**Revolt® (selamectin)**  
**Topical Parasiticide for Dogs and Cats**

**CAUTION:**

US Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**DESCRIPTION:**

Revolt (selamectin) Topical Parasiticide is available as a colorless to yellow, ready to use solution in single dose tubes for topical (dermal) treatment of dogs six weeks of age and older and cats eight weeks of age and older. The content of each tube is formulated to provide a minimum of 2.7 mg/lb (6 mg/kg) of body weight of selamectin. The chemical composition of selamectin is (5Z,25S)-25-cyclohexyl-4'-O-de(2,6-dideoxy-3-O-methyl- $\alpha$ -L-arabino-hexopyranosyl)-5-demethoxy-25-de(1-methylpropyl)-22,23-dihydro-5-hydroxyiminoavermectin A<sub>1a</sub>.

**INDICATIONS:**

Revolt is recommended for use in dogs six weeks of age or older and cats eight weeks of age and older for the following parasites and indications:

**Dogs:**

Revolt kills adult fleas and prevents flea eggs from hatching for one month and is indicated for the prevention and control of flea infestations (*Ctenocephalides felis*), prevention of heartworm disease caused by *Dirofilaria immitis*, and the treatment and control of ear mite (*Otodectes cynotis*) infestations. Revolt also is indicated for the treatment and control of sarcoptic mange (*Sarcoptes scabiei*) and for the control of tick infestations due to *Dermacentor variabilis*.

**Cats:**

Revolt kills adult fleas and prevents flea eggs from hatching for one month and is indicated for the prevention and control of flea infestations (*Ctenocephalides felis*), prevention of heartworm disease caused by *Dirofilaria immitis*, and the treatment and control of ear mite (*Otodectes cynotis*) infestations. Revolt is also indicated for the treatment and control of roundworm (*Toxocara cati*) and intestinal hookworm (*Ancylostoma tubaeforme*) infections in cats.

**DOSAGE AND ADMINISTRATION:**

The recommended minimum dose is 2.7 mg selamectin per pound (6 mg/kg) of body weight.

Administer the entire contents of a single dose tube (or two tubes used in combination for dogs weighing over 130 pounds) of Revolt topically in accordance with the following tables.

<b>Cats (lb)</b>	<b>Package Color</b>	<b>mg per tube</b>	<b>Potency (mg/mL)</b>	<b>Administered volume (mL)</b>
Up to 5	Rose	15 mg	60	0.25
5.1-15	Blue	45 mg	60	0.75
15.1-22	Taupe	60 mg	60	1.0

For cats over 22 lbs use the appropriate combination of tubes.

<b>Dogs (lb)</b>	<b>Package Color</b>	<b>mg per tube</b>	<b>Potency (mg/mL)</b>	<b>Administered volume (mL)</b>
Up to 5	Rose	15 mg	60	0.25
5.1-10	Purple	30 mg	120	0.25
10.1-20	Brown	60 mg	120	0.5
20.1-40	Maroon	120 mg	120	1.0
40.1-85	Teal	240 mg	120	2.0
85.1-130	Plum	360 mg	120	3.0

For dogs over 130 lbs use the appropriate combination of tubes. Recommended for use in dogs 6 weeks of age and older and in cats 8 weeks of age and older.

A veterinarian or veterinary technician should demonstrate or instruct the pet owner regarding the appropriate technique for applying Revolt topically to dogs and cats prior to first use.

While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube. To administer the product, part the hair on the back of the animal at the base of the neck in front of the shoulder blades until the skin is visible. Place the tip of the tube on the skin and squeeze the tube 3 or 4 times to empty its entire contents directly onto the skin in one spot. Keeping the tube squeezed, drag it away from the liquid and lift to remove. Check the tube to ensure that it is empty.

Do not massage the product into the skin. Due to alcohol content, do not apply to broken skin. Avoid contact between the product and fingers. Do not apply when the haircoat is wet. Bathing or shampooing the dog 2 or more hours after treatment will not

reduce the effectiveness of Revolt against fleas or heartworm. Bathing or shampooing the cat 2 hours after treatment will not reduce the effectiveness of Revolt against fleas. Bathing or shampooing the cat 24 hours after treatment will not reduce the effectiveness of Revolt against heartworm.

Stiff hair, clumping of hair, hair discoloration, or a slight powdery residue may be observed at the treatment site in some animals. These effects are temporary and do not affect the safety or effectiveness of the product. Discard empty tubes in your ordinary household refuse.

### **Flea Control in Dogs and Cats**

For the prevention and control of flea infestations, Revolt should be administered at monthly intervals throughout the flea season, starting one month before fleas become active. In controlled laboratory studies >98% of fleas were killed within 36 hours. Results of clinical field studies using selamectin monthly demonstrated >90% control of flea infestations within 30 days of the first dose. Dogs and cats treated with selamectin, including those with pre-existing flea allergy dermatitis, showed improvement in clinical signs associated with fleas as a direct result of eliminating the fleas from the animals and their environment.

If the dog or cat is already infested with fleas when the first dose of Revolt is administered, adult fleas on the animal are killed and no viable fleas hatch from eggs after the first administration. However, an environmental infestation of fleas may persist for a short time after beginning treatment with Revolt because of the emergence of adult fleas from pupae.

### **Heartworm Prevention in Dogs and Cats**

For the prevention of heartworm disease, Revolt must be administered on a monthly basis. Revolt may be administered year-round or at least within one month after the animal's first exposure to mosquitoes and monthly thereafter until the end of the mosquito season. The final dose must be given within one month after the last exposure to mosquitoes. If a dose is missed and a monthly interval between dosing is exceeded then immediate administration of Revolt and resumption of monthly dosing will minimize the opportunity for the development of adult heartworms. When replacing another heartworm preventive product in a heartworm disease prevention program, the first dose of Revolt must be given within a month of the last dose of the former medication.

Selamectin, the active ingredient in Revolt, is a macrocyclic lactone compound. These compounds effectively prevent the development of adult heartworms when administered to dogs and cats within one month of exposure to infective (L3) *Dirofilaria immitis* larvae. Efficacy of macrocyclic lactones decreases below 100% in dogs, however, if first administered >2 months after exposure to infective larvae. Thus, in heartworm endemic regions, delaying initiation of heartworm prevention using Revolt beyond 2 months of first exposure to infective larvae (e.g., starting puppies and kittens at >8 weeks of age), or gaps of >2 months in the administration of Revolt during periods of heartworm transmission, increases the risk of the animal acquiring heartworms. Animals with unknown heartworm history that test negative for heartworms prior to the initiation of Revolt may be harboring pre-patent infections at the time Revolt was started. Testing such animals 3-4 months after initiation of Revolt would be necessary to confirm their negative heartworm status.

At the discretion of the veterinarian, cats  $\geq$  6 months of age may be tested to determine the presence of existing heartworm infections before beginning treatment with Revolt.

Cats already infected with adult heartworms can be given Revolt monthly to prevent further infections.

### **Ear Mite Treatment in Dogs and Cats**

For the treatment of ear mite (*O. cynotis*) infestations in dogs and cats, Revolt should be administered once as a single topical dose. A second monthly dose may be required in some dogs. Monthly use of Revolt will control any subsequent ear mite infestations. In the clinical field trials ears were not cleaned, and many animals still had debris in their ears after the second dose. Cleansing of the infested ears is recommended to remove the debris.

### **Sarcoptic Mange Treatment in Dogs**

For the treatment of sarcoptic mange (*S. scabiei*) in dogs, Revolt should be administered once as a single topical dose. A second monthly dose may be required in some dogs. Monthly use of Revolt will control any subsequent sarcoptic mange mite infestations. Because of the difficulty in finding sarcoptic mange mites on skin scrapings, effectiveness assessments also were based on resolution of clinical signs. Resolution of the pruritus associated with the mite infestations was observed in approximately 50% of the dogs 30 days after the first treatment and in approximately 90% of the dogs 30 days after the second monthly treatment.

### **Tick Control in Dogs**

For the control of tick (*Dermacentor variabilis*) infestations in dogs, Revolt should be administered on a monthly basis. In heavy tick infestations, complete efficacy may not be achieved after the first dose. In these cases, one additional dose may be administered two weeks after the previous dose, with monthly dosing continued thereafter.

### **Nematode Treatment in Cats**

For the treatment and control of intestinal hookworm (*A. tubaeforme*) and roundworm (*T. cati*) infections, Revolt should be applied once as a single topical dose.

## **WARNINGS:**

### **User Safety Warnings**

**Not for human use. Keep out of the reach of children. In humans, Revolt may be irritating to skin and eyes.**

Reactions such as hives, itching and skin redness have been reported in humans. Individuals with known hypersensitivity to Revolt should use the product with caution or consult a health care professional. Revolt contains isopropyl alcohol and the preservative butylated hydroxytoluene (BHT). Wash hands after use and wash off any product in contact with the skin immediately with soap and water. If contact with eyes occurs, then flush eyes copiously with water; if wearing contact lenses, rinse eyes first then remove contact lenses and continue to rinse for 5-10 minutes and seek medical attention. In case of ingestion by a human, contact a physician immediately. The safety data sheet (SDS) provides more detailed occupational safety information. To obtain a SDS contact Aurora Pharmaceutical at 1-888-215-1256 or [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com).

Flammable - Keep away from heat, sparks, open flames or other sources of ignition.

### **Animal Safety Warnings**

**Do not use in sick, debilitated or underweight animals (see TARGET ANIMAL SAFETY).**

## **PRECAUTIONS:**

Prior to administration of Revolt, dogs should be tested for existing heartworm infections. At the discretion of the veterinarian, infected dogs should be treated to remove adult heartworms. Revolt is not effective against adult *D. immitis* and, while the number of circulating microfilariae may decrease following treatment, Revolt is not effective for microfilariae clearance.

Hypersensitivity reactions have not been observed in dogs with patent heartworm infections administered selamectin (**See TARGET ANIMAL SAFETY**).

## **ADVERSE REACTIONS:**

### **Pre-approval clinical trials:**

Following treatment with selamectin, transient localized alopecia with or without inflammation at or near the site of application was observed in approximately 1% of 691 treated cats. Other signs observed ( $\leq 0.5\%$  of 1743 treated cats and dogs) included vomiting, loose stool or diarrhea with or without blood, anorexia, lethargy, salivation, tachypnea, and muscle tremors.

### **Post-approval experience (2021):**

The following adverse events are based on post-approval adverse drug experience reporting for selamectin. Not all adverse events are reported to FDA/CVM. It is not always possible to reliably estimate the adverse event frequency or establish a causal relationship to product exposure using this data.

The following adverse events reported for **dogs** are listed in decreasing order of reporting frequency: Lethargy, vomiting, diarrhea, anorexia, generalized pruritus, seizures, application site reactions (including alopecia, lesions, erythema, pruritis, and inflammation), tremors, ataxia, death, and dermatitis.

The following adverse events reported for **cats** are listed in decreasing order of reporting frequency: Application site reactions (including alopecia, lesions, erythema, pruritus, inflammation, vesicles, blisters, and excoriations), lethargy, anorexia, vomiting, death, generalized pruritus, diarrhea, ataxia, fever, generalized alopecia, tremors, hypersalivation, dermatitis, and seizures.

### **Contact Information:**

To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Aurora Pharmaceutical at 1-888-215-1256 or [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae)

## **TARGET ANIMAL SAFETY:**

Selamectin has been tested safe in over 100 different pure and mixed breeds of healthy dogs and over 15 different pure and mixed breeds of healthy cats, including pregnant and lactating females, breeding males and females, puppies six weeks of age and older, kittens eight weeks of age and older, and avermectin-sensitive collies. A kitten, estimated to be 5–6 weeks old (0.3 kg), died 8 1/2 hours after receiving a single treatment of selamectin at the recommended dosage. The kitten displayed clinical signs which included muscle spasms, salivation and neurological signs. The kitten was a stray with an unknown history and was malnourished and underweight (see **WARNINGS**).

**DOGS:** In safety studies, selamectin was administered at 1, 3, 5, and 10 times the recommended dose to six-week-old puppies, and no adverse reactions were observed. The safety of selamectin administered orally also was tested in case of accidental oral ingestion. Oral administration of selamectin at the recommended topical dose in 5- to 8-month-old beagles did not cause any adverse reactions. In a pre-clinical study selamectin was dosed orally to ivermectin-sensitive collies. Oral administration of 2.5, 10, and 15 mg/kg in this dose escalating study did not cause any adverse reactions; however, eight hours after receiving 5 mg/kg orally, one avermectin-sensitive collie became ataxic for several hours, but did not show any other adverse reactions after receiving subsequent doses of 10 and 15 mg/kg orally. In a topical safety study conducted with avermectin-sensitive collies at 1, 3 and 5 times the recommended dose of selamectin, salivation was observed in all treatment groups, including the vehicle control. Selamectin also was administered at 3 times the recommended dose to heartworm infected dogs, and no adverse effects were observed.

**CATS:** In safety studies, selamectin was applied at 1, 3, 5, and 10 times the recommended dose to six-week-old kittens. No adverse reactions were observed. The safety of selamectin administered orally also was tested in case of accidental oral ingestion. Oral administration of the recommended topical dose of selamectin to cats caused salivation and intermittent vomiting. Selamectin also was applied at 4 times the recommended dose to patent heartworm infected cats, and no adverse reactions were observed.

In well-controlled clinical studies, selamectin was used safely in animals receiving other frequently used veterinary products such as vaccines, anthelmintics, antiparasitics, antibiotics, steroids, collars, shampoos and dips.

## **STORAGE CONDITIONS:**

Store below 30°C (86°F).

## **HOW SUPPLIED:**

Available in eight separate dose strengths for dogs and cats of different weights (see **DOSAGE**). Revolt for puppies and kittens is available in cartons containing 3 single dose tubes. Revolt for cats and dogs is available in cartons containing 3 or 6 single dose tubes.

Approved by FDA under ANADA # 200-673

**Revolt® (selamectin)**

Manufactured By:  
Aurora Pharmaceutical, Inc.  
NORTHFIELD, MN 55057  
www.aurorapharmaceutical.com  
Aurora Pharmaceutical®  
Manufactured In U.S.A

**Client Information About Revolt® (selamectin)  
Revolt (pronounced “Re-volt”)  
Generic name: selamectin (“sel-a-mec-tin”)**

This summary contains important information about Revolt. You should read this information before you start using Revolt on your dog or cat and review it each time your prescription is refilled. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or if you want to know more about Revolt.

**What is Revolt?**

Revolt is a topical parasiticide that is applied to the skin of dogs six weeks of age and older and cats eight weeks of age and older to kill adult fleas and prevent flea eggs from hatching, prevent heartworm disease and protect your pet against other parasites (see below).

**Why has my veterinarian prescribed Revolt?**

Revolt has been prescribed by your veterinarian to treat, prevent and/or control the following parasites in your dog or cat:

**Dog Parasites:**

- Control and prevention of flea infestation (*Ctenocephalides felis*)
- Prevention of heartworm disease (*Dirofilaria immitis*)
- Treatment and control of ear mite infestation (*Otodectes cynotis*)
- Treatment and control of sarcoptic mange (*Sarcoptes scabiei*)
- Control of the American Dog Tick (*Dermacentor variabilis*)

**Cat Parasites:**

- Control and prevention of flea infestation (*Ctenocephalides felis*)
- Prevention of heartworm disease (*Dirofilaria immitis*)
- Treatment and control of ear mite infestation (*Otodectes cynotis*)
- Treatment and control of intestinal worms - Roundworm (*Toxocara cati*), Hookworm (*Ancylostoma tubaeforme*)

**What should I discuss with my veterinarian before Revolt is prescribed?**

Your veterinarian is best suited to discuss and recommend appropriate medications for your dog or cat. It is important to discuss your pet's health history with your veterinarian so he/she can decide if Revolt is right for your animal.

**Revolt should not be used in sick, debilitated or underweight animals.**

Dogs should be tested for heartworm disease prior to giving Revolt. If your dog tests

positive for adult heartworms, your veterinarian can recommend appropriate treatment. Dogs infected with adult heartworms can safely be given Revolt.

If your cat is older than six months of age, your veterinarian may decide to test him/her for heartworm disease before prescribing Revolt. Cats infected with adult heartworms can be given Revolt to prevent further infections.

### **What dose of Revolt do I use on my dog or cat?**

Your veterinarian will recommend the appropriate dose for your dog or cat based on your animal's body weight. You should not administer Revolt to dogs younger than 6 weeks of age or cats younger than 8 weeks of age. Revolt is available in eight separate dose strengths for dogs and cats of different weights.

### **What should I do if I do not give Revolt on time or miss a dose?**

If you forget to apply a monthly dose of Revolt, immediately apply Revolt, resume monthly applications, and notify your veterinarian.

### **What if I administer more than the prescribed amount of Revolt to my dog or cat?**

Contact your veterinarian if you administer more than the prescribed amount of Revolt.

### **How should Revolt be applied?**



1. Remove the Revolt tube from its protective package.

2. While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.
3. Part the hair on the back of the animal at the base of the neck, in front of the shoulder blades, until the skin is visible.
4. Apply the tip of the Revolt tube directly to the skin. Squeeze the tube firmly 3 to 4 times in one spot until empty. Keep the tube compressed on the final squeeze to avoid drawing liquid back into tube. Avoid contact between Revolt and your fingers.
5. While keeping tube squeezed, drag it away from liquid and lift up to remove.
6. Ensure tube is empty.

**Do not massage Revolt into the skin.**

**Do not apply when the haircoat is wet.**

**Do not apply to broken skin - Revolt contains alcohol.**

Stiff hair, clumping of hair, hair discoloration, or a slight powdery residue may be observed at the site in some animals. These effects are usually temporary and do not affect the safety or effectiveness of the product.

**Can I give my pet a bath after applying Revolt?**


Yes. Bathing or shampooing the dog 2 or more hours after treatment will not reduce the effectiveness of Revolt against fleas or heartworm. Bathing or shampooing the cat 2 hours after treatment will not reduce the effectiveness of Revolt against fleas. Bathing or shampooing the cat 24 hours after treatment will not reduce the effectiveness of Revolt against heartworm.

**When can I play with my pet following treatment with Revolt?**

You should avoid contact with application site when wet. You may hold or play with your pet any time after the area on which Revolt was applied is dry.

# FLEAS

*Ctenocephalides felis*



## I see fleas on my dog or cat. Is Revolt working?

Revolt kills adult fleas and prevents flea eggs from hatching. You may occasionally see a few fleas on dogs or cats treated with Revolt but more than 98% of adult fleas are killed within 36 hours.

Immature stages of the flea called pupae may be present in your pets' environment (yard, flooring, carpet, bedding, etc.). These pupae are not killed by parasiticides (including Revolt) and as such may emerge as adult fleas. These adult fleas may hop onto your pet at any time. They must be exposed to Revolt on your dog or cat before being killed. It can take from 3 to 5 weeks (or longer depending on environmental conditions) for most fleas to complete their 4-stage life cycle (egg, larvae, pupae, and adult) and reach the adult stage before being seen on your pet. Due to the presence of immature flea stages in infested environments it can take up to 2 to 3 monthly applications for Revolt to maximally control the infestation of fleas in the environment. Once the flea population is controlled you will be less likely to see fleas.

### **I see ticks on my dog. Is Revolt working?**

Revolt controls tick infestations only due to the American Dog Tick (*Dermacentor variabilis*), a tick commonly found on dogs. There are other common species of ticks that are not killed or controlled by Revolt. Your veterinarian can recommend appropriate products to control or kill ticks common to your area. For the control of the American Dog Tick, Revolt should be applied once a month; however, your veterinarian may recommend a second administration applied 14 days after the first dose if your dog has a heavy tick infestation and/or recommend additional tick control methods. It may take up to 5 days to kill the majority of ticks on your dog.

### **What are the possible side effects of Revolt?**

Like all medicines, Revolt has some side effects. The most common is hair loss at the site of application with or without inflammation (redness, flaking) in cats. Other side effects reported in cats and dogs include vomiting, diarrhea with or without blood, anorexia (decreased appetite), lethargy (sluggishness), salivation, rapid breathing, pruritus (itching), urticaria (welts, hives), erythema (skin redness), ataxia (incoordination), fever and rare instances of death. There have also been rare reports of seizures in dogs.

If you have additional questions about possible side effects, talk to your veterinarian.

### **Can Revolt be given with other medicines?**

In well-controlled clinical studies, selamectin was used safely in dogs and cats receiving other veterinary products such as vaccines, anthelmintics, antiparasitics, antibiotics, steroids, collars, shampoos and dips.

Tell your veterinarian about all medicines you have given your dog or cat in the past, and any medicines that you are planning to use with Revolt. This should include other medicines that you can get without a prescription. Your veterinarian may want to check that all of your dog's or cat's medicines can be given together.

### **How should Revolt be stored?**

Revolt is flammable - Keep away from heat, sparks, open flames or other sources of ignition. Store below 30°C (86°F). After application, empty tubes can be placed in your normal household refuse for disposal.

### **What else should I know about Revolt?**

**Revolt is not for use in humans. Revolt should be kept out of reach of children. In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin redness have been reported in humans in rare instances. Individuals with known hypersensitivity to Revolt should use the product with caution or consult a health care professional. Revolt contains isopropyl alcohol and the preservative butylated hydroxytoluene (BHT). Wash hands after use and wash off any product in contact with the skin immediately with soap and water. In case of human ingestion contact a doctor immediately.**

**Approved by FDA under ANADA #200-673**

**Revised: 08/2023**

**PRINCIPAL DISPLAY PANEL - 15 MG TUBE CARTON**

For **puppies ≤ 5 lbs** in weight and at least 6 weeks of age and **kittens ≤ 5 lbs** in weight and at least 8 weeks of age

Three single-dose tubes each containing 0.25 mL (15 mg) of selamectin

NDC 51072-097-01

0.25 mL

## Revolt® (selamectin)

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

### 3 Doses

IN 40-1577

Rev. 08/2023

REORDER NO: 17003

**Revolt® (selamectin)**

**Dose and administration:**  
Revolt is recommended for use in puppies 6 weeks of age and older and kittens 8 weeks of age and older. The recommended minimum dose is 2.2 mg of selamectin per lb (5 mg/kg) of body weight on a monthly basis.  
To facilitate application of Revolt, follow these basic steps:

- Remove one dose applicator tube from the package.
- While holding the tube in an upright position, break the seal.
- Press the cap on the tube to break the seal.
- Place the tip of the tube on the back and place tip of tube on the side of the neck.
- Squeeze tube 3-4 times in one spot to ensure the dose is compressed on the back.
- Draw the final dose to avoid drawing liquid back into tube.

**How to apply:**  
Revolt should be applied when the pet's hair coat is dry. However, bathing or immersing your puppy in water 2 hours after application will not affect the effectiveness of this treatment. A veterinarian or veterinary technician should demonstrate or instruct the pet owner regarding the appropriate technique for applying Revolt topically to dogs and cats prior to first use.  
Discard empty tubes in your ordinary household refuse.  
See package insert for full product information.

**Revolt® (selamectin)**  
Topical solution for puppies & kittens

Manufactured By:  
Aurora Pharmaceutical, Inc.  
Northfield, MN 55057  
www.aurorapharmaceutical.com

Made in the USA.

**Revolt® (selamectin)**

**3 Doses**

Topical solution for puppies ≤ 5 lbs in weight and at least 6 weeks of age and kittens ≤ 5 lbs in weight and at least 8 weeks of age

These single-dose tubes each containing 0.25 mL (15 mg) of selamectin

NDC 51072-097-01

**Revolt® (selamectin)**

Administer topically once a month to prevent **heartworm disease** and prevent and control **flea infestations**

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.  
For use in animals only. Keep out of reach of children.  
Approved by FDA under ANADA # 200-673

**Indications:**  
Revolt is recommended for use in puppies 6 weeks of age and older and kittens 8 weeks of age and older. The recommended minimum dose is 2.2 mg of selamectin per lb (5 mg/kg) of body weight on a monthly basis.  
To facilitate application of Revolt, follow these basic steps:  
Remove one dose applicator tube from the package.  
While holding the tube in an upright position, break the seal.  
Press the cap on the tube to break the seal.  
Place the tip of the tube on the back and place tip of tube on the side of the neck.  
Squeeze tube 3-4 times in one spot to ensure the dose is compressed on the back.  
Draw the final dose to avoid drawing liquid back into tube.

**How to apply:**  
Revolt should be applied when the pet's hair coat is dry. However, bathing or immersing your puppy in water 2 hours after application will not affect the effectiveness of this treatment. A veterinarian or veterinary technician should demonstrate or instruct the pet owner regarding the appropriate technique for applying Revolt topically to dogs and cats prior to first use.  
Discard empty tubes in your ordinary household refuse.  
See package insert for full product information.

**Warnings:**  
Use Safety Warnings: Keep Revolt away from reach of children. Revolt is for use in animals only. Revolt is not for use in humans. Conditions such as theft, tampering and other incidents have been reported in humans.

**Contraindications:**  
Individuals with known hypersensitivity to Revolt should use the product with caution or consult a health care professional. Revolt contains propylparaben and the preservative butylated hydroxytoluene (BHT) which may cause allergic reactions in some individuals. If contact with eyes occurs, then flush eyes copiously with water if wearing contact lenses, then the eyes first then remove contact lenses.

**Adverse Reactions:**  
In cases of irritation by a human, contact a physician immediately. The safety data sheet (SDS) provides more detailed occupational safety information. For more information on the safety of Revolt, contact Aurora Pharmaceutical at 1-888-215-1258 or www.aurorapharmaceutical.com. For additional information, consult your drug experience reporting for animal drugs, contact FDA at 1-888-DVA-4133 or online at www.fda.gov/oc/ohrtadvice. Revolt® - keep away from heat, sparks, open flames, and other source of ignition.

**Animal Safety Warnings:**  
Do not use in sick, debilitated or underweight animals. Score below 30°C (86°F).

## PRINCIPAL DISPLAY PANEL - 30 MG TUBE CARTON

For **dogs 5.1-10 lbs** in weight and at least 6 weeks of age

Three single-dose tubes each containing 0.25 mL (30 mg) of selamectin

NDC 51072-098-06

0.25 mL

# Revolt® (selamectin)

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

## 3 Doses

IN 40-1597

Rev. 02/2021

REORDER NO: 16009

**Revolt® (selamectin)**

**3 Doses**  
For dogs 5.1-10 lbs in weight and at least 6 weeks of age

NDC 51072-097-04

Three single-dose tubes each containing 0.75 mL (45 mg) of selamectin

**Revolt® (selamectin)**

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian. For use in animals only. Keep out of reach of children. Approved by FDA under ANADA # 200-673

**Dose and Administration:**  
Revolt is recommended for use in dogs six weeks of age and older. Administer 0.75 mL (45 mg) of selamectin per lb (16 mg/kg) of body weight once a month. To facilitate application of Revolt, follow these basic steps:  
• Remove one dose applicator tube from the package.  
• While holding the tube in an upright position, twist the cap to break the seal.  
• Press the base of neck and tip of tube on visible skin area.  
• Rub the tube 3-4 times in one spot until empty. Keep tube compressed on the final final rub to avoid drawing liquid back into tube.

**Parasiticide for Dogs at least 6 weeks of age**  
Revolt is recommended for use in dogs six weeks of age and older.  
• Fleas, ticks, and preventive dogs from hatching for one month.  
• Prevents and controls flea (Dipylidium immitis) infestations.  
• Prevents heartworm disease caused by *Dirofilaria immitis*.  
• Treats and controls ear mite (Otodectes cynotis) infestations.  
• Treats and controls sarcoptic mange due to *Sarcoptes scabiei*.  
• Controls *Demodex* mite infestations.

**Warnings:**  
Not for human use. Keep out of reach of children. In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin rashes have been reported in humans in rare instances.

**Manufactured By:**  
Aurora Pharmaceuticals, Inc.  
Aurora Park, MN 55007  
www.aurorapharmaceutical.com  
Manufactured in USA.

Individuals with known hypersensitivity to Revolt should use the product with caution or avoid it. A health care professional, Revolt should be used with caution. Wash hands after use and avoid any contact with the skin immediately with soap and water. If contact with eye occurs, flush eyes copiously with water. This safety data sheet (SDS) provides more detailed occupational safety information. To report a suspected adverse drug event, for technical information or to obtain a copy of the SDS, contact Aurora Pharmaceutical at 1-888-888-8888. For more information on Revolt, visit our website, contact FDA at 1-888-FDA-1088 or online at [www.fda.gov/imp/online.htm](http://www.fda.gov/imp/online.htm). **Flammable** - Keep away from heat, sparks, open flame, and other sources of ignition. Do not use in hot, disinfected or underventilated animals. (See below for CLP).

## PRINCIPAL DISPLAY PANEL - 45 MG TUBE CARTON

Topical solution for **cats 5.1-15 lbs** in weight and at least 8 weeks of age

Three single-dose tubes each containing 0.75 mL (45 mg) of selamectin

NDC 51072-097-04

0.75 mL

## Revolt® (selamectin)

Administer topically once a month for protection against **fleas** and **heartworm disease** and treatment and control of **hookworms, roundworms, and ear mites**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

### 3 Doses

IN 40-1595

Rev. 08/2023

REORDER NO: 17004

**Revolt® (selamectin)**

**3 Doses**

Topical solution for cats 5.1-15 lbs in weight and at least 8 weeks of age

Three single-dose tubes each containing 0.75 mL (45 mg) of selamectin

NDC 51072-097-04

Administer topically once a month for protection against fleas and heartworm disease and treatment and control of hookworms, roundworms, and ear mites.

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian. For use in animals only. Keep out of reach of children. Approved by FDA under ANADA # 200-673

**Revolt® (selamectin)**

**3 Doses**

Topical solution for cats 5.1-15 lbs in weight and at least 8 weeks of age

IN 40-1595  
Rev. 08/2023  
REORDER NO: 17004

**Parasite Indications for Cats at Least 8 Weeks of Age**

Indicated for the treatment and control of the following parasites in cats:

- Fleas and cat flea ova (Ctenocephalides felis felis)
- Heartworm disease (Dirofilaria immitis)
- Roundworms (Toxocara cati)
- Hookworms (Uncinaria stenocephala)
- Ear mites (Otodectes cheylei)

**How to Use:**

1. Remove one dose tube from the pack.  
2. While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.  
3. Part hair at base of neck and place tip of tube on visible skin area.  
4. Squeeze tube 3-4 times in one spot until empty. Keep tube compressed on the final squeeze to avoid drawing liquid back into tube.  
5. While holding tube, release, drag it away from liquid and lift to remove.

**Do not use:**

- In cats with known hypersensitivity to selamectin or any of the components of this product.
- In cats with known hypersensitivity to selamectin or any of the components of this product.
- In cats with known hypersensitivity to selamectin or any of the components of this product.

**Caution:**

- Avoid contact with eyes, nose, mouth, and other mucous membranes.
- Avoid contact with water, swimming, or bathing.
- Avoid contact with other animals.
- Avoid contact with food, water, or other sources of infection.

**Animal Safety:**

• Do not use in cats with known hypersensitivity to selamectin or any of the components of this product.

**Storage:**

• Store at 20°C (68°F).

**Manufactured By:**

Aurora Pharmaceutical, Inc.  
www.aurorapharm.com

Manufactured in U.S.A.

## PRINCIPAL DISPLAY PANEL - 60 MG [1 ML] TUBE CARTON

For cats 15.1-22 lbs in weight and at least 8 weeks of age

Three single-dose tubes each containing 1.0 mL (60 mg) of selamectin

NDC 51072-097-05

1.0 mL

### Revolt® (selamectin)

Administer once a month for protection against fleas and heartworm disease and treatment and control of hookworms, roundworms, and ear mites

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

### 3 Doses

IN 40-1596

Rev. 02/2021

REORDER NO: 17006

**Revolt® (selamectin)**

**3 Doses**  
For cats 15.1-22 lbs in weight and at least 8 weeks of age  
Three single-dose tubes each containing 1.0 mL (60 mg) of selamectin  
NDC 51072-087-05

**Revolt® (selamectin)**  
3 Doses  
For cats 15.1-22 lbs in weight and at least 8 weeks of age

Administer once a month for protection against fleas and heartworm disease and treatment and control of hookworms, roundworms, and ear mites.

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian. For use in animals only. Approved for use in children. Approved by FDA under ANADA # 200-673

**aurora** pharmaceutical®

**3 Doses**

**Revolt® (selamectin)**

**Indications:**  
Parasiticide for cats at least 8 weeks of age.  
Revolt is recommended for use in cats eight weeks of age and older.  
Revolt is indicated for the treatment and control of:  
• Fleas and fleas and prevents flea eggs from hatching for one month.  
• Heartworm and controls flea (Ctenocephalides felis) infestations.  
• Prevents heartworm disease caused by *Dirofilaria immitis*.  
• Treats and controls ear mite (Otodectes cynotis) infestations.  
• Treats and controls intestinal hookworm (Ancylostoma tubaeforme) and roundworm (Toxocara cati) infections.

**Warnings:**  
Not for human use. Keep out of reach of children.  
In humans, Revolt may be irritating to skin and open reactions may occur. If irritation occurs, wash the affected area with soap and water. If irritation persists, consult your veterinarian.  
Do not use in sick, debilitated or underweight animals.  
Store below 30°C (86°F).

**How to Use:**  
Remove one dose applicator tube from the package.  
While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.  
Part hair at base of neck and place tip of tube on visible skin area.  
Squeeze tube 3-4 times in one spot until the applicator tube is empty. Do not attempt to empty the tube by squeezing the tube.  
While keeping tube squeezed, drag it away from liquid and lift up to remove.

**Dose and Administration:**  
Revolt is recommended for use in cats eight weeks of age and older.  
Revolt contains 0.5 mL (60 mg) of selamectin per 1 mL (60 mg) of body weight once a month.  
To facilitate application of Revolt, follow these basic steps:  
• Remove one dose applicator tube from the package.  
• While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.  
• Part hair at base of neck and place tip of tube on visible skin area.  
• Squeeze tube 3-4 times in one spot until the applicator tube is empty. Do not attempt to empty the tube by squeezing the tube.  
• While keeping tube squeezed, drag it away from liquid and lift up to remove.

**Ensure tube is empty.**  
Revolt should be applied when the pet's haircoat is dry. However, bathing or immersing your pet in water 2 hours after application will not reduce the effectiveness of this treatment.  
Revolt is a topical treatment. Do not attempt to use Revolt by applying Revolt topically to dogs and cats prior to fleas, ticks, or other parasites.  
Discard empty tubes in your ordinary household refuse.  
See package insert for full product information.

Manufactured by:  
**aurora pharmaceutical, Inc.**  
Hennepin, MN 55857  
www.aurorapharm.com

Manufactured in USA.

### PRINCIPAL DISPLAY PANEL - 60 MG [0.5 ML] TUBE CARTON

For **dogs 10.1-20 lbs** in weight and at least 6 weeks of age

Three single-dose tubes each containing 0.5 mL (60 mg) of selamectin

NDC 51072-098-07

0.5 mL

### Revolt® (selamectin)

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

### 3 Doses

IN 40-1598

Rev. 02/2021

REORDER NO: 16011

**Revolt® (selamectin)**

**Dose and Administration:**

- Revolt is recommended for use in dogs six weeks of age and older.
- Apply 1.0 mL (120 mg) of selamectin to the skin on the back of the dog once a month.
- To facilitate application of Revolt, follow these basic steps:
  - Remove one dose applicator tube from the package.
  - While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.
  - Part the hair at base of neck and apply the tube on visible skin area.
  - Squeeze tube 3-4 times in one spot until empty. Keep tube compressed on the final squeeze to avoid drawing liquid back into tube.

**Revolt® (selamectin)**

**Indications:**

- Prevent and control flea infestations
- Prevent and control heartworm disease caused by *Dirofilara immitis*
- Treat and control ear mite (Otodectes cynotis) infestations
- Treat and control scabies (Sarcoptes scabiei) infestations
- Control Demodex mite infestations

**Warnings:**

- Not for human use. Keep out of reach of children.
- In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin rashes have been reported in humans in use studies.

**Manufacturer:**

**aurora pharmaceutical, Inc.**  
 Boothfield, MA 02007  
[www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com)

Manufactured in USA.

While keeping tube sealed, drag it away from liquid and lift up to remove.

- Ensure tube is empty.
- Revolt should be applied when the pet's haircoat is dry. However, bathing or immersing your pet in water 2 hours after application will not reduce the effectiveness of this treatment.
- Apply Revolt to the skin on the back of the dog. Do not instruct the pet owner regarding the appropriate technique for applying Revolt topically to dogs and cats prior to first use.
- Discard empty tubes in your ordinary household refuse.
- See package insert for full product information.

Individuals with known hypersensitivity to Revolt should use the product with caution or avoid it. Health care professional Revolt Hydrochloride (HCl) which binds after use and wash off of any product in contact with the skin immediately with soap and water.

Revolt is not for use in cats, dogs, or other animals.

Revolt is not for use in dogs with eye sores, skin eye sores, or with water.

Do not use in sick, debilitated or underweight animals.

Store below 30°C (86°F).

**3 Doses**  
 For dogs 10.1-20 lbs in weight and at least 6 weeks of age

**Revolt® (selamectin)**

IN 40-1599  
 Rev. 02/2021  
 REORDER NO: 16013

**3 Doses**  
 For dogs 10.1-20 lbs in weight and at least 6 weeks of age

Three single-dose tubes each containing 0.5 mL (60 mg) of selamectin

NDC 51072-098-07

**Revolt® (selamectin)**

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

**aurora pharmaceutical**

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.  
 For use in animals only. Keep out of reach of children.  
 Approved by FDA under ANADA # 200-673

## PRINCIPAL DISPLAY PANEL - 120 MG TUBE CARTON

For **dogs 20.1-40 lbs** in weight and at least 6 weeks of age

Three single-dose tubes each containing 1.0 mL (120 mg) of selamectin

NDC 51072-098-08

1.0 mL

**Revolt® (selamectin)**

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

**3 Doses**

IN 40-1599

Rev. 02/2021


REORDER NO: 16013



**Dose and Administer Instructions:**  
Revolt is recommended for use in dogs six weeks of age and older. Administer 2.0 mL (240 mg) of selamectin once a month to dogs weighing 16 to 60 mg of body weight once a month.

To facilitate application of Revolt, follow these basic steps:

- Remove one dose applicator tube from the package.
- While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.
- Part the base of the back of the dog's neck to reveal a visible skin area.
- Insert the applicator tube on the skin area.
- Apply Revolt to the skin area until empty. Keep the tube compressed until the final squeeze to avoid drawing liquid back into the tube.




**Revolt® (selamectin)**

For dogs 20.1-40 lbs in weight and at least 9 weeks of age

Manufactured by: Aurora Pharmaceutical, Inc.  
Burlington, MN 56007  
www.aurorapharmaceutical.com

Manufactured in USA.

**Parasiticide for Dogs at least 6 weeks of age**

Revolt is recommended for use in dogs six weeks of age and older.

**Indications:**

- Kill adult fleas and prevent flea eggs from hatching for one month.
- Prevent and control ticks (Dermacentor and Ixodes) infestations.
- Prevent heartworm disease caused by *Dirofilaria immitis*.
- Prevent and control ear mite (Otodectes) infestations.
- Prevent and control sarcoptic mange due to *Sarcoptes scabiei*.
- Control Demodex mite infestations.

**Warnings:**

Not for human use. Keep out of reach of children. In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin redness have been reported in humans in rare instances.

Do not use in sick, debilitated or underweight animals. Store below 30°C (86°F).

**Revolt® (selamectin)**

3 Doses  
For dogs 20.1-40 lbs in weight and at least 6 weeks of age

1.0 mL

For dogs 20.1-40 lbs in weight and at least 6 weeks of age  
Three single-dose tubes each containing 1.0 mL (120 mg) of selamectin

NDC 51072-098-09

**Revolt® (selamectin)**

3 Doses

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.  
For use in animals only. Keep out of reach of children.  
Approved by FDA under ANADA # 200-675

## PRINCIPAL DISPLAY PANEL - 240 MG TUBE CARTON

For **dogs 40.1-85 lbs** in weight and at least 6 weeks of age

Three single-dose tubes each containing 2.0 mL (240 mg) of selamectin

NDC 51072-098-09

2.0 mL

**Revolt® (selamectin)**

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-675

**3 Doses**

IN 40-1600

Rev. 02/2021

REORDER NO: 16015



**Dose and Administration:**  
Revolt is recommended for use in dogs 6 weeks of age and older. Administer once a month to prevent and control heartworm disease and prevent and control flea infestations. The recommended dosage is 10 mg (0.35 mg/lb) of selamectin once a month. For dogs 16 lb (7.3 kg) or less, use the 1.67 mL (0.06 oz) of 2.7 mg/mL (0.1 mg/lb) of selamectin once a month. To facilitate application of Revolt, follow these basic steps:

- Remove one dose applicator tube from the package.
- While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.
- Part the hair of neck and place tip of tube on visible skin area.
- Squeeze tube 3-4 times in one spot until empty. Keep tube compressed on the final squeeze to avoid drawing liquid back into tube.





While keeping tube squeezed, drag it away from liquid and fit up to remove.

Ensure tube is empty.

Revolt should be applied when the pet's haircoat is dry. However, bathing or immersing your pet in water 2 hours after application will not reduce the effectiveness of this treatment. For more information on the proper use of Revolt, please refer to the instructions on the product packaging. For more information on applying Revolt, please refer to the instructions on the product packaging. Revolt is specifically for dogs and cats prior to first use. Discard empty tubes in your ordinary household refuse. See package insert for full product information.

**Parasiticide for Dogs at least 6 weeks of age**

**Indications:**

- Kill adult fleas and prevent flea eggs from hatching for one month.
- Prevent and control heartworm disease caused by *Dirofilara immitis*.
- Prevent and control ear mite (*Otodectes cynotis*) infestations.
- Treat and control sarcoptic mange due to *Sarcoptes scabiei*.
- Control Demodex mite infestations.

**Warnings:**

Not for human use. Keep out of reach of children. In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin redness have been reported in humans in rare instances.

**Manufactured by:** **aurora pharmaceutical, Inc.**  
Northfield, MN 55057  
www.aurorapharmaceutical.com

Manufactured in USA.



For dogs **40.1-85 lbs**  
in weight and at least  
**9** weeks of age



**3 Doses**  
For dogs **40.1-85 lbs**  
in weight and at least  
**6** weeks of age

Individuals with known hypersensitivity to Revolt should use the product with caution or consult a health care professional. Revolt contains benzimidazole derivatives and may cause drowsiness, hypotension (BHT). Wash hands after use and avoid use of any product in contact with the skin immediately with soap and water. In rare instances, hypersensitivity reactions may occur. If such reactions occur, discontinue use immediately. This safety sheet (SSS) provides more detailed occupational safety information. To report a suspected adverse drug event, for technical information, or a copy of this SSS, contact the manufacturer at 1-866-3-2286 or visit our website at [www.aurorapharmaceutical.com](http://www.aurorapharmaceutical.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VEIS or online at [www.fda.gov/oc/ohrt/whistleblowers](http://www.fda.gov/oc/ohrt/whistleblowers). **Flammable** - Keep away from heat, sparks, open flame, and other sources of ignition.

Do not use in sick, debilitated or underweight animals.

Store below 30°C (86°F).



Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**



**3 Doses**

**3 Doses**  
For dogs **40.1-85 lbs**  
in weight and at least  
**6** weeks of age

For dogs **40.1-85 lbs** in weight and at least 6 weeks of age  
Three single-dose tubes each containing 3.0 mL (240 mg) of selamectin

NDC 51072-098-09



**Dose and Administration:**  
Revolt is recommended for use in dogs 6 weeks of age and older. Administer once a month to prevent and control heartworm disease and prevent and control flea infestations. The recommended dosage is 10 mg (0.35 mg/lb) of selamectin once a month. For dogs 16 lb (7.3 kg) or less, use the 1.67 mL (0.06 oz) of 2.7 mg/mL (0.1 mg/lb) of selamectin once a month. To facilitate application of Revolt, follow these basic steps:

- Remove one dose applicator tube from the package.
- While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.
- Part the hair of neck and place tip of tube on visible skin area.
- Squeeze tube 3-4 times in one spot until empty. Keep tube compressed on the final squeeze to avoid drawing liquid back into tube.





While keeping tube squeezed, drag it away from liquid and fit up to remove.

Ensure tube is empty.

Revolt should be applied when the pet's haircoat is dry. However, bathing or immersing your pet in water 2 hours after application will not reduce the effectiveness of this treatment. For more information on the proper use of Revolt, please refer to the instructions on the product packaging. For more information on applying Revolt, please refer to the instructions on the product packaging. Revolt is specifically for dogs and cats prior to first use. Discard empty tubes in your ordinary household refuse. See package insert for full product information.

**Parasiticide for Dogs at least 6 weeks of age**

**Indications:**

- Kill adult fleas and prevent flea eggs from hatching for one month.
- Prevent and control heartworm disease caused by *Dirofilara immitis*.
- Prevent and control ear mite (*Otodectes cynotis*) infestations.
- Treat and control sarcoptic mange due to *Sarcoptes scabiei*.
- Control Demodex mite infestations.

**Warnings:**

Not for human use. Keep out of reach of children. In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin redness have been reported in humans in rare instances.

**Manufactured by:** **aurora pharmaceutical, Inc.**  
Northfield, MN 55057  
www.aurorapharmaceutical.com

Manufactured in USA.

## PRINCIPAL DISPLAY PANEL - 360 MG TUBE CARTON

For **dogs 85.1-130 lbs** in weight and at least 6 weeks of age

Three single-dose tubes each containing 3.0 mL (360 mg) of selamectin

NDC 51072-098-10

3.0 mL

**Revolt® (selamectin)**

Administer once a month to prevent **heartworm disease** and prevent and control **flea infestations**

aurora pharmaceutical®

Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

For use in animals only. Keep out of reach of children.

Approved by FDA under ANADA # 200-673

**3 Doses**

IN 40-1601

Rev. 02/2021

REORDER NO: 16017



**Dose and administration:**  
Revolt is recommended for use in dogs 6 weeks of age and older. The recommended dose is 10 mg of selamectin per lb (22 mg/kg) of body weight once a month. To facilitate application of Revolt, follow these basic steps:

- Remove one dose applicator tube from the package.
- While holding the tube in an upright position, twist the cap to break the seal. The cap will remain on the tube.
- Part hair at base of neck and apply the tube on visible skin area.
- Squeeze tube 3-4 times in one spot until empty. Keep tube compressed on the final squeeze to avoid drawing liquid back into tube.



While keeping tube squeezed, drag it away from liquid and lift up to remove.

- Ensure tube is empty.
- Revolt should be applied when the pet's haircoat is dry. However, bathing or rinsing your pet's water 2 hours after application will not reduce the effectiveness of this treatment.
- For more information on the appropriate technique for instructing the pet owner regarding the appropriate technique for applying Revolt to dogs and cats, prior to first use, discard empty tube in your or your household refuse.
- See package insert for full product information.

**Parasiticide for Dogs at least 6 weeks of age**

**Indications:**

- Kills adult fleas and prevents their eggs from hatching for one month.
- Prevents and controls flea (Ceratophyllus) flea infestations.
- Prevents heartworm disease caused by *Dirofilaria immitis*.
- Treats and controls ear mite (Otodectes) infestations.
- Treats and controls sarcoptic mange due to *Sarcoptes scabiei*.
- Controls Demodex mite infestations.

**Warnings:**  
Not for human use. Keep out of reach of children. In humans, Revolt may be irritating to skin and eyes. Reactions such as hives, itching and skin rashes have been reported in humans in case reports.

**Manufactured By:**  
Zoetis Animal Health, Inc.  
Kendallville, PA 15867  
www.zoetis.com

Manufactured in USA.



**For dogs 85.1-130 lbs in weight and at least 6 weeks of age**



**3 Doses**  
For dogs 85.1-130 lbs in weight and at least 6 weeks of age



IN 491091  
REV. 02/2021  
REORDER NO: 19817

NDC 51072-097-10

For dogs 85.1-130 lbs in weight and at least 6 weeks of age  
Three single-dose tubes each containing 3.0 mL (360 mg) of selamectin



**3 Doses**

Administer once a month to prevent heartworm disease and prevent and control flea infestations



Caution: U.S. Federal law restricts this drug to use by or on the order of a licensed veterinarian. For use in animals only. Keep out of reach of children. Approved by FDA under ANADA # 200-073

REVOLT selamectin solution				
Product Information				
Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:51072-097	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
SELAMECTIN (UNII: A2669OWX9N) (SELAMECTIN - UNII:A2669OWX9N)		SELAMECTIN	60 mg in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ISOPROPYL ALCOHOL (UNII: ND2M416302)				
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51072-097-01	3 in 1 CARTON		
1		0.25 mL in 1 TUBE		
2	NDC:51072-097-02	6 in 1 CARTON		
2		0.75 mL in 1 TUBE		
3	NDC:51072-097-04	3 in 1 CARTON		
3		0.75 mL in 1 TUBE		
4	NDC:51072-097-03	6 in 1 CARTON		
4		1.0 mL in 1 TUBE		

5	NDC:51072-097-05	3 in 1 CARTON		
5		1.0 mL in 1 TUBE		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA	ANADA200673		08/26/2020	

## REVOLT

selamectin solution

<b>Product Information</b>			
<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:51072-098
<b>Route of Administration</b>	TOPICAL		

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
SELAMECTIN (UNII: A2669OWX9N) (SELAMECTIN - UNII:A2669OWX9N)	SELAMECTIN	120 mg in 1 mL

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:51072-098-01	6 in 1 CARTON		
1		0.25 mL in 1 TUBE		
2	NDC:51072-098-06	3 in 1 CARTON		
2		0.25 mL in 1 TUBE		
3	NDC:51072-098-02	6 in 1 CARTON		
3		0.5 mL in 1 TUBE		
4	NDC:51072-098-07	3 in 1 CARTON		
4		0.5 mL in 1 TUBE		
5	NDC:51072-098-03	6 in 1 CARTON		
5		1.0 mL in 1 TUBE		
6	NDC:51072-098-08	3 in 1 CARTON		
6		1.0 mL in 1 TUBE		
7	NDC:51072-098-04	6 in 1 CARTON		
7		2.0 mL in 1 TUBE		
8	NDC:51072-098-09	3 in 1 CARTON		
8		2.0 mL in 1 TUBE		

<b>9</b>	NDC:51072-098-05	6 in 1 CARTON		
<b>9</b>		3.0 mL in 1 TUBE		
<b>10</b>	NDC:51072-098-10	3 in 1 CARTON		
<b>10</b>		3.0 mL in 1 TUBE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200673	08/26/2020	

**Labeler** - Aurora Pharmaceutical, Inc. (832848639)

## Establishment

Name	Address	ID/FEI	Business Operations
Aurora Pharmaceutical, Inc.		832848639	manufacture

Revised: 6/2025

Aurora Pharmaceutical, Inc.