

**UNISHIELD ASPIRIN- aspirin tablet, coated**  
**UNISHIELD ASPIRIN- aspirin tablet, film coated**  
**Unishield**

-----  
**Unishield Aspirin**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or to any other pain reliever/fever reducer

***Ask a doctor before use if***

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- drink a full glass of water with each dose

**Adults and children 12 years and over:** Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

Children under 12 years: Consult a doctor

### Other information

- store at room temperature
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

### Inactive ingredients

corn starch, hypromellose\*, microcrystalline cellulose\*, polyethylene glycol\* povidone\*, propylene glycol\*

\* may contain

Questions or comments? 800-480-5855

## Unishield Aspirin Label

Unishield

Aspirin 325 mg

Pain Reliever/Fever Reducer (NSAID)

100 Tablets/2 Tablets Per Packet

Pull To Open

This Package is for Households Without Young Children.

- Relieves headaches, muscle pain and minor arthritis pain

Compare active ingredient to:

Genuine Bayer®

Registered Trademark of Bayer Corp.

Tamper Evident Unit Dose Packets

**Drug Facts**  
**Active ingredient (in each tablet) Purpose**  
Aspirin 325 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves  
■ headache ■ muscle pain  
■ minor pain of arthritis ■ pain and fever of colds  
■ toothache ■ menstrual pain

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling  
■ asthma (wheezing) ■ shock

**Drug Facts (continued)**  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcohol drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or to any other pain reliever/fever reducer.

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems such as heartburn  
■ you have high blood pressure, heart disease, liver problems, or kidney disease  
■ you are taking a diuretic  
■ you have asthma (continued on opposite panel)

**Compare active ingredient to:**  
Genuine Bayer®  
Registered Trademark of Bayer Corp.

**Drug Facts (continued)**  
**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better

■ pain gets worse or lasts more than 10 days  
■ fever lasts more than 3 days  
■ ringing in the ears or loss of hearing occurs  
■ redness or swelling is present  
■ new symptoms appear

**Directions**  
■ drink a full glass of water with each dose

**Adults and children (12 years and over)** Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.

**Children under 12 years:** Consult a doctor

**Other information**  
■ store at room temperature  
■ avoid excessive heat and humidity  
■ tamper evident sealed packets  
■ do not use any opened or torn packets

**Inactive ingredients**  
corn starch, hypromellose\*, microcrystalline cellulose\*, polyethylene glycol\*, povidone\*, propylene glycol\*

\*may contain

**Questions or comments?** 800-480-5855

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

Retain carton for complete product information

Retain carton for complete product information

Reorder #11133  
Packaged for UniShield  
San Fernando, CA 91340 USA  
1(800) 480-5855  
www.unishieldusa.com

## Unishield Aspirin Label

Aspirin 325 mg

# Pain Reliever/Fever Reducer (NSAID)

100 Tablets/2Tablets Per Packet

Pull To Open

This Package is for Households Without Young Children.

- Relieves headaches, muscle pain and minor arthritis pain

Compare active ingredient to:

Genuine Bayer®

Registered Trademark of Bayer Corp.

Tamper Evident Unit Dose Packets

**Drug Facts**  
**Active ingredient (in each tablet)** Purpose  
 Aspirin 325 mg (NSAID)\*...Pain reliever/fever reducer  
 \*nonsteroidal anti-inflammatory drug

**Uses**  
 temporarily relieves  
 ■ headache ■ muscle pain  
 ■ minor pain of arthritis ■ pain and fever of colds  
 ■ toothache ■ menstrual pain

**Warnings**  
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
 ■ hives ■ facial swelling  
 ■ asthma (wheezing) ■ shock

**Drug Facts (continued)**  
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcohol drinks every day while using this product.  
 ■ take more or for a longer time than directed

**Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer.**  
 Ask a doctor before use if:  
 ■ stomach bleeding warning applies to you  
 ■ you have a history of stomach problems such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
 ■ you are taking a diuretic  
 ■ you have asthma (continued on opposite panel)

**Directions**  
 ■ drink a full glass of water with each dose  
**Adults and children (12 years and over)** Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.

**Children under 12 years:** Consult a doctor

**Other information**  
 ■ store at room temperature  
 ■ avoid excessive heat and humidity  
 ■ tamper evident sealed packets  
 ■ do not use any opened or torn packets

**Inactive ingredients**  
 corn starch, hydroxyethylcellulose, microcrystalline cellulose, polyethylene glycol, povidone, propylene glycol  
 \*may contain

**Questions or comments?** 800-485-5855

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

Relieves headaches, muscle pain and minor arthritis pain

Compare active ingredient to: **Bayer** Registered trademark of Bayer Corp.

Retain carton for complete product information

UNISHIELD ASPIRIN			
aspirin tablet, coated			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49314-1173
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
Inactive Ingredients			
Ingredient Name			Strength
POVIDONE (UNII: FZ 989GH94E)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
STARCH, CORN (UNII: O8232NY3SJ)			
Product Characteristics			
<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm

<b>Flavor</b>		<b>Imprint Code</b>	FR21	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49314-1173-3	50 in 1 BOX	03/01/2021	04/01/2024
1		2 in 1 PACKET; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	03/01/2021	04/01/2024	

<b>UNISHIELD ASPIRIN</b>			
aspirin tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49314-1163
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name	Basis of Strength	Strength	
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
<b>Inactive Ingredients</b>			
Ingredient Name	Strength		
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)			
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)			
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)			
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)			
<b>Product Characteristics</b>			
<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49314-1163-3	50 in 1 BOX	03/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/01/2021	

**Labeler** - Unishield (790677053)

**Registrant** - Unifirst First Aid Corporation (832947092)

## Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(49314-1163, 49314-1173)

Revised: 9/2025

Unishield