BENZO-JEL- benzocaine gel Henry Schein Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200 mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep Out of Reach of Children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59°-86°F (15-30°C). Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

1-800-472-4346

Distributed by (in US only): HENRY SCHEIN INC. 135 DURYEA ROAD MELVILLE, NY 11747 USA

For Professional Use Only

Made in USA



mouth symptoms do not improve if swelling, rash or fever develops, redness persists or worsens; or directed by a physician. If sore in 7 days; irritation, pain, or see your physician promptly

 in children under 2 years of age When using this product for teething

contact occurs, flush with water. Avoid contact with eyes. If

> may cause methemoglobinemia, a serious condition that must be

warning: Use of this product

Methemoglobinemia

Warnings

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed,

If pregnant or breast feeding (eep out of reach of children. Poison Control Center right away. get medical help or contact a ask a physician before use.

 Apply only amount needed to Directions

attention if you or a child in your

pale, gray or blue colored skin

care develops:

 children under 2 years of age: the oral mucosa to prevent or relieve pain.

do not use

Store at 59° - 86°F (15 - 30°C). Other information Protect from freezing.

dizziness or lightheadedness

shortness of breath · rapid heart rate

 headache (cyanosis)

flavoring, PEG 3350, PEG 400, sodium saccharin, water. May **Inactive** ingredients

yellow #5, (tartrazine), yellow #6, #5, red #3, red #28, red #40, as a color additive.

contain blue #1, green #3, green

Questions or comments?

-800-472-4346

99093451 Rev 2018/06

Allergy Alert: Do not use on allergies to local anesthetics such as procaine, butacaine, fatique or lack of energy patients with a history of

Do not use anesthetics.

benzocaine or other "caine'

for more than 7 days unless

BENZO-JEL

Benzocaine 200 mg (in each g)

Active ingredients

Drug Facts

benzocaine gel

Product Information

HUMAN OTC DRUG NDC:0404-0036 **Product Type** Item Code (Source)

occur even if you have used this

product before. Stop use and

seek immediate medical

reduces the amount of oxygen

treated promptly because it

carried in the blood. This can

DENTAL Route of Administration

For oral mucosal use only, as

Oral Anesthetic

Purpose

emporary relief of pain due

directed by dentist. For the

o minor dental procedures.

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZ OCAINE** 200 mg in 1 g

Inactive Ingredients

Ingredient Name Strength POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)

POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)

SACCHARIN SODIUM (UNII: SB8ZUX40TY)

WATER (UNII: 059QF0KO0R)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics			
Color	pink	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0404-0036- 30	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018	

Marketing In	Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph not final	part356	05/01/2016			

Labeler - Henry Schein Inc. (012430880)

Revised: 1/2022 Henry Schein Inc.