

SALICYLIC ACID- one-step corn remover patch
Premier Brands of America Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Premier Brands One-Step Corn Removers Tan

Active ingredient

Salicylic acid 40%w/w

Purpose

Corn remover

Use

- for the removal of corns

Warnings

For external use only.

Do not use

- if you are a diabetic
- have poor blood circulation
- on irritated skin or any area that is infected or reddened

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

If discomfort persists see your doctor or podiatrist.

Directions

- wash affected area: may soak corn in warm water for 5 minutes
- dry area thoroughly
- remove bandage strips by peeling backing off at slit
- carefully adhere bandage with the medicated portion directly over corn
- after 48 hours remove medicated bandage strip
- repeat this procedure every 48 hours as needed (until corn is removed) for up to 14 days

Other information

store between 59° to 86°F (15° to 30°C)

Inactive ingredients

acrylic adhesive, brown iron oxide, polyethylene foam

Questions?

call 1-866-964-0939

Principal Display Panel

Premier

Medicated

ONE-STEP CORN

REMOVERS

Salicylic Acid

Effective corn removal treatment

Medicated bandage conceals & protects

One-step, no mess application

Tan Bandage

8 Medicated Bandages



Drug Facts

Active ingredient Salicylic acid 40% w/w.....	PurposeCorn remover
Use ■ for the removal of corns	
Warnings For external use only.	
Do not use ■ if you are diabetic ■ have poor blood circulation ■ on irritated skin or any area that is infected or reddened	
Keep out of reach of children If swallowed, get medical help or contact a Poison Control Center right away. If discomfort persists see your doctor or podiatrist.	
Directions ■ wash affected area, may soak corn in warm water for 5 minutes ■ dry area thoroughly ■ remove bandage strips by peeling backing off at slit ■ carefully adhere bandage with the medicated portion directly over corn ■ after 48 hours remove medicated bandage strip ■ repeat this procedure every 48 hours as needed (until corn is removed) for up to 14 days	
Other information store between 59° and 86°F (15° and 30°C)	
Inactive ingredients acrylic adhesive, brown iron oxide, polyethylene foam	
Questions? Call 1-866-964-0939	

Care ■ Avoid surrounding skin when applying product

Distributed by:
©2021 Premier Brands of America Inc.
White Plains, New York 10601
www.premier-brands.com
Questions? Call 1-866-964-0939
MADE IN USA WITH U.S.
AND IMPORTED PARTS

SALICYLIC ACID

one-step corn remover patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56104-047
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	40 mg in 6

Inactive Ingredients

Ingredient Name			Strength	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)				
BROWN IRON OXIDE (UNII: 1N032N7MFO)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56104-047-08	8 in 1 BOX; Type 0: Not a Combination Product	02/15/2021	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part358F	02/15/2021		

Labeler - Premier Brands of America Inc. (080051232)

Revised: 2/2021

Premier Brands of America Inc.