

RINGL- acetaminophen tablet
Sato Pharmaceutical Co., Ltd.

Ringl Tablet (final packaged product)

Active ingredient (in each tablet)
Acetaminophen 325 mg

Purpose Pain reliever-fever reducer

Uses

- for the temporary relief of minor aches and pains associated with:
 - a cold
 - sore throat
 - headache
 - toothache
 - muscular aches
 - backache
 - the premenstrual and menstrual periods (dysmenorrhea)
- for minor pain from arthritis
- to reduce fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 12 tablets in 24 hours, which is the maximum daily amount for this product
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is

- taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain persists for more than 10 days (for adults) or 5 days (for children)
- sore throat persists for more than 2 days
- any of the following occurs (these could be signs of a serious condition):
 - fever persists for more than 3 days
 - a severe sore throat
 - sore throat is accompanied or followed by fever, headache, rash, nausea, or vomiting
- pain or fever persists or gets worse
- new symptoms occur
- redness or swelling is present
- when using for pain or arthritis:
 - pain persists for more than 10 days
 - redness is present
 - in conditions affecting children under 12 years of age

Do not give to children under 3 years of age or to children for the pain of arthritis.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over: 1 to 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses or 12 tablets in 24 hours, or as directed by a doctor

children 11 to under 12 years of age: 1 to 1.5 tablets every 4 hours, while symptoms persist, not to exceed 5 doses or 7.5 tablets in 24 hours, or as directed by a doctor

children 9 to under 11 years of age: 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses or 5 tablets in 24 hours, or as directed by a doctor

children 6 to under 9 years of age: 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses or 5 tablets in 24 hours, or as directed by a doctor

children 4 to under 6 years of age: 3/4 tablet every 4 hours, while symptoms persist, not to exceed 5 doses or 3.75 tablets in 24 hours, or as directed by a doctor

children 3 to under 4 years of age: 1/2 tablet every 4 hours, while symptoms persist, not to exceed 5 doses or 2.5 tablets in 24 hours, or as directed by a doctor

children under 3 years of age: ask a doctor

Other information

■ keep tightly sealed ■ store between 15° to 30°C (59° to 86° F)

Inactive ingredients hydroxypropyl cellulose, microcrystalline cellulose, magnesium stearate, and polyvinyl alcohol.

★ Do not use if printed bottle neck seal is broken or missing. TO OPEN: See the top of cap. ★ 瓶口の密封シールが破れていたり、ついでにふたがなかった場合は使用しないでください。開け方: キャップをグッと押し下げ、たまま、矢印の方向にまわしてください。



Drug Facts

Active ingredient (in each tablet) Purpose
Acetaminophen 325 mg Pain reliever-fever reducer

Uses
■ for the temporary relief of minor aches and pains associated with: ■ a cold ■ sore throat ■ headache ■ toothache ■ muscular aches ■ backache ■ the premenstrual and menstrual periods (dysmenorrhea) ■ for minor pain from arthritis ■ to reduce fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
■ adult takes more than 12 tablets in 24 hours, which is the maximum daily amount for this product
■ child takes more than 5 doses in 24 hours
■ taken with other drugs containing acetaminophen
■ adult has 3 or more alcoholic drinks everyday while using this product

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has liver disease
Ask a doctor or pharmacist before use if the user is
■ taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain persists for more than 10 days (for adults) or 5 days (for children)
■ sore throat persists for more than 2 days
■ any of the following occurs (these could be signs of a serious condition):
■ fever persists for more than 3 days ■ a severe sore throat ■ sore throat is accompanied or followed by fever, headache, rash, nausea, or vomiting ■ pain or fever persists or gets worse ■ new symptoms occur ■ redness or swelling is present

Drug Facts (continued)

■ when using for pain of arthritis:
■ pain persists for more than 10 days ■ redness is present ■ in conditions affecting children under 12 years of age

Do not give to children under 3 years of age or to children for the pain of arthritis.
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over	1 to 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses or 12 tablets in 24 hours, or as directed by a doctor
children 11 to under 12 years of age	1 to 1.5 tablets every 4 hours, while symptoms persist, not to exceed 5 doses or 7.5 tablets in 24 hours, or as directed by a doctor
children 9 to under 11 years of age	1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses or 5 tablets in 24 hours, or as directed by a doctor
children 6 to under 9 years of age	1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses or 5 tablets in 24 hours, or as directed by a doctor
children 4 to under 6 years of age	3/4 tablet every 4 hours, while symptoms persist, not to exceed 4 doses or 3.75 tablets in 24 hours, or as directed by a doctor
children 3 to under 4 years of age	1/2 tablet every 4 hours, while symptoms persist, not to exceed 3 doses or 2.5 tablets in 24 hours, or as directed by a doctor
children under 3 years of age	ask a doctor

Other information
■ keep tightly sealed ■ store between 15° to 30°C (59° to 86° F)
Inactive ingredients hydroxypropyl cellulose, microcrystalline cellulose, magnesium stearate, and polyvinyl alcohol

Pain Reliever Fever Reducer 鎮痛・解熱
RINGL 錠剤
Acetaminophen アセトアミノフェン
CHILD-RESISTANT PACKAGING
錠剤キップ使用
30 TABLETS (錠)
See new warnings information
使用上の注意が新しくなりました

SATO NDC 49873 08201

医薬品表示

有効成分(1錠当たり) 用途
アセトアミノフェン 325mg 鎮痛・解熱剤

効能
■ 下記の症状に伴う軽度から中等度の鎮痛:
・発熱(頭痛、腰痛、歯痛、関節痛、1年病)
・風邪・インフルエンザによる発熱(熱さまし)
■ 関節炎による軽度から中等度の鎮痛
■ 解熱

注意
肝臓に障害する警告: 本製品はアセトアミノフェンを含有しています。この場合、過量・頻回服用を避ける必要があります。
■ 成人が24時間以内の本製品の最大日量、10錠を5回服用する
■ 小児が24時間以内の本製品の最大日量、7.5錠を5回服用する
■ アセトアミノフェンを含有する他の製剤と併用する
■ 成人が本製品の処方中にアルコール飲料を併用しないよう注意する。
■ アセトアミノフェンを含有する他の薬品(処方せん薬、非処方せん薬)を併用する。薬剤師がアセトアミノフェンを含有しているかどうかを知らせる場合は、医師または薬剤師に相談してください。
■ 次の場合は併用前に医師に相談してください。
■ 肝臓病がある場合は併用前に医師に相談してください。
■ 腎臓病がある場合は併用前に医師に相談してください。
■ 糖尿病がある場合は併用前に医師に相談してください。
■ 次のいずれかの症状がある場合は併用前に医師に相談してください。
・発熱が3日以上続く。・発熱が繰り返す。・発熱が3日以上続く。
・発熱が繰り返す。・発熱が繰り返す。・発熱が繰り返す。
■ 併用前に医師に相談してください。
■ 併用前に医師に相談してください。
■ 併用前に医師に相談してください。

用法・用量
成人の小児 1錠を4時間おきに1回(1日最大3回、24時間以内の最大日量10錠)服用してください。また併用前に医師に相談してください。
小児の小児 1.5錠を4時間おきに1回(1日最大5回、24時間以内の最大日量7.5錠)服用してください。また併用前に医師に相談してください。
小児の小児 1錠を4時間おきに1回(1日最大5回、24時間以内の最大日量5錠)服用してください。また併用前に医師に相談してください。
小児の小児 1錠を4時間おきに1回(1日最大5回、24時間以内の最大日量5錠)服用してください。また併用前に医師に相談してください。
小児の小児 1錠を4時間おきに1回(1日最大5回、24時間以内の最大日量5錠)服用してください。また併用前に医師に相談してください。
小児の小児 1錠を4時間おきに1回(1日最大5回、24時間以内の最大日量5錠)服用してください。また併用前に医師に相談してください。

その他の情報
■ 室温(15°C-30°C)で保管してください。
■ 15°C-30°C(59°F-86°F)で保管してください。

添加物
ヒドロキシプロピルセルロース、結晶セルロース、ステアリン酸マグネシウム、ポリビニルアルコール

Reports of serious side effects associated with use of this product can be sent to:
SATO PHARMACEUTICAL, INC.
20695 S. Western Ave., Suite 240, Torrance, CA 90501
Manufactured by
SATO PHARMACEUTICAL CO., LTD.
1-5-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN
製造元 佐藤薬業株式会社
日本医薬品動向協会 17号5番2号

EXP DATE 使用期限
LOT 製造記号

US-U

RINGL
acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49873-082
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	

Inactive Ingredients

Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE, LOW SUBSTITUTED (UNII: 2165RE0K14)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	SATO;N
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49873-082-01	1 in 1 CARTON	05/09/1996	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/09/1996	

Labeler - Sato Pharmaceutical Co., Ltd. (690575642)

Establishment

Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-082) , label(49873-082) , pack(49873-082)

Revised: 12/2023

Sato Pharmaceutical Co., Ltd.