STOMACH RELIEF- bismuth subsalicylate tablet, chewable SAFEWAY

Stomach Relief Bismuth Subsalicylate 262 mg Chewable Tablets 30 Counts Drug Facts

Active ingredient (in each chewable tablet)

Bismuth subsalicylate 262 mg

Purpose

Antidiarrheal/ upset stomach reliever

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Do not use if you are

- allergic to salicylate (including aspirin)
- taking other salicylate products

Do not use if you have

an ulcer

- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately at 1-800-222-1222

Directions

- chew completely or dissolve tablets in mouth. Do not swallow tablets whole.
- adults and children 12 years and over:
- 2 tablets (1 dose) every ½ hour to 1 hour or 4 tablets (2 doses) every hour as needed for diarrhea
- 2 tablets (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- children under 12 years: ask a doctor

Other information

- each tablet contains: salicylate 102 mg
- Store at room temperature
- avoid excessive heat
- keep this carton for future reference on full labeling

Inactive ingredients

D&C Red No. 27, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, sorbitol, sucralose.

Questions or comments?

1-866-467-2748

Principal Display Panel

Compare to the active ingredient in Pepto-Bismol® Chewables*

Stomach Relief

Chewable Tablets

Upset Stomach Reliever/Antidiarrheal

Original flavor

5 Symptoms Digestive Relief:

- Heartburn
- Indigestion
- Nausea
- Upset Stomach
- Diarrhea

30 Chewable Tablets

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR OF STRIP UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Procter & Gamble Inc., the distributor of Pepto-Bismol® Chewables.

Distributed by:



STOMACH RELIEF

bismuth subsalicylate tablet, chewable

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-241
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg	

Inactive Ingredients

Strength

Product Characteristics			
Color	PINK (light-pink)	Score	no score
Shape	ROUND	Size	16mm
Flavor		Imprint Code	RP124
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-241- 30	5 in 1 CARTON	10/01/2020	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/01/2020	

Labeler - SAFEWAY (009137209)

Revised: 11/2024 SAFEWAY