

**DIRECT SAFETY ASPIRIN- aspirin tablet, coated**  
**DIRECT SAFETY ASPIRIN- aspirin tablet, film coated**  
**Direct Safety**

-----  
**Direct Safety Aspirin**

***Drug Facts***

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose



Direct Safety®

Aspirin

Pain Reliever

Aspirin 325 mg (NSAID)

Aspirina 325 mg (AINE)

This Package is for Households without Young Children.

Este Paquete Es Hogares Sin Niños Pequeños

Industrial Pack 500

(250 Packets of 2 Tablets)

# ASPIRIN

## Pain Reliever

Aspirin 325 mg (NSAID)  
Aspirina 325 mg (AINE)

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Aspirin 325 mg (NSAID) Pain reliever/fever reducer</p> <p><b>Uses</b> temporarily relieves ● headache ● muscle pain ● minor pain of arthritis ● pain and fever of colds ● toothache ● menstrual pain</p> <p><b>Warnings</b> Highly sensitive: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Warnings</b> Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include ● hives ● facial swelling ● asthma (wheezing) ● shock If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinner (anticoagulant or blood thinner) ● take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 2 or more alcohol drinks every day while using this product ● take more for a longer time than directed.</p> <p><b>Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer.</b></p> <p><b>Ask a doctor before use if</b> ● stomach bleeding warning applies to you ● you have a history of stomach problems such as heartburn ● you have high blood pressure, heart disease, liver problems, or kidney disease ● you are taking a blood thinner ● you have asthma</p> <p><b>Ask a doctor or pharmacist before use if you are taking a prescription drug for</b> ● gout ● diabetes ● arthritis</p>	<p><b>Drug Facts (continued)</b></p> <p>How to use: Ask your doctor if ● you have an allergic reaction occurs. Seek medical help right away. ● you experience any of the following signs of stomach bleeding: ● hot heart ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● pain gets worse or lasts more than 10 days ● lasts less than 3 days ● if ringing in the ears or loss of hearing occurs ● if hives or swelling is present ● new symptoms appear</p> <p><b>If pregnant or breast feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> ● Drink a full glass of water with each dose.</p> <p><b>Adults and children (12 years and older)</b> ● Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours. Not to exceed 12 tablets in 24 hours.</p> <p><b>Children under 12 years:</b> ● Consult a doctor.</p> <p><b>Other information</b> ● Store at room temperature. ● Avoid exposure to heat and humidity. ● Keep in original sealed packets. ● Do not use any opened or torn packets.</p> <p><b>Inactive ingredients</b> corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, polypropylene glycol, "may contain"</p> <p><b>Questions or comments? 1-800-356-9100</b></p>	<p><b>Información del medicamento</b></p> <p><b>Ingrediente activo (en cada tableta)</b> Aspirina 325 mg (AINE) Analgésico antiinflamatorio no esteroideo</p> <p><b>Propiedad</b> Alivia el dolor/Reduce la fiebre</p> <p><b>Usos:</b> alivia temporal de ● dolor de cabeza ● dolores musculares ● dolor y fiebre de resaca ● dolor menstrual ● dolor de dientes ● dolor de oídos ● dolor de garganta</p> <p><b>Advertencias:</b> El síndrome de Reye: No debe administrarse a niños y adolescentes enfermos o en recuperación de los síntomas de la varicela o de la influenza o de la gripe. Si con el consumo de este medicamento aparecen cambios en el comportamiento de la persona junto con náuseas y vómitos, consulte con un doctor ya que estos síntomas pueden señalar el comienzo del síndrome de Reye, una enfermedad que siempre es poco común en la gripe.</p> <p><b>Atención a las alergias:</b> La aspirina puede causar severas reacciones alérgicas las cuales pueden incluir: ● urticaria ● inflamación facial ● asma (respiración con silbido) ● shock Si se presenta una reacción alérgica, de inmediato suspenda su consumo y obtenga atención médica.</p> <p><b>Advertencia acerca del sangrado estomacal:</b> Este producto contiene un medicamento antiinflamatorio no esteroideo (AINE), el cual pudiera causar sangrado estomacal. Las posibilidades de que ocurra este suceso son: ● si tiene 60 o más años de edad ● si ha tenido úlceras estomacales o problemas de sangrado ● si toma medicamentos para el adelgazamiento de la sangre (anticoagulantes) o esteroides ● si toma otros medicamentos con su receta médica que contengan un AINE (aspirina, naproxeno, ibuprofeno, u otros) ● si toma 2 o más bebidas alcohólicas diariamente mientras consume este producto ● si se consume más o por más tiempo de lo indicado</p> <p><b>no se administre si usted es alérgico a la aspirina o a cualquier otro analgésico/reductor de fiebre.</b></p> <p><b>Consulte con un médico antes de consumir, si</b> ● la advertencia acerca del sangrado estomacal se aplica en su caso. ● tiene antecedentes de problemas estomacales tales como acidez estomacal ● toma la presión sanguínea alta, enfermedad del corazón, anemia del hígado o enfermedad de los riñones ● está tomando un estudio de presión de sangre</p>	<p><b>Información del medicamento (continuación)</b></p> <p>Consulte con un médico o farmacéutico antes de administrarlo si usted está tomando un medicamento para: ● la gripe ● diabetes ● la artritis</p> <p><b>Deje de tomar este producto y consulte con el médico si</b> ● ocurre una reacción alérgica. Busque ayuda médica inmediatamente. ● se presentan cualquiera de los siguientes síntomas de sangrado estomacal: ● se siente débil ● vomita sangre ● si los excrementos contienen sangre o tienen un aspecto negro ● tiene dolor estomacal que no mejora ● el dolor empeora o persiste por más de 10 días ● si tiene pérdida por más de 3 días ● ocurre un zumbido en los oídos o pérdida de la audición ● aparece un eritema facial</p> <p><b>Si está embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto. Es sobre todo importante que no se consuma aspirina durante los últimos 3 meses del embarazo, a menos que el médico se lo haya especificado, ya que esto podría causar problemas a la criatura que está por nacer o traer complicaciones durante el parto.</b></p> <p><b>Monitoreo en y todos los medicamentos hacen del alcance de los riñones. En caso de ocurrir una sobredosis obtenga inmediata atención médica o llame a un centro de control de envenenamiento (Poison Control Center).</b></p> <p><b>Indicaciones</b> ● con cada dosis debe un vaso lleno con agua</p> <p><b>Aplicación y dosis (ver "Tiene 12 tablets/Children 3 tablets cada 6 horas, no exceda 12 tablets en 24 horas. Niños menores de 12 años Consultar al médico")</b></p> <p><b>Otra información</b> ● almacenamiento: conserve a temperatura ambiente ● evite exposición al calor y humedad excesiva ● papeles con sellado evidente ● no use ningún paquete abierto o roto</p> <p><b>Ingredientes inactivos</b> amidon de maíz, "hipromelosa", celulosa microcristalina, "polietileno glicol", "propileno glicol", "povidona"</p> <p><b>¿Tiene alguna pregunta o comentario? Llame al 1-800-356-9100</b></p>
---	--	--	--

Retain carton for complete product information

**ASPIRIN, PAIN RELIEVER (NSAID)**

Order No. 23578  
Reorder: 800-356-9100  
cousinc.com

DISTRIBUTED BY:  
COUNNER SAFETY PRODUCTS  
MADISON, WI 53744-4190

Industrial Pack 500  
(250 Packets of 2 Tablets)

PAQUETES CON SELLADO EVIDENTE.  
NO USE NINGÚN PAQUETE  
ABIERTO O ROTO.

Order No. 23578  
Reorder: 800-356-9100  
cousinc.com

DISTRIBUIDO POR:  
COUNNER SAFETY PRODUCTS  
MADISON, WI 53744-4190

<b>DIRECT SAFETY ASPIRIN</b>			
aspirin tablet, coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:61245-620
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
<b>Inactive Ingredients</b>			
<b>Ingredient Name</b>	<b>Strength</b>		
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			
POVIDONE (UNII: FZ989GH94E)			
STARCH, CORN (UNII: O8232NY3S)			

## Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61245-620-13	250 in 1 BOX	02/15/2021	04/01/2024
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2021	04/01/2024

## DIRECT SAFETY ASPIRIN

aspirin tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61245-616
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

## Product Characteristics

Color	white	Score	no score
-------	-------	-------	----------

<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61245-616-13	250 in 1 BOX	02/15/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2021	

**Labeler** - Direct Safety (023316151)

**Registrant** - Unifirst First Aid Corporation (832947092)

### Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(61245-616, 61245-620)

Revised: 9/2025

Direct Safety