

**DIRECT SAFETY ASPIRIN- aspirin tablet, coated**  
**DIRECT SAFETY ASPIRIN- aspirin tablet, film coated**  
**Direct Safety**

-----  
**Direct Safety Aspirin**

***Drug Facts***

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever lasts more than 3 days
  - if ringing in the ears or loss of hearing occurs
  - redness or swelling is present
  - new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose

**Adults and children 12 years and over:** take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

**Children under 12 years:** consult a doctor

**Other information**

- store at room temperature
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

corn starch, hypromellose\*, microcrystalline cellulose\*, polyethylene glycol\* povidone\*, propylene glycol\*

\* may contain

**Questions or comments? 1-800-356-9100**

Direct Safety®

Aspirin

Pain Reliever

Aspirin 325 mg (NSAID)

Aspirina 325 mg (AINE)

This Package is for Households without Young Children.

Este Paquete Es Hogares Sin Niños Pequeños

Industrial Pack 500

(250 Packets of 2 Tablets)

# ASPIRIN

## Pain Reliever

**Aspirin 325 mg (NSAID)**  
**Aspirina 325 mg (AINE)**

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)  
Pain reliever/fever reducer  
Nonsteroidal anti-inflammatory drug

**Uses**  
Temporarily relieves:  
■ headache ■ muscle pain  
■ pain and fever of colds ■ toothache ■ minor pain of arthritis

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Always "shut" Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
In allergic reaction victims, stop use and seek medical help right away.  
Research bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood-thinning (anticoagulant or blood-thin) drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 2 or more alcohol drinks every day while using this product  
■ take more or for a longer time than directed  
Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer.  
Ask a doctor before use if:  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems such as heartburn  
■ you have high blood pressure, heart disease, liver ulcers, or kidney disease  
■ you are taking a dietic  
■ you have asthma.  
Ask a doctor or pharmacist before use if you are taking a prescription drug for:  
■ pain ■ diabetes ■ asthma

**Drug Facts (continued)**

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ hot heart, ■ vomit blood, ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you get a severe rash more than 10 days  
■ fever lasts more than 3 days  
■ if ringing in the ears or loss of hearing occurs  
■ redness or swelling is present  
■ no symptoms appear

■ pregnant or breast feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.  
Directions  
■ Drink a full glass of water with each dose.  
Adults and children (12 years and older): Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.  
Children under 12 years: Consult a doctor.  
Other information  
■ Store at room temperature  
■ avoid excessive heat and humidity  
■ longer evident sealed packets  
■ do not use any opened or torn packets  
Inactive ingredients  
corn starch, hypromellose\*, microcrystalline cellulose\*, polyethylene glycol\*, povidone\*, propylene glycol\*  
\*may contain  
Questions or comments? 1-800-356-9100

**Información del medicamento**

**Ingrediente activo (en cada tableta)**  
Aspirina 325 mg (AINE)  
Alivia el dolor/Reduce la fiebre  
Medicamento anti-inflamatorio no esteroideo

**Propiedad**  
Analgésico y antipirético

**Usos:**  
Alivia temporalmente:  
■ dolor de cabeza ■ dolores musculares  
■ los dolores de artritis ■ dolor de muelas  
■ dolor dental ■ dolores menstruales

**Advertencias:**  
El síndrome de Reye: Los niños administrados a niños y adolescentes enfermos o en recuperación de los síntomas de la varicela o de la influenza A la gripe. Si con el consumo de este medicamento se presentan cambios en el comportamiento de la persona junto con náuseas y vómitos, consulte con un doctor ya que estos síntomas pueden señalar el comienzo del síndrome de Reye, una enfermedad, que aunque es poco común es grave.  
Alerta a las alergias: La aspirina puede causar severas reacciones alérgicas tan como pueden incluir:  
■ urticaria ■ inflamación facial ■ asma (respiración con silbido) ■ shock  
Si se presenta una reacción alérgica, de inmediato suspenda su consumo y obtenga atención médica.  
Advertencia acerca del sangrado estomacal: Este producto contiene un medicamento anti-inflamatorio no esteroideo (AINE), el cual puede causar sangrado estomacal. Las posibilidades de que ocurra este sangrado:  
■ tiene 60 o más años de edad  
■ ha tenido úlceras estomacales o problemas de sangrado  
■ toma medicamentos para el dolor (anticoagulantes) o esteroideos  
■ toma otra medicina con o sin receta médica que contenga un AINE (aspirina, ibuprofeno, naproxeno, u otros)  
■ toma 2 o más bebidas alcohólicas diariamente mientras consume este producto  
■ si se consume más o por más tiempo de lo indicado  
No se administre a usted si es alérgico a la aspirina o a cualquier otro analgésico/reductor de fiebre.  
Consulte con un médico antes de consumir, si la advertencia acerca del sangrado estomacal debe aplicarse en su caso.  
Evite el uso de productos de protección estomacal tales como: ácido estomacal  
■ tiene la presión sanguínea alta, enfermedad del corazón, crisis del hígado o enfermedad de los riñones  
■ está tomando un diurético  
■ prefiere de aspirina

**Información del medicamento (continuación)**  
Consulte con un médico o farmacéutico antes de administrarle al usted está tomando un medicamento recetado para:  
■ la ginecología ■ la diabetes ■ la artritis  
Dado de tomar este producto consulte con el médico si:  
■ ocurre una reacción alérgica. Busque ayuda médica inmediatamente.  
■ se presentan cambios en los siguientes síntomas de sangrado estomacal:  
■ se siente débil ■ vómito sanguinoso  
■ o si su excremento contiene sangre o tiene un aspecto negro  
■ tiene dolor estomacal que no mejora  
■ el dolor empeora o persiste por más de 10 días  
■ la fiebre persiste por más de 10 días  
■ ocurre zumbido en los oídos o pérdida de la audición  
■ aparece enrojecimiento o hinchazón  
■ aparece un síntoma nuevo  
Si está embarazada o en periodo de lactancia, consulte la opinión de un médico antes de consumir este producto. Es sobre todo importante que no se consuma aspirina durante los últimos 3 meses del embarazo, a menos que el médico así lo haya especificado, ya que esto podría causar problemas a la criatura que está por nacer o traer complicaciones durante el parto.  
Barridos: Evite beber los medicamentos fuera del alcance de los niños. En caso de ocurrir una sobredosis obtenga inmediata atención médica o llame a un centro de control de envenenamiento (Poison Control Center).

**Indicaciones**  
■ con cada dosis beba un vaso lleno con agua  
Adultos y niños (de 12 años y mayores): Tome 1 o 2 tabletas 4 horas o 3 tabletas cada 6 horas, no exceda 12 tabletas en 24 horas.  
Niños menores de 12 años: Consulte al médico.

**Otra información**  
■ el medicamento conservarlo a temperatura ambiente  
■ evite exponerlo al calor y humedad excesiva  
■ mantévalo con el sello sellado  
■ no use ningún paquete abierto o roto

**Ingredientes inactivos**  
maiz de almidón, croscelina, celulosa microcristalina\*, polietilenglicol\*, povidona\*, propilenglicol\*,  
\*puede contener:  
■ ¿Tiene alguna pregunta o comentario? Llame al 1-800-356-9100

Retain carton for complete product information

**ASPIRIN, PAIN RELIEVER (NSAID)**

Reorder No. 23578  
Reorder: 800-356-9100  
cuney.com

DISTRIBUTED BY:  
CONNELLY SAFETY PRODUCTS  
MADISON, WI 53744-4190

Industrial Pack 500  
(250 Packets of 2 Tablets)

Conservar la caja para obtener la información completa del producto

PAQUETES CON SELLADO EVIDENTE.  
NO USE NINGÚN PAQUETE  
ABIERTO O ROTO.

Reorder No. 23578  
Reorder: 800-356-9100  
cuney.com

DISTRIBUIDO POR:  
CONNELLY SAFETY PRODUCTS  
MADISON, WI 53744-4190

Direct Safety®

Aspirin

Pain Reliever

Aspirin 325 mg (NSAID)

Aspirina 325 mg (AINE)

This Package is for Households without Young Children.

Este Paquete Es Hogares Sin Niños Pequeños

Industrial Pack 500

(250 Packets of 2 Tablets)

**DIRECT SAFETY. ASPIRIN Pain Reliever DIRECT SAFETY.**

**Aspirin 325 mg (NSAID) Aspirina 325 mg (AINE)**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)  
\*nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever/fever reducer

**Uses**  
temporarily relieves  
headache  
pain and fever of colds

**Warnings**  
Haley's syndrome: Children and teenagers who have not recovered from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older  
have had stomach ulcers or bleeding problems  
take a blood thinner (anticoagulant or blood drug)  
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
have 2 or more alcohol drinks every day while using this product  
take more for a longer time than directed

**Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer**

**Ask a doctor before use if:**  
stomach bleeding warning applies to you  
you have a history of stomach problems such as heartburn  
you have high blood pressure, heart disease, liver problems, or kidney disease  
you are taking a diuretic  
you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:**  
cold, flu, or sinus  
arthritis

**ASPIRIN**

**Pain Reliever**

Aspirin 325 mg (NSAID)  
Aspirina 325 mg (AINE)

**THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.**

**Información del medicamento**

**Ingrediente activo (en cada tableta)**  
Aspirina 325 mg (AINE)  
\*antiinflamatorio no esteroideo

**Propiedad**  
Alivio de dolor/febres y fiebre

**Usos:**  
alivia temporal de  
dolor de cabeza  
fiebre debida de resaca  
dolor de muelas

**Advertencias:**  
El síndrome de Haley: No debe administrarse a niños y adolescentes enfermos o en recuperación de los síntomas de la varicela o de la influenza o de la gripe. Si con el consumo de este medicamento aparecen cambios en el comportamiento de la persona junto con náuseas y vómitos, consulte con un doctor ya que estos síntomas pueden señalar el comienzo del síndrome de Reye, una enfermedad que aunque es poco común es grave.

**Atención a las alergias:** La aspirina puede causar severas reacciones alérgicas las cuales pueden incluir:  
urticaria  
inflamación facial  
ataque

**Advertencia acerca del sangrado estomacal:** Este producto contiene un medicamento antiinflamatorio no esteroideo (AINE), el cual pudiera causar sangrado estomacal. Las posibilidades de que ocurra este suceso son:  
si tiene 60 o más años de edad  
si ha tenido úlceras estomacales o problemas de sangrado  
si toma medicamentos para el adelgazamiento de la sangre (anticoagulantes) o esteroides  
si toma otros medicamentos con su receta médica que contengan un AINE (aspirina, naproxeno, ibuprofeno, u otros)  
si toma 2 o más bebidas alcohólicas diariamente mientras consume este producto  
si se consume más o por más tiempo de lo indicado

**no se administre si usted es alérgico a la aspirina o a cualquier otro analgésico/reductor de fiebre**

**Consulte con un médico antes de consumir, si:**  
la advertencia acerca del sangrado estomacal se aplica en su caso  
tiene antecedentes de problemas estomacales tales como acidez estomacal  
si toma la presión sanguínea alta, enfermedad del corazón, ataques del tigrado o enfermedad de los riñones  
está tomando un estudio  
padece de asma

**Retain carton for complete product information**

**ASPIRIN, PAIN RELIEVER (NSAID)**

TAMPER EVIDENT PACKETS.  
DO NOT USE IF OPEN OR TORN.

Reorder No. 23578  
Reorder: 800-356-9100  
cvs.com

DISTRIBUTED BY:  
CONNER SAFETY PRODUCTS  
MADISON, WI 53744-4190

Industrial Pack 500  
(250 Packets of 2 Tablets)

PAQUETES CON SELLADO EVIDENTE.  
NO USE NINGÚN PAQUETE  
ABIERTO O ROTO.

Reorder No. 23578  
Reorder: 800-356-9100  
cvs.com

DISTRIBUIDO POR:  
CONNER SAFETY PRODUCTS  
MADISON, WI 53744-4190

# DIRECT SAFETY ASPIRIN

aspirin tablet, coated

## Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:61245-620

Route of Administration

ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R3D61U)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3S)	

## Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61245-620-13	250 in 1 BOX	02/15/2021	04/01/2024
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2021	04/01/2024

## DIRECT SAFETY ASPIRIN

aspirin tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61245-616
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

## Product Characteristics

Color	white	Score	no score
-------	-------	-------	----------

<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61245-616-13	250 in 1 BOX	02/15/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2021	

**Labeler** - Direct Safety (023316151)

**Registrant** - Unifirst First Aid Corporation (832947092)

### Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(61245-616, 61245-620)

Revised: 10/2024

Direct Safety