

ALEVE ARTHRITIS PAIN GEL- diclofenac sodium gel
Bayer HealthCare LLC

Aleve Arthritis Pain Gel UI 1614804

Drug Facts

Active Ingredient

Diclofenac sodium (NSAID*) 1% (equivalent to 0.93% diclofenac)

*nonsteroidal anti-inflammatory drug

Arthritis pain reliever

Uses

- for temporary relief of arthritis pain ONLY in the following areas:
 - hand, wrist, elbow (upper body areas)
 - foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

For external use only

Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Liver Warning: This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- apply more or for longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises, or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under 18 years.

Ask a doctor or pharmacist before use

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- avoid contact with eyes, nose, or mouth
- if eye contact occurs, rinse thoroughly with water

Stop use and ask a doctor if

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling

- weakness in one part or side of body
- slurred speech

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use this product during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

Use up to 21 days unless directed by your doctor.

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily

For your arthritis pain:

- Use 4 times per day every day
- Do not use on more than 2 body areas at the same time

Per Dose

Use ENCLOSED DOSING CARD to measure a dose

- For each upper body area (hand, wrist or elbow)- Squeeze out 2.25 inches (2 grams)
 - For each lower body (foot, ankle or knee)- Squeeze out 4.5 inches (4 grams)
-

Read the enclosed User Guide for complete instructions:

- use only as directed
- do not use more than directed for longer than directed
- apply to only clean, dry skin that does not have cuts, open wounds, infections or rashes
- do not apply in the same area as any other product
- do not apply with external heat such as heating pad
- do not apply to bandage over the treated area
- store ENCLOSED DOSING CARD with your ALEVE Arthritis Pain. The dosing cards is re-usable.

Other information

- store at 20-25°C (68-77°F). Keep from freezing.
- read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients

CARBOMER HOMOPOLYMER TYPE C, COCOYL CAPRYLOCAPRATE, FRAGRANCE,

ISOPROPYL ALCOHOL; MINERAL OIL; POLYOXYL 20 CETOSTEARYL ETHER; PROPYLENE GLYCOL, PURIFIED WATER, STRONG AMONIA SOLUTION

Questions or Comments 1-800-395-0689

NEW!

DIFFERENT

INGREDIENT

DICLOFENAC

SODIUM

ALEVE® ARTHRITIS

PAIN GEL

Diclofenac Sodium Topical Gel 1% (NSAID)

Arthritis Pain Reliever

FOR DAILY

TREATMENT OF

ARTHRITIS PAIN

ANTI-INFLAMMATORY

FOR EXTERNAL USE ONLY

NET WT 3.53 OZ (100 g)

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DICLOFENAC SODIUM (UNII: QTG126297Q) (DICLOFENAC - UNII:14408QL0L1)	DICLOFENAC SODIUM	1 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
WATER (UNII: 059QF0K00R)	
POLYOXYL 20 CETOSTEARYL ETHER (UNII: YRC528SWUY)	
COCOYL CAPRYLOCAPRATE (UNII: 8D9H4QU99H)	
AMMONIA (UNII: 5138Q19F1X)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
MINERAL OIL (UNII: T5L8T28FGP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0039-01	1 in 1 CARTON	06/01/2021	
1		100 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0280-0039-02	1 in 1 CARTON	06/01/2021	
2		50 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211253	06/01/2021	

Labeler - Bayer HealthCare LLC (112117283)

Revised: 12/2025

Bayer HealthCare LLC