

**TRAVEL SAVVY ASPIRIN- aspirin tablet, coated**  
**Bellegrove Medical Supply, Inc.**

-----  
**Travel Savvy Aspirin**

***Drug Facts***

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves

- headache
- muscle pain
- minor pain of arthritis
- pain and fever of colds
- toothache
- menstrual pain

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose

**Adults and children 12 years and older:** Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

**Children under 12 years:** Consult a doctor

***Other information***

- store at room temperature
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

***Inactive ingredients***

corn starch, hypromellose\*, microcrystalline cellulose\*, polyethylene glycol\* povidone\*, propylene glycol\*

\* may contain

***Questions or comments?*** 1-800-323-1319

BTravelSavvy.com

Travel Savvy™

Convenient, Single-Dose

Aspirin

for aches and fevers

Aspirin • 325 mg • NSAID

Pain Reliever • Fever Reducer


Temporarily reduces fever and relieves minor aches and pains from headache, arthritis and inflammation.

1000

Tablets

(500 x 2)

BTravelSavvy.com BTravelSavvy.com



Convenient, Single-Dose  
**Aspirin**  
for aches and fevers

Aspirin • 325 mg • NSAID  
Pain Reliever • Fever Reducer

Compare to  
Genuine Bayer® aspirin  
active ingredient

Aspirin • 325 mg • NSAID  
Pain Reliever • Fever Reducer


1000 Tablets (500 x 2) 1000 Tablets (500 x 2)

Temporarily reduces fever and relieves minor aches and pains from headache, arthritis and inflammation.

**Single-Dose Aspirin Pain Reliever Fever Reducer**

in convenient to-go packets


Temporarily reduces fever and relieves minor aches and pains from headache, arthritis and inflammation.



THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN. THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

BTravelSavvy.com  
Travel Savvy™  
Convenient, Single-Dose  
Aspirin  
for aches and fevers  
Aspirin • 325 mg • NSAID  
Pain Reliever • Fever Reducer  
Temporarily reduces fever and relieves minor aches and pains from headache, arthritis and inflammation.  
1000  
Tablets  
(500 x 2)

BTravelSavvy.com BTravelSavvy.com



Convenient, Single-Dose  
**Aspirin**  
for aches and fevers

Aspirin • 325 mg • NSAID  
Pain Reliever • Fever Reducer

Compare to  
Genuine Bayer® aspirin  
active ingredient

Aspirin • 325 mg • NSAID  
Pain Reliever • Fever Reducer


1000 Tablets (500 x 2) 1000 Tablets (500 x 2)

Temporarily reduces fever and relieves minor aches and pains from headache, arthritis and inflammation.

**Single-Dose Aspirin Pain Reliever Fever Reducer**

in convenient to-go packets

Temporarily reduces fever and relieves minor aches and pains from headache, arthritis and inflammation.



THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN. THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

**TRAVEL SAVVY ASPIRIN**  
aspirin tablet, coated

DRUG FACTS		DRUG FACTS (CONTINUED)
<b>Active ingredient (in each tablet)</b> Aspirin 325 mg (NSAID) <sup>1</sup>	<b>Purpose</b> Pain reliever/fever reducer nonsteroidal anti-inflammatory drug	<b>Stop use and ask a doctor if</b> • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • red, black, or vomit blood • hard black or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • have lightheadedness or loss of hearing occurs • redness or swelling is present • new symptoms appear
<b>Uses</b> Temporarily relieves • headache • muscle pain • minor pain of arthritis • pain and fever of colds • toothache • menstrual pain	<b>Warnings</b> <b>Reyes' syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reyes' syndrome, a rare but serious illness. <b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • rash • asthma (wheezing) If an allergic reaction occurs, stop use and seek medical help right away. <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcohol drinks every day while using this product • take more or for a longer time than directed	<b>Directions</b> • drink a full glass of water with each dose <b>Adults and children:</b> Take 1 or 2 tablets every 4 hours or 3 tablets (12 years and older) every 6 hours, not to exceed 12 tablets in 24 hours <b>Children under 12 years:</b> Consult a doctor
<b>Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer</b> <b>Ask a doctor before use if</b> • stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn or kidney disease • you are taking a diuretic • you have asthma <b>Ask a doctor or pharmacist before use if you are taking a prescription drug for</b> • gout • diabetes • arthritis	<b>Other information</b> • avoid excessive heat and humidity • tamper evident sealed packets • do not use any opened or torn packets <b>Inactive ingredients</b> corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, polydextrose, polypropylene glycol *may contain <b>Questions or comments?</b> 1-800-323-1319	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.

Retain carton for complete product information

Distributed by  
**Bellevue medical**  
8349 154th Ave NE, Redmond, WA 98052

Retailer #59961-129-02  
Packaged for Bellevue by Medique Products  
Fort Meyers, FL, USA 33967

5996112902

DRUG FACTS		DRUG FACTS (CONTINUED)
<b>Active ingredient (in each tablet)</b> Aspirin 325 mg (NSAID) <sup>1</sup>	<b>Purpose</b> Pain reliever/fever reducer nonsteroidal anti-inflammatory drug	<b>Stop use and ask a doctor if</b> • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • red, black, or vomit blood • hard black or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • have lightheadedness or loss of hearing occurs • redness or swelling is present • new symptoms appear
<b>Uses</b> Temporarily relieves • headache • muscle pain • minor pain of arthritis • pain and fever of colds • toothache • menstrual pain	<b>Warnings</b> <b>Reyes' syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reyes' syndrome, a rare but serious illness. <b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • rash • asthma (wheezing) If an allergic reaction occurs, stop use and seek medical help right away. <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcohol drinks every day while using this product • take more or for a longer time than directed	<b>Directions</b> • drink a full glass of water with each dose <b>Adults and children:</b> Take 1 or 2 tablets every 4 hours or 3 tablets (12 years and older) every 6 hours, not to exceed 12 tablets in 24 hours <b>Children under 12 years:</b> Consult a doctor
<b>Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer</b> <b>Ask a doctor before use if</b> • stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn or kidney disease • you are taking a diuretic • you have asthma <b>Ask a doctor or pharmacist before use if you are taking a prescription drug for</b> • gout • diabetes • arthritis	<b>Other information</b> • avoid excessive heat and humidity • tamper evident sealed packets • do not use any opened or torn packets <b>Inactive ingredients</b> corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, polydextrose, polypropylene glycol *may contain <b>Questions or comments?</b> 1-800-323-1319	<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.

Retain carton for complete product information

Distributed by  
**Bellevue medical**  
8349 154th Ave NE, Redmond, WA 98052

Retailer #59961-129-02  
Packaged for Bellevue by Medique Products  
Fort Meyers, FL, USA 33967

5996112902

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59961-620
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	FR21
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59961-620-15	500 in 1 BOX	02/15/2021	04/01/2024
1		2 in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	02/15/2021	04/01/2024

**TRAVEL SAVVY ASPIRIN**

aspirin tablet, coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59961-616
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

**Product Characteristics**

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59961-616-15	500 in 1 BOX	02/15/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2021	

**Labeler** - Bellegrove Medical Supply, Inc. (141647094)**Registrant** - Unifirst First Aid Corporation (832947092)**Establishment**

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(59961-620, 59961-616)

**Establishment**

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(59961-616, 59961-620)

