

**BOVAGUARD BARRIER DIP- iodine solution**  
**AgroChem Inc**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

**BovaGuard Barrier Dip**

**BOVAGUARD BARRIER DIP**

**IODINE**

**TEAT DIP**

**WARNING**

**CAUSES SERIOUS EYE IRRITATION**

**FOR FARM AND INDUSTRIAL USE ONLY.**

**KEEP OUT OF REACH OF CHILDREN.**

**KEEP OUT OF DIRECT SUNLIGHT.**

**AVOID FREEZING**

**PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.**

**FIRST AID / RESPONSE:**

**IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.**

**STORAGE: Store locked up.**

**DISPOSAL: Dispose of contents/ container to an approved waste disposal plant**

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**INGREDIENTS:**

**Active Ingredient - Iodine 1.0 % (10.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution**

**Emollients: Glycerine 10%**

**BovaGuard is made with IoPlex technology and is NPE free.**

**Use BovaGuard Barrier to reduce the spread of mastitis causing organisms. BovaGuard Barrier is an effective and economical 1% iodine barrier teat dip which is both non-drip and highly water resistant. Recommended for post milking applications.**

**Directions:**

**General Use**

**Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.**

**Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.**

**If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.**

#### **Pre Dipping Use**

**Perform forestripping and inspect teats for clinical mastitis.**

**Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.**

#### **Post Dipping Use**

**Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.**

#### **Extended Use**

**When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.**

# BOVAGUARD BARRIER DIP

## IODINE TEAT DIP

### WARNING

**CAUSES SERIOUS EYE IRRITATION  
FOR FARM AND INDUSTRIAL USE ONLY.  
KEEP OUT OF REACH OF CHILDREN.  
KEEP OUT OF DIRECT SUNLIGHT.  
AVOID FREEZING**

**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

**FIRST AID / RESPONSE:**

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/ container to an approved waste disposal plant

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**INGREDIENTS:**

Active Ingredient - Iodine 1.0 % (10.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution  
Emollients: Glycerine 10%

BovaGuard is made with loPlex technology and is NPE free.

**NET CONTENTS:**



3170B-BULK



Use BovaGuard Barrier to reduce the spread of mastitis causing organisms. BovaGuard Barrier is an effective and economical 1% iodine barrier teat dip which is both non-drip and highly water resistant. Recommended for post milking applications.

**Directions:**

**General Use**

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

**Pre Dipping Use**

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

**Post Dipping Use**

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

**Extended Use**

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.



**AgroChem**

26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

# BOVAGUARD BARRIER DIP

## IODINE TEAT DIP

### WARNING



**CAUSES SERIOUS EYE IRRITATION  
FOR FARM AND INDUSTRIAL USE ONLY.  
KEEP OUT OF REACH OF CHILDREN.  
KEEP OUT OF DIRECT SUNLIGHT.  
AVOID FREEZING**

**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

**FIRST AID / RESPONSE:**

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/ container to an approved waste disposal plant

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**INGREDIENTS:**

Active Ingredient - Iodine 1.0 % (10.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution  
Emollients: Glycerine 10%

BovaGuard is made with loPlex technology and is NPE free.

**NET CONTENTS:**



31708-BULK

Use BovaGuard Barrier to reduce the spread of mastitis causing organisms. BovaGuard Barrier is an effective and economical 1% iodine barrier teat dip which is both non-drip and highly water resistant. Recommended for post milking applications.

**Directions:**

**General Use**

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

**Pre Dipping Use**

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

**Post Dipping Use**

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

**Extended Use**

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.



## AgroChem

26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

## BOVAGUARD BARRIER DIP

iodine solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17307-0001
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	.10 kg in 1 kg

### Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	.83 kg in 1 kg
GLYCERIN (UNII: PDC6A3C0OX)	.04 kg in 1 kg
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	.019 kg in 1 kg

BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	.01 kg in 1 kg
XANTHAN GUM (UNII: TTV12P4NEE)	.0037 kg in 1 kg
ALGINIC ACID (UNII: 8C3Z4148WZ)	.002 kg in 1 kg

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-0001-1	19 kg in 1 PACKAGE		
2	NDC:17307-0001-2	58 kg in 1 DRUM		
3	NDC:17307-0001-3	116 kg in 1 DRUM		
4	NDC:17307-0001-4	214.5 kg in 1 DRUM		
5	NDC:17307-0001-5	1068 kg in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		07/01/2013	

**Labeler** - AgroChem Inc (805427700)

### Establishment

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	manufacture, api manufacture

Revised: 2/2018

AgroChem Inc