

**MAXIMUM STRENGTH DM MAX- dextromethorphan hydrobromide,  
guaifenesin liquid  
H E B**

-----  
**H E B Maximum Strength DM MAX**

***Drug Facts***

***Active ingredients (in each 20 mL)***

Dextromethorphan HBr 20 mg  
Guaifenesin 400 mg

***Purpose***

Dextromethorphan HBr ... Cough suppressant  
Guaifenesin .....Expectorant

***Uses***

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - cough due to minor throat and bronchial irritation

***Warnings***

Do not use • for children under 12 years of age  
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

***Ask a doctor before use if you have***

- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.
- cough that occurs with too much phlegm (mucus)

**When using this product**

- do not use more than directed

**Stop use and ask a doctor if**

- cough lasts for more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- take only as recommended
- use dosage cup
- mL = milliliter
- do not take more than 6 doses in any 24-hour period

<b>Age</b>	<b>Dose</b>
Adults & children 12 years and older	20 ml every 4 hours
Children under 12 Years of age	Do not use

**Other information**

- each 20 mL contains: sodium 12 mg
- dosage cup provided
- store between 15-30° C (59-86° F)
- do not refrigerate

**Inactive ingredients**

anhydrous citric acid, dextrose, D&C Red #33, FD&C Red #40, flavors, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate,

sucralose, Xanthan gum

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at

**1-877-798-5944**

**Product Label**

**Compare** to **Mucinex<sup>®</sup> FAST-MAX<sup>™</sup>**

**DM MAX** active ingredients\*

NDC 37808-518-25

**H-E-B<sup>®</sup>**

**Maximum Strength  
DM MAX**

**Dextromethorphan HBr /  
Cough Suppressant  
Guaifenesin / Expectorant**

**Congestion & Cough**

**Controls Cough  
Thins & Loosens Mucus  
Relieves Chest Congestion**

**Adults**

**For Ages 12 & Over**

**6 FL OZ (177mL)**

LF 021

***Peel Corner to Read Complete  
Drug Facts and Information →***

**DO NOT USE IF PRINTED SEAL  
UNDER CAP IS TORN OR MISSING**

**H-E-B<sup>®</sup>**

**100%  
GUARANTEE  
*promise***

**If you aren't completely pleased  
with this product, we'll be happy to  
replace it or refund your money.  
You have our word on it.**

LB-021

12194-1905

0 41220 83197 5

**Lot:**  
**Exp:**

**\*This product is not manufactured or distributed by Reekitt Benckiser Inc., distributor of Mucinex® FAST-MAX™ DM MAX**

**MADE WITH PRIDE & CARE FOR H-E-B®  
SAN ANTONIO, TEXAS, 782044**



Peel Corner to Read Complete Drug Facts and Information



DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING



12194-1905



Lot:  
Exp:

**Drug Facts**

Active ingredients (in each 20 mL)	Purpose
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to clear bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - cough due to minor throat and bronchial irritation

**Warnings**

- Do not use for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not show if your prescription drug contains an MAOI, ask a

**Drug Facts (continued)**

doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- cough that lasts or is chronic such as occur with smoking, asthma, chronic bronchitis or emphysema,
- cough that occurs with too much phlegm (mucus)

**When using this product**

- do not use more than directed

**Stop use and ask a doctor if**

- cough lasts for more than 7 days, comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**

**Directions**

- take only as recommended
- use dosage cup
- mL = milliliter
- do not take more than 6 doses in any 24-hour period

Age	Dose
Adults & children 12 years and older	20 mL every 4 hours
Children under 12 Years of age	Do not use

**Other information**

- each 30 mL contains: sodium 12 mg
- dosage cap provided
- store between 15-30°C (50-86°F)
- do not refrigerate

**Inactive ingredients**

anhydrous citric acid, disodium phosphate, FD&C Red #33, FD&C Red #40, flavors, glycerin, methylcellulose, polyethylene glycol, purified water, saccharin sodium, sodium benzoate, sucralose, Xanthan gum

**Drug Facts (continued)**

**Questions?** Call weekdays from 9:30 AM to 4:30 PM CST at 1-877-798-5344

\* This product is not manufactured or distributed by Reekitt Benckiser Inc., distributor of Mucinex® FAST-MAX™ DM MAX

**MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204**

res

**MAXIMUM STRENGTH DM MAX**

dextromethorphan hydrobromide, guaifenesin liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-518
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-518-25	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/14/2012	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M012	09/14/2012	
--------------------	------	------------	--

**Labeler -** H E B (007924756)

**Registrant -** AptaPharma Inc. (790523323)

**Establishment**

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(37808-518)

Revised: 12/2023

H E B