

PEPTO BISMOL CHEWS- bismuth subsalicylate tablet, chewable
The Procter & Gamble Manufacturing Company

Pepto Bismol® Chews

Drug Facts

Active ingredient (in each chewable tablet)

Bismuth subsalicylate 262 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer

- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- fully chew before swallowing
- adults and children 12 years and over:
- 2 chewable tablets every ½ hour or 4 chewable tablets every hour as needed for diarrhea
- 2 chewable tablets every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 chewable tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each chewable tablet contains:**
- calcium 50 mg • salicylate 117 mg • low sodium

- Store at no greater than 25°C. Avoid excessive heat (over 40°C.)

Inactive ingredients

acacia, anhydrous citric acid, calcium carbonate, caprylic/capric triglyceride, carmine, carnauba wax, corn starch, corn syrup, dextrin, flavor, hydrogenated coconut oil, maltodextrin, potassium hydroxide, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide, water, white wax

Questions?

1-800-717-3786

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.

**DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202**



PRINCIPAL DISPLAY PANEL - 24 CHEWABLE TABLETS

Pepto Bismol®

Bismuth Subsalicylate - Upset Stomach Reliever/Antidiarrheal

CHEWS

24 CHEWABLE TABLETS



Drug Facts

Active ingredient (in each chewable tablet) Bismuth subsalicylate 262 mg

Purpose Upset stomach reliever and antidiarrheal

Uses relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are
 • allergic to salicylates (including aspirin)
 • taking other salicylate products

Do not use if you have • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
 • fully chew before swallowing
 • adults and children 12 years and over:
 • 2 chewable tablets every 1/2 hour or 4 chewable tablets every hour as needed for diarrhea
 • 2 chewable tablets every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 • do not exceed 8 doses (16 chewable tablets) in 24 hours
 • use until diarrhea stops but not more than 2 days
 • children under 12 years: ask a doctor
 • drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Inactive ingredients acacia, anhydrous citric acid, calcium carbonate, caprylic/capric triglyceride, carmine, carnauba wax, corn starch, corn syrup, dextrin, flavor, hydrogenated coconut oil.

maltodextrin, potassium hydroxide, propylene glycol, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide, water, white wax

Other information • each chewable tablet contains:
 • calcium 50 mg • salicylate 117 mg • low sodium
 • Store at no greater than 25°C. Avoid excessive heat (over 40°C).

Questions? 1-800-717-3786

P&G
 Patents: www.pg.com/patents

| PEPTO BISMOL CHEWS | | | |
|--|----------------|---------------------------|-----------------|
| bismuth subsalicylate tablet, chewable | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:37000-996 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |
| Ingredient Name | | Basis of Strength | Strength |
| BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ, BISMUTH CATION - UNII:ZS9CD118YE) | | BISMUTH SUBSALICYLATE | 262 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SORBIC ACID (UNII: X045WJ989B) | |
| WATER (UNII: 059QF0KO0R) | |
| HYDROGENATED COCONUT OIL (UNII: JY81OXM1OM) | |
| SORBITOL (UNII: 506T60A25R) | |
| ICODEXTRIN (UNII: 2NX48Z0A9G) | |
| ACACIA (UNII: 5C5403N26O) | |
| WHITE WAX (UNII: 7G1J5DA97F) | |
| CALCIUM CARBONATE (UNII: H0G9379FGK) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| CORN SYRUP (UNII: 9G5L16BK6N) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) | |
| SHELLAC (UNII: 46N107B71O) | |
| CARMINIC ACID (UNII: CID8Z8N95N) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| SUCROSE (UNII: C151H8M554) | |
| CAPRYLIC/CAPRIC TRIGLYCERIDE (UNII: C9H2L21V7U) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| SOYBEAN LECITHIN (UNII: 1DI56QDM62) | |

Product Characteristics

| | | | |
|-----------------|-------------|---------------------|----------|
| Color | pink | Score | no score |
| Shape | ROUND | Size | 18mm |
| Flavor | WINTERGREEN | Imprint Code | PB |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:37000-996-24 | 24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 09/01/2020 | |
| 2 | NDC:37000-996-72 | 3 in 1 CELLO PACK | 09/01/2020 | |
| 2 | | 24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M008 | 09/01/2020 | |

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 3/2025

The Procter & Gamble Manufacturing Company