

**DELFLEx- dextrose monohydrate, sodium chloride, sodium lactate, calcium chloride, magnesium chloride solution**  
**DELFLEx- dextrose monohydrate, sodium chloride, sodium lactate, calcium chloride, magnesium chloride solution**  
**Fresenius Medical Care North America**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use DELFLEx<sup>®</sup> safely and effectively. See full prescribing information for DELFLEx<sup>®</sup>

**DELFLEx (dextrose) peritoneal dialysis solution**

Initial U.S. Approval: 1984

**DELFLEx Low Magnesium, Low Calcium (dextrose) peritoneal dialysis solution**

Initial U.S. Approval: 1992

----- **INDICATIONS AND USAGE** -----

For treatment of chronic kidney failure. ( 1)

----- **DOSAGE AND ADMINISTRATION** -----

For intraperitoneal dialysis only. ( 2)

----- **DOSAGE FORMS AND STRENGTHS** -----

DELFLEx solutions are available in multiple compositions, calculated osmolarity, pH, and ionic concentrations. See full prescribing information for detailed descriptions of each formulation. ( 3, 11)

----- **CONTRAINDICATIONS** -----

None (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Monitor patient for electrolyte, fluid, and nutrition imbalances. ( 5.1)
- Encapsulating Peritonitis Sclerosis (EPS) ( 5.2)
- Peritonitis: Initiate appropriate antimicrobial therapy ( 5.2)
- Monitor for Lactic Acidosis in patients at risk. ( 5.3)

----- **ADVERSE REACTIONS** -----

Adverse reactions may include peritonitis, catheter site infection, electrolyte and fluid imbalances, hypovolemia, hypervolemia, hypertension, disequilibrium syndrome, muscle cramping, abdominal pain, abdominal distension, and abdominal discomfort. ( 6)

**To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Medical Care North America at 1-800-323-5188 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**See 17 for PATIENT COUNSELING INFORMATION.**

Revised: 3/2023

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1. INDICATIONS AND USAGE**

**2. DOSAGE AND ADMINISTRATION**

- 2.1 Basic Dosing Information
- 2.2 Administration Instructions
- 2.3 Compatible Medications

**3. DOSAGE FORMS AND STRENGTHS**

**4. CONTRAINDICATIONS**

**5. WARNINGS AND PRECAUTIONS**

- 5.1 Electrolyte, Fluid and Nutrition Imbalances
- 5.2 Peritonitis and Encapsulating Peritoneal Sclerosis
- 5.3 Lactic Acidosis
- 5.4 Over Infusion

**6. ADVERSE REACTIONS**

**8. USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use

**11. DESCRIPTION**

**12. CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

**13. NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**16. HOW SUPPLIED/STORAGE AND HANDLING**

**17. PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### 1. INDICATIONS AND USAGE

DELFLEx® is indicated in the treatment of chronic kidney failure in patients being maintained on peritoneal dialysis.

### 2. DOSAGE AND ADMINISTRATION

#### 2.1 Basic Dosing Information

DELFLEx® is intended for intraperitoneal administration only. Not for intravenous or intra-arterial administration.

The mode of therapy, frequency of treatment, formulation, exchange volume, duration of dwell, and length of dialysis should be selected by the physician responsible for the treatment of the individual patient.

Utilize the peritoneal dialysis solution with lowest level of osmolarity consistent with the fluid removal requirements for that exchange.

Do not store solutions containing additives.

#### 2.2 Administration Instructions

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.


Do not heat in a microwave oven.

#### Get Ready

1. Clean work surface.
2. Gather supplies:
  - DELFLEx peritoneal dialysis bag(s).
  - Prescribed medication(s), if ordered by your healthcare provider.
  - Mask.
- 3.

#### PVC


Tear the overwrap from the slit edge down the length of the inner bag to open.



Wipe away any moisture from the solution bags. Some opacity may be observed in the plastic of the bag and/or tubing and is due to moisture absorption during the sterilization process. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually.



Locate pull tabs on overwrap. Grasping one tab in each hand, pull outward, down the length of the inner bag to open.



Wipe away any moisture from the solution bags.

#### Inspect DELFLEx Solution Bag

4. After removing the overwrap, check your DELFLEX solution bag(s) for strength, clarity, amount, leaks, and expiration date. Do not use DELFLEX solution if leaks are found, the solution bag is damaged, and/or the solution is cloudy or discolored, or the product is expired. Color may vary from clear to slightly yellow but does not affect efficacy and may be used.
5. Visually check that the solution bag tubing is free from kinks. If kinks are present, straighten tubing to allow the solution to flow freely.  
**Note: Retain DELFLEX peritoneal dialysis bag sample for manufacturer evaluation and notify your healthcare provider if any of the above defects are found.**

**Note: DELFLEX peritoneal dialysis solutions utilize the Safe-Lock® Connection System. This unique system consists of two Safe-Lock connectors, one located on the administration port of the bag, and the mating connector is located on the cyclor set. The Safe-Lock connectors were designed to reduce the potential risk of touch contamination of the internal connection components.**

6. Put on mask. Wash your hands.
7. If you will be adding medications(s):
  - Clean hands (as per facility's protocol)
  - Clean the medication port as instructed by your healthcare provider.
  - Add the medicine(s).
  - Turn the bag upside down several times to mix the medicine(s).
8. To connect the bag(s) to the cyclor set, unscrew the protective caps of the administration port and the cyclor set solution line connector. Secure these two connectors with a twisting motion to lock in place, so that the cyclor set connector is seated over the administration port O-ring to assure a firm and tight fit.
9. After completing Step 8, wait for the cyclor prompt to break the administration port cone and initiate solution flow. Do this by placing the thumb firmly on the tube over the cone and pressing towards the outer wall of the tube and away from the bag.



10. Perform your treatment as prescribed.
11. At the end of your treatment, throw away the fluid and used set as instructed by your healthcare provider. **In case of cloudiness, save the fluid and the used set and immediately contact your healthcare provider.** Dispose of your empty solution bag according to your local recycling program. Empty solution bags may not be recyclable in your area.

### 2.3 Compatible Medications

Compatible medications can be added via the medication port [see *Dosage and Administration ( 2.2)*]. The following medications have demonstrated stability with DELFLEX solutions: cefazolin, ceftazidime, gentamicin, and vancomycin [see *Clinical Pharmacology ( 12.3)*].

### 3. DOSAGE FORMS AND STRENGTHS

DELFLX peritoneal dialysis solutions are available in single-dose flexible bags comprised of either polyvinyl chloride (PVC), or a proprietary blend of polyolefins called Biofine®. All DELFLX peritoneal dialysis solutions have overfills declared on the bag label.

DELFLX peritoneal dialysis solutions are available in the sizes and formulations shown in Table 1.

**Table 1. DELFLX peritoneal dialysis solution sizes and formulations**

PVC				Biofine®		
2L	3L	5L	6L	3L	5L	6L

DELFLEX Standard with 1.5% Dextrose			X	X		X	X
DELFLEX Standard with 2.5% Dextrose			X	X		X	X
DELFLEX Low Magnesium, Low Calcium with 1.5% Dextrose	X	X	X	X	X	X	X
DELFLEX Low Magnesium, Low Calcium with 2.5% Dextrose	X	X	X	X	X	X	X
DELFLEX Low Magnesium, Low Calcium with 4.25% Dextrose	X	X	X	X	X	X	X

#### 4. CONTRAINDICATIONS

None.

#### 5. WARNINGS AND PRECAUTIONS

##### 5.1 Electrolyte, Fluid and Nutrition Imbalances

Peritoneal dialysis may affect a patient's protein, water-soluble vitamin, potassium, sodium, chloride, bicarbonate, and magnesium levels and volume status. Monitor electrolytes and blood chemistry periodically and take appropriate clinical action.

Potassium is omitted from DELFLEX solutions because dialysis may be performed to correct hyperkalemia. In situations where there is a normal serum potassium level or hypokalemia, the addition of potassium chloride (up to a concentration of 4 mEq/L) may be indicated to prevent severe hypokalemia.

To avoid the risk of severe dehydration or hypovolemia and to minimize the loss of protein, it is advisable to select the peritoneal dialysis solution with lowest level of osmolarity consistent with the fluid removal requirements for that exchange.

Significant loss of protein, amino acids and water-soluble vitamins may occur during peritoneal dialysis. Replacement therapy should be provided as necessary.

##### 5.2 Peritonitis and Encapsulating Peritoneal Sclerosis

Infectious and aseptic peritonitis has been associated with peritoneal dialysis therapy. Following DELFLEX use, inspect the drained fluid for the presence of fibrin or cloudiness, which may indicate the presence of peritonitis. Improper clamping or priming sequence may result in infusion of air into the peritoneal cavity, which may result in abdominal pain and/or peritonitis. If peritonitis occurs, treat with appropriate therapy.

Encapsulating peritoneal sclerosis (EPS), sometimes fatal, is a complication of peritoneal dialysis therapy.

##### 5.3 Lactic Acidosis

Monitor patients with conditions known to increase the risk of lactic acidosis [e.g., severe hypotension or sepsis that can be associated with acute kidney failure, inborn errors of metabolism, treatment with drugs such as nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs)] for lactic acidosis before the start of treatment and during treatment with DELFLEX.

Solutions containing the lactate ion should be used with great care in patients with metabolic or respiratory alkalosis. Lactate should be administered with great care in those conditions in which there is an increased level or an impaired utilization of this ion, such as severe hepatic insufficiency.

##### 5.4 Over Infusion

Over infusion of peritoneal dialysis solution volume into the peritoneal cavity may be characterized by abdominal distention, feeling of fullness and/or shortness of breath. Drain the peritoneal dialysis solution from the peritoneal cavity to treat over infusion.


#### 6. ADVERSE REACTIONS

Solution related adverse reactions may include peritonitis, catheter site infection, electrolyte and fluid imbalances, hypovolemia, hypervolemia, hypertension, hypotension, disequilibrium syndrome, muscle cramping, abdominal pain, abdominal distension, and



Dextrose	2.5 g	538 mg	448 mg	18.4 mg	5.08 mg	394	5.5	132	2.5	0.5	95	40
DELFLX Low Magnesium, Low Calcium with 4.25% Dextrose	4.25g	538 mg	448 mg	18.4 mg	5.08 mg	483	5.5	132	2.5	0.5	95	40

Dextrose, USP, is chemically designated D-glucose monohydrate ( $C_6H_{12}O_6 \cdot H_2O$ ) a hexose sugar freely soluble in water. The structural formula is shown here:



Calcium chloride, USP, is chemically designated calcium chloride dihydrate ( $CaCl_2 \cdot 2H_2O$ ) white fragments or granules freely soluble in water.

Magnesium chloride, USP, is chemically designated magnesium chloride hexahydrate ( $MgCl_2 \cdot 6H_2O$ ) colorless flakes or crystals very soluble in water.

Sodium lactate solution, USP, is chemically designated ( $CH_3CH(OH)COONa$ ), a 60% aqueous solution miscible in water.

Sodium chloride, USP, is chemically designated ( $NaCl$ ), a white, crystalline compound freely soluble in water.

Water for injection, USP, is chemically designated ( $H_2O$ ).

Hydrochloric Acid or Sodium Hydroxide may be added for pH adjustment. pH is  $5.5 \pm 0.5$ .

Exposure to temperatures above  $25^\circ C$  ( $77^\circ F$ ) during transport and storage will lead to minor losses in moisture content. Higher temperatures lead to greater losses. It is unlikely that these minor losses will lead to clinically significant changes within the expiration period. Since the inner bag is compounded from flexible plastic, water may permeate from the inner bag into the overwrap in quantities insufficient to affect the solution significantly. Solutions in contact with the plastic inner bag can cause certain chemical components of the bag to leach out in very small amounts; however, the safety of the plastic formulation is supported by biological tests for plastic containers.

## 12. CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

DELFLX peritoneal dialysis solutions are hypertonic peritoneal dialysis solutions containing dextrose, a monosaccharide, as the primary osmotic agent. An osmotic gradient must be created between the peritoneal membrane and the dialysis solution in order for ultrafiltration to occur. The hypertonic concentration of glucose in DELFLX solutions exert an osmotic pressure across the peritoneal membrane resulting in transcapillary ultrafiltration. Like other peritoneal dialysis solutions, DELFLX solutions contain electrolytes to facilitate the correction of acid-base and electrolyte abnormalities. DELFLX solutions contain a buffer, lactate, to help normalize acid-base abnormalities.

### 12.3 Pharmacokinetics

#### Absorption

Glucose can be rapidly absorbed from the peritoneal cavity by diffusion and appears quickly in the circulation due to the high glucose concentration gradient between DELFLX solutions compared to blood capillary glucose level. Absorption per unit time will be the highest at the start of an exchange and decreases over time. The rate of glucose absorption will be dependent upon the transport characteristics of the patient's peritoneal membrane as determined by a peritoneal equilibration test (PET). Glucose absorption will also depend upon the concentration of glucose used for the exchange

and the length of the dwell. Transport of other molecules will be dependent upon the molecular size of the solute, the concentration gradient, and the effective peritoneal surface area as determined by the PET.

#### Metabolism and Elimination

Glucose is metabolized by normal cellular pathways (i.e., glycolysis). Metabolism of lactate occurs in the liver and results in the generation of the bicarbonate. Glucose not absorbed during PD exchange procedure is removed by drainage of the PD solution from the peritoneal cavity.

#### Drug Interaction Studies

##### Antibiotics

No formal clinical drug interaction studies have been performed. In vitro studies of the following medications have demonstrated stability with DELFLEX solutions: cefazolin, ceftazidime, gentamicin, and vancomycin.

### **13. NONCLINICAL TOXICOLOGY**

#### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long term animal studies with DELFLEX peritoneal dialysis solutions have not been performed to evaluate the carcinogenic potential, mutagenic potential or effect on fertility.

### **16. HOW SUPPLIED/STORAGE AND HANDLING**

DELFLX peritoneal dialysis solutions are available in the sizes and formulations shown in Table 1 [see *Dosage Forms and Strengths ( 3)* ] .

**Table 3. DELFLX peritoneal dialysis NDC designations**

	PVC				Biofine <sup>®</sup>		
	2L	3L	5L	6L	3L	5L	6L
Standard 1.5% Dextrose			49230-188-50	49230-188-60		49230-188-52	49230-188-62
Standard 2.5% Dextrose			49230-191-50	49230-191-60		49230-191-52	49230-191-62
Low Mg/Low Ca 1.5% Dextrose	49230-206-20	49230-206-30	49230-206-50	49230-206-60	49230-206-32	49230-206-52	49230-206-62
Low Mg/Low Ca 2.5% Dextrose	49230-209-23	49230-209-30	49230-209-50	49230-209-60	49230-209-32	49230-209-52	49230-209-62
Low Mg/Low Ca 4.25% Dextrose	49230-212-23	49230-212-30	49230-212-50	49230-212-60	49230-212-32	49230-212-52	49230-212-62

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (between 59°F and 86°F). See USP Controlled Room Temperature. Brief exposure to temperatures up to 40°C (104°F) may be tolerated provided the mean kinetic temperature does not exceed 25°C (77°F); however, such exposure should be minimized.

Keep DELFLX and all medicines out of the reach of children.

### **17. PATIENT COUNSELING INFORMATION**

Aseptic technique must be used throughout the procedure and at its termination in order to reduce the possibility of infection.

The solution bag should remain in the carton and the overwrap intact until time of use.

Use only after checking for strength, clarity, amount, leaks, and expiration date.

Advise patients that DELFLX peritoneal dialysis solution should not be heated in a microwave oven.

Care should be taken to ensure that there is not any leakage around the catheter, since if not controlled, the leakage can create edema from subcutaneous infiltration of the dialysis solution. The leakage will also create an inaccurate fluid balance measurement. If any leakage is identified, advise the patient not to proceed with infusion and notify your

physician.



**FRESENIUS  
MEDICAL CARE**

Fresenius Medical Care North America  
920 Winter Street  
Waltham, MA 02451  
1-800-323-5188

© 2020, 2022 Fresenius Medical Care. All Rights Reserved. Fresenius Medical Care, triangle logo, DELFLEX, Safe-Lock, Biofine are trademarks of Fresenius Medical Care Holding, Inc. or affiliated companies.

### Principal Display Panel - 1.5% Dextrose 2000 mL Bag Label

**DELFLX®**

PERITONEAL DIALYSIS SOLUTION  
WITH  
1.5% DEXTROSE  
LOW MAGNESIUM • LOW CALCIUM  
CAT. NO. 044-20221  
NDC 49230-206-20  
2000 mL  
(Approx. 90 mL excess)



<b>DELFLX®</b>	
PERITONEAL DIALYSIS SOLUTION WITH 1.5% DEXTROSE LOW MAGNESIUM LOW CALCIUM	
<b>CAT. NO. 044-20221</b>	<b>2000 mL</b>
NDC 49230-206-20	(Approx. 40 mL excess)
Each 100 mL contains:	
Dextrose, Hydrrous, USP	1.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	344 mOsmol/Liter (calculated)
May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.	
Approximate Milliequivalents Per Liter:	
Sodium 132	Magnesium 0.5
Calcium 2.5	Chloride 95
Lactate 40	
STERILE AND NON PYROGENIC, FOR INTRAPERITONEAL ADMINISTRATION ONLY. USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN. Potassium Chloride to be added only under the direction of a physician. LATEX: This product and its packaging does not contain latex materials. CAUTION: Use only if solution is clear and container is undamaged. Discard any unused portion. Read package insert for full information. Recommended Storage: Room Temperature (25°C). Avoid Excessive Heat. Protect From Freezing. See Insert. Rx only DPL-113 PLASTIC 89-909-04 Rev 01/07	
Fresenius Medical Care	
Fresenius Medical Care NA Waltham, MA 02451 1-800-323-5188	

### Principal Display Panel - 1.5% Dextrose 3000 mL Bag Label

**DELFLX®**

PERITONEAL DIALYSIS SOLUTION  
WITH



1.5% DEXTROSE

LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-30321

NDC 49230-206-30

3000 mL

(Approx. 60 mL excess)



**DELFLEX<sup>®</sup>**  
PERITONEAL DIALYSIS SOLUTION  
WITH  
1.5% DEXTROSE  
LOW MAGNESIUM LOW CALCIUM

CAT. NO. 044-30321                                  3000 mL  
NDC 49230-206-30                                        (Approx. 60 mL excess)

Each 100 mL contains:

Dextrose, Hydrous, USP	1.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.

pH 5.5 (5.0 - 6.0)    344 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:  
Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40

**STERILE AND NON PYROGENIC,  
FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.**

Potassium Chloride to be added only under the direction of a physician.

**LATEX:** This product and its packaging does not contain latex materials.

**CAUTION:** Use only if solution is clear and container is undamaged.  
Discard any unused portion. Read package insert for full information.

**Recommended Storage:** Room Temperature (25°C).  
Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only                                  DPL-113 PLASTIC 89-909-10 Rev 01/07

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

Fresenius Medical Care

**Principal Display Panel - 1.5% Dextrose 5000 mL Bag Label**

**DELFLEX<sup>®</sup>**

PERITONEAL DIALYSIS SOLUTION

WITH

1.5% DEXTROSE


LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-50521

NDC 49230-206-50

5000 mL

(Approx. 100 mL excess)



**DELFLIX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
**1.5% DEXTROSE**  
 LOW MAGNESIUM    LOW CALCIUM

**CAT. NO. 044-50521** **5000 mL**  
 NDC 49230-206-50 (Approx. 100 mL excess)

---

Each 100 mL contains:

Dextrose, Hydrus, USP	1.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5,5 (5,0 - 6,0)	344 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:


Sodium 132	Magnesium 0.5	Calcium 2.5	Chloride 95	Lactate 40
------------	---------------	-------------	-------------	------------

---

STERILE AND NON PYROGENIC,  
 FOR INTRAPERITONEAL ADMINISTRATION ONLY,  
 USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.  
 Potassium Chloride to be added only under the direction of a physician.  
 LATEX: This product and its packaging does not contain latex materials.  
**CAUTION:** Use only if solution is clear and container is undamaged.  
 Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).  
 Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only DPL-113 PLASTIC 89-909-13 Rev 01/07

  
 Fresenius Medical Care

Fresenius Medical Care NA  
 Waltham, MA 02451  
 1-800-323-5188

**Principal Display Panel - 2.5% Dextrose 2000 mL Bag Label**

**DELFLIX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
 2.5% DEXTROSE  
 LOW MAGNESIUM • LOW CALCIUM  
 CAT. NO. 044-20322  
 NDC 49230-209-23  
 2000 mL/3 Liter Bag  
 (Approx. 40 mL excess)



(01) 003 49230 209 23 5

**DELFLEX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
**2.5% DEXTROSE**  
 LOW MAGNESIUM    LOW CALCIUM

**CAT. NO. 044-20322**                      **2000 mL /3 Liter Bag**  
 NDC 49230-209-23                              (Approx. 40 mL excess)

---

Each 100 mL contains:

Dextrose, Hydrous, USP	2.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	394 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132	Magnesium 0.5	Calcium 2.5	Chloride 95	Lactate 40
------------	---------------	-------------	-------------	------------

---

STERILE AND NON PYROGENIC.  
 FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
 USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.  
 Potassium Chloride to be added only under the direction of a physician.  
 LATEX: This product and its packaging does not contain latex materials.  
**CAUTION:** Use only if solution is clear and container is undamaged.  
 Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).  
 Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only                      DPL-113 PLASTIC    89-909-05 Rev 01/07



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 2.5% Dextrose 3000 mL Bag Label**

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION

WITH

2.5% DEXTROSE

LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-30322

NDC 49230-209-30

3000 mL

(Approx. 60 mL excess)



(01) 003 49230 209 30 3

**DELFLEX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
**2.5% DEXTROSE**  
 LOW MAGNESIUM    LOW CALCIUM

**CAT. NO. 044-30322** **3000 mL**  
 NDC 49230-209-30 (Approx. 60 mL excess)

---

Each 100 mL contains:

Dextrose, Hydrous, USP	2.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	394 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132	Magnesium 0.5	Calcium 2.5	Chloride 95	Lactate 40
------------	---------------	-------------	-------------	------------

---

STERILE AND NON PYROGENIC.  
 FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
 USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.  
 Potassium Chloride to be added only under the direction of a physician.  
 LATEX: This product and its packaging does not contain latex materials.  
 CAUTION: Use only if solution is clear and container is undamaged.  
 Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).  
 Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only DPL-113 PLASTIC 89-909-11 Rev 01/07



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 2.5% Dextrose 5000 mL Bag Label**

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION  
WITH  
2.5% DEXTROSE


LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-50522

NDC 49230-209-50

5000 mL

(Approx. 100 mL excess)



**DELFLIX®**

**PERITONEAL DIALYSIS SOLUTION  
WITH  
2.5% DEXTROSE  
LOW MAGNESIUM    LOW CALCIUM**

**CAT. NO. 044-50522**

**5000 mL**

NDC 49230-209-50

(Approx. 100 mL excess)

Each 100 mL contains:

Dextrose, Hydrus, USP	2.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	394 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132    Magnesium 0.5    Calcium 2.5    Chloride 95    Lactate 40

**STERILE AND NON PYROGENIC.  
FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.**

Potassium Chloride to be added only under the direction of a physician.

LATEX: This product and its packaging does not contain latex materials.

**CAUTION: Use only if solution is clear and container is undamaged.  
Discard any unused portion. Read package insert for full information.**

Recommended Storage: Room Temperature (25°C).

Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only

DPL-113 PLASTIC    89-909-14 Rev 01/07



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 4.25% Dextrose 2000 mL Bag Label**

**DELFLIX®**

**PERITONEAL DIALYSIS SOLUTION  
WITH  
4.25% DEXTROSE**

**LOW MAGNESIUM • LOW CALCIUM**

**CAT. NO. 044-20324**

**NDC 49230-212-23**

**2000 mL/3 Liter Bag  
(Approx. 40 mL excess)**



(01) 003 49230 212 23 5

**DELFLEX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
**4.25% DEXTROSE**  
 LOW MAGNESIUM    LOW CALCIUM

**CAT. NO. 044-20324**                      **2000 mL /3 Liter Bag**  
 NDC 49230-212-23                              (Approx. 40 mL excess)

---

Each 100 mL contains:

Dextrose, Hydrous, USP	4.25 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	483 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132	Magnesium 0.5	Calcium 2.5	Chloride 95	Lactate 40
------------	---------------	-------------	-------------	------------

---

STERILE AND NON PYROGENIC,  
 FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
 USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.  
 Potassium Chloride to be added only under the direction of a physician.  
 LATEX: This product and its packaging does not contain latex materials.  
 CAUTION: Use only if solution is clear and container is undamaged.  
 Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).  
 Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only                      DPL-113 PLASTIC    89-909-06 Rev 01/07



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 4.25% Dextrose 3000 mL Bag Label**

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION  
WITH  
4.25% DEXTROSE

LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-30324

NDC 49230-212-30

3000 mL  
(Approx. 60 mL excess)



(01) 003 49230 212 30 3

**DELFLEX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
**4.25% DEXTROSE**  
 LOW MAGNESIUM    LOW CALCIUM

**CAT. NO. 044-30324** **3000 mL**  
 NDC 49230-212-30 (Approx. 60 mL excess)

---

Each 100 mL contains:

Dextrose, Hydrous, USP	4.25 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	483 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132	Magnesium 0.5	Calcium 2.5	Chloride 95	Lactate 40
------------	---------------	-------------	-------------	------------

---

STERILE AND NON PYROGENIC.  
 FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
 USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.  
 Potassium Chloride to be added only under the direction of a physician.  
 LATEX: This product and its packaging does not contain latex materials.  
**CAUTION:** Use only if solution is clear and container is undamaged.  
 Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).  
 Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only DPL-113 PLASTIC 89-909-12 Rev 01/07



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 4.25% Dextrose 5000 mL Bag Label**

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION

WITH

4.25% DEXTROSE


LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-50524

NDC 49230-212-50

5000 mL

(Approx. 100 mL excess)



(01) 003 49230 212 50 1

**DELFLEX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
**4.25% DEXTROSE**  
 LOW MAGNESIUM    LOW CALCIUM

**CAT. NO. 044-50524** **5000 mL**  
 NDC 49230-212-50 (Approx. 100 mL excess)

---

Each 100 mL contains:

Dextrose, Hydrus, USP	4.25 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	483 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132	Magnesium 0.5	Calcium 2.5	Chloride 95	Lactate 40
------------	---------------	-------------	-------------	------------

---

STERILE AND NON PYROGENIC.  
 FOR INTRAPERITONEAL ADMINISTRATION ONLY.  
 USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.  
 Potassium Chloride to be added only under the direction of a physician.  
 LATEX: This product and its packaging does not contain latex materials.  
**CAUTION:** Use only if solution is clear and container is undamaged.  
 Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).  
 Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only DPL-113 PLASTIC 89-909-15 Rev 01/07



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 2.5% Dextrose 6000 mL Bag Label**

**DELFLEX®**  
 PERITONEAL DIALYSIS SOLUTION  
 WITH  
 2.5% DEXTROSE  
 LOW MAGNESIUM • LOW CALCIUM  
 CAT. NO. 044-60622  
 NDC 49230-209-60  
 6000 mL  
 (Approx. 180 mL excess)



**DELFLIX®**

PERITONEAL DIALYSIS SOLUTION  
WITH  
2.5% DEXTROSE  
LOW MAGNESIUM LOW CALCIUM

CAT. NO. 044-60622 6000 mL

NDC 49230-209-60 (Approx. 180 mL excess)

Each 100 mL contains:

Dextrose, Hydrus, USP	2.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.

pH 5.5 (5.0 - 6.0) 394 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40

STERILE AND NON PYROGENIC.

FOR INTRAPERITONEAL ADMINISTRATION ONLY.

USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.

Potassium Chloride to be added only under the direction of a physician.

LATEX: This product and its packaging does not contain latex materials.

CAUTION: Use only if solution is clear and container is undamaged.  
Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).

Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only

DPL-113 PLASTIC 89-909-62 Rev 05/17



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 4.25% Dextrose 6000 mL Bag Label**

**DELFLIX®**

PERITONEAL DIALYSIS SOLUTION

WITH

4.25% DEXTROSE

LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-60624

NDC 49230-212-60

6000 mL

(Approx. 180 mL excess)

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION  
WITH  
4.25% DEXTROSE  
LOW MAGNESIUM LOW CALCIUM

CAT. NO. 044-60624 6000 mL

NDC 49230-212-60 (Approx. 180 mL excess)

Each 100 mL contains:

Dextrose, Hydrous, USP	4.25 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	483 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40

STERILE AND NON PYROGENIC.

FOR INTRAPERITONEAL ADMINISTRATION ONLY.

USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.

Potassium Chloride to be added only under the direction of a physician.

LATEX: This product and its packaging does not contain latex materials.

CAUTION: Use only if solution is clear and container is undamaged.  
Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).

Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only DPL-113 PLASTIC 89-909-63 Rev 05/17



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 1.5% Dextrose 6000 mL Bag Label**

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION

WITH

1.5% DEXTROSE

LOW MAGNESIUM • LOW CALCIUM

CAT. NO. 044-60621

NDC 49230-206-60

6000 mL

(Approx. 180 mL excess)

**DELFLEX®**

PERITONEAL DIALYSIS SOLUTION  
WITH  
1.5% DEXTROSE  
LOW MAGNESIUM LOW CALCIUM

CAT. NO. 044-60621

6000 mL

NDC 49230-206-60

(Approx. 180 mL excess)

Each 100 mL contains:

Dextrose, Hydrous, USP	1.5 g
Sodium Chloride, USP	538 mg
Sodium Lactate	448 mg
Calcium Chloride, USP	18.4 mg
Magnesium Chloride, USP	5.1 mg
Water for Injection, USP	q.s.
pH 5.5 (5.0 - 6.0)	344 mOsmol/Liter (calculated)

May contain Hydrochloric Acid or Sodium Hydroxide for pH adjustment.

Approximate Milliequivalents Per Liter:

Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40

STERILE AND NON PYROGENIC.

FOR INTRAPERITONEAL ADMINISTRATION ONLY.

USE ASEPTIC TECHNIQUE. DOSAGE AS DIRECTED BY A PHYSICIAN.

Potassium Chloride to be added only under the direction of a physician.

LATEX: This product and its packaging does not contain latex materials.

CAUTION: Use only if solution is clear and container is undamaged.  
Discard any unused portion. Read package insert for full information.

Recommended Storage: Room Temperature (25°C).

Avoid Excessive Heat. Protect From Freezing. See Insert.

Rx only

DPL-113 PLASTIC

89-909-61 Rev 05/17



Fresenius Medical Care

Fresenius Medical Care NA  
Waltham, MA 02451  
1-800-323-5188

**Principal Display Panel - 1.5% Dextrose 3000 mL Bag Label**

**DELFLEX®**

Peritoneal Dialysis Solution  
in Biofine® container

1.5% Dextrose

Low Magnesium • Low Calcium

Cat. No. 077-30321

NDC 49230-206-32

3000 mL

(Approx. 90 mL Excess)





2.5% Dextrose

Low Magnesium • Low Calcium

Cat. No. 077-50622

NDC 49230-209-52

5000 mL

(Approx. 150 mL Excess)

<b>DELFLEX<sup>®</sup></b> Peritoneal Dialysis Solution in Biofine <sup>®</sup> container <b>2.5% Dextrose</b> Low Magnesium • Low Calcium Cat. No. 077-50622 5000 mL NDC 49230-209-52 (Approx. 150 mL Excess)		<ul style="list-style-type: none"><li>• Dosage as directed by physician.</li><li>• Single dose container.</li><li>• Sterile and non-pyrogenic.</li><li>• Use aseptic technique.</li><li>• For intraperitoneal administration only.</li><li>• Potassium chloride to be added only under the direction of a physician.</li><li>• Latex: this product and its packaging does not contain latex materials.</li><li>• Caution: use only if solution is clear and container is undamaged. Discard any unused portion.</li><li>• Recommended storage: store at room temperature (25°C); avoid excessive heat; and protect from freezing.</li><li>• See prescribing information.</li></ul> Rx Only 89-940-21 Rev 08/2022
Each 100 mL Contains: Dextrose, Hydrous, USP 2500 mg Sodium Chloride, USP 538 mg Sodium Lactate, USP 448 mg Calcium Chloride, USP 18.4 mg Magnesium Chloride, USP 5.1 mg Water For Injection, USP q.s.  Approximate milliequivalents per liter: Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40 pH 5.5 (5.0 - 6.0)* 394 mOsmol/Liter (Calculated) <small>*sodium hydroxide or hydrochloric acid may have been used for pH adjustment.</small>		

**Principal Display Panel - 2.5% Dextrose 6000 mL Bag Label**

**DELFLEX<sup>®</sup>**

Peritoneal Dialysis Solution  
in Biofine<sup>®</sup> container

2.5% Dextrose

Low Magnesium • Low Calcium

Cat. No. 077-60622

NDC 49230-209-62

6000 mL

(Approx. 180 mL Excess)

<b>DELFLEX<sup>®</sup></b> Peritoneal Dialysis Solution in Biofine <sup>®</sup> container <b>2.5% Dextrose</b> Low Magnesium • Low Calcium Cat. No. 077-60622 6000 mL NDC 49230-209-62 (Approx. 180 mL Excess)		<ul style="list-style-type: none"><li>• Dosage as directed by physician.</li><li>• Single dose container.</li><li>• Sterile and non-pyrogenic.</li><li>• Use aseptic technique.</li><li>• For intraperitoneal administration only.</li><li>• Potassium chloride to be added only under the direction of a physician.</li><li>• Latex: this product and its packaging does not contain latex materials.</li><li>• Caution: use only if solution is clear and container is undamaged. Discard any unused portion.</li><li>• Recommended storage: store at room temperature (25°C); avoid excessive heat; and protect from freezing.</li><li>• See prescribing information.</li></ul> Rx Only 89-940-14 Rev 08/2022
Each 100 mL Contains: Dextrose, Hydrous, USP 2500 mg Sodium Chloride, USP 538 mg Sodium Lactate, USP 448 mg Calcium Chloride, USP 18.4 mg Magnesium Chloride, USP 5.1 mg Water For Injection, USP q.s.  Approximate milliequivalents per liter: Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40 pH 5.5 (5.0 - 6.0)* 394 mOsmol/Liter (Calculated) <small>*sodium hydroxide or hydrochloric acid may have been used for pH adjustment.</small>		

**Principal Display Panel - 4.25% Dextrose 3000 mL Bag Label**

**DELFLEX®**

Peritoneal Dialysis Solution  
in Biofine® container

4.25% Dextrose

Low Magnesium • Low Calcium

Cat. No. 077-30324

NDC 49230-212-32

3000 mL

(Approx. 90 mL Excess)

<p><b>DELFLEX®</b> Peritoneal Dialysis Solution in Biofine® container <b>4.25% Dextrose</b> Low Magnesium • Low Calcium</p>		<ul style="list-style-type: none"><li>• Dosage as directed by physician.</li><li>• Single dose container.</li><li>• Sterile and non-pyrogenic.</li><li>• Use aseptic technique.</li><li>• For intraperitoneal administration only.</li><li>• Potassium chloride to be added only under the direction of a physician.</li><li>• Latex: this product and its packaging does not contain latex materials.</li><li>• Caution: use only if solution is clear and container is undamaged. Discard any unused portion.</li><li>• Recommended storage: store at room temperature (25°C); avoid excessive heat; and protect from freezing.</li><li>• See prescribing information.</li></ul>												
<p>Cat. No. 077-30324 NDC 49230-212-32</p>	<p>3000 mL (Approx. 90 mL Excess)</p>													
<p>Each 100 mL Contains:</p> <table><tr><td>Dextrose, Hydrous, USP</td><td>4250 mg</td></tr><tr><td>Sodium Chloride, USP</td><td>538 mg</td></tr><tr><td>Sodium Lactate, USP</td><td>448 mg</td></tr><tr><td>Calcium Chloride, USP</td><td>18.4 mg</td></tr><tr><td>Magnesium Chloride, USP</td><td>5.1 mg</td></tr><tr><td>Water For Injection, USP</td><td>q.s.</td></tr></table>		Dextrose, Hydrous, USP	4250 mg	Sodium Chloride, USP	538 mg	Sodium Lactate, USP	448 mg	Calcium Chloride, USP	18.4 mg	Magnesium Chloride, USP	5.1 mg	Water For Injection, USP	q.s.	
Dextrose, Hydrous, USP	4250 mg													
Sodium Chloride, USP	538 mg													
Sodium Lactate, USP	448 mg													
Calcium Chloride, USP	18.4 mg													
Magnesium Chloride, USP	5.1 mg													
Water For Injection, USP	q.s.													
<p>Approximate milliequivalents per liter: Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40</p>														
<p>pH 5.5 (5.0 - 6.0)*                      483 mOsmol/Liter (Calculated) *sodium hydroxide or hydrochloric acid may have been used for pH adjustment.</p>		<p>Rx Only 89-940-12 Rev 08/2022</p>												

**Principal Display Panel - 4.25% Dextrose 5000 mL Bag Label**

**DELFLEX®**

Peritoneal Dialysis Solution  
in Biofine® container

4.25% Dextrose

Low Magnesium • Low Calcium

Cat. No. 077-50624

NDC 49230-212-52

5000 mL

(Approx. 150 mL Excess)

<p><b>DELFLX®</b> Peritoneal Dialysis Solution in Biofine® container</p> <p><b>4.25% Dextrose</b> Low Magnesium • Low Calcium</p>		<ul style="list-style-type: none"> <li>• Dosage as directed by physician.</li> <li>• Single dose container.</li> <li>• Sterile and non-pyrogenic.</li> <li>• Use aseptic technique.</li> <li>• For intraperitoneal administration only.</li> <li>• Potassium chloride to be added only under the direction of a physician.</li> <li>• Latex: this product and its packaging does not contain latex materials.</li> <li>• Caution: use only if solution is clear and container is undamaged. Discard any unused portion.</li> <li>• Recommended storage: store at room temperature (25°C); avoid excessive heat; and protect from freezing.</li> <li>• See prescribing information.</li> </ul>
Cat. No. 077-50624 NDC 49230-212-52	5000 mL (Approx. 150 mL Excess)	
Each 100 mL Contains:		Rx Only 89-940-22 Rev 08/2022
Dextrose, Hydrous, USP Sodium Chloride, USP Sodium Lactate, USP Calcium Chloride, USP Magnesium Chloride, USP Water For Injection, USP	4250 mg 538 mg 448 mg 18.4 mg 5.1 mg q.s.	
Approximate milliequivalents per liter: Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40		
pH 5.5 (5.0 - 6.0)*                      483 mOsmol/Liter (Calculated) <small>*sodium hydroxide or hydrochloric acid may have been used for pH adjustment.</small>		

**Principal Display Panel - 4.25% Dextrose 6000 mL Bag Label**

**DELFLX®**  
Peritoneal Dialysis Solution  
in Biofine® container

4.25% Dextrose

Low Magnesium • Low Calcium

Cat. No. 077-60624

NDC 49230-212-62

6000 mL  
(Approx. 180 mL Excess)

<p><b>DELFLX®</b> Peritoneal Dialysis Solution in Biofine® container</p> <p><b>4.25% Dextrose</b> Low Magnesium • Low Calcium</p>		<ul style="list-style-type: none"> <li>• Dosage as directed by physician.</li> <li>• Single dose container.</li> <li>• Sterile and non-pyrogenic.</li> <li>• Use aseptic technique.</li> <li>• For intraperitoneal administration only.</li> <li>• Potassium chloride to be added only under the direction of a physician.</li> <li>• Latex: this product and its packaging does not contain latex materials.</li> <li>• Caution: use only if solution is clear and container is undamaged. Discard any unused portion.</li> <li>• Recommended storage: store at room temperature (25°C); avoid excessive heat; and protect from freezing.</li> <li>• See prescribing information.</li> </ul>
Cat. No. 077-60624 NDC 49230-212-62	6000 mL (Approx. 180 mL Excess)	
Each 100 mL Contains:		Rx Only 89-940-15 Rev 08/2022
Dextrose, Hydrous, USP Sodium Chloride, USP Sodium Lactate, USP Calcium Chloride, USP Magnesium Chloride, USP Water For Injection, USP	4250 mg 538 mg 448 mg 18.4 mg 5.1 mg q.s.	
Approximate milliequivalents per liter: Sodium 132 Magnesium 0.5 Calcium 2.5 Chloride 95 Lactate 40		
pH 5.5 (5.0 - 6.0)*                      483 mOsmol/Liter (Calculated) <small>*sodium hydroxide or hydrochloric acid may have been used for pH adjustment.</small>		

**DELFLX**

dextrose monohydrate, sodium chloride, sodium lactate, calcium chloride, magnesium chloride solution

Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:49230-206
<b>Route of Administration</b>	INTRAPERITONEAL		



Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SL0G7R00K)		DEXTROSE MONOHYDRATE	1.5 g in 100 mL	
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)		SODIUM CHLORIDE	538 mg in 100 mL	
SODIUM LACTATE (UNII: TU7HW0W0QT) (SODIUM CATION - UNII:LYR4M0NH37, LACTIC ACID - UNII:33X04XA5AT)		SODIUM LACTATE	448 mg in 100 mL	
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)		CALCIUM CHLORIDE	18.4 mg in 100 mL	
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)		MAGNESIUM CHLORIDE	5.08 mg in 100 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0K00R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49230-206-20	6 in 1 CARTON	08/19/1992	
1		2000 mL in 1 BAG; Type 0: Not a Combination Product		
2	NDC:49230-206-30	4 in 1 CARTON	08/19/1992	
2		3000 mL in 1 BAG; Type 0: Not a Combination Product		
3	NDC:49230-206-32	4 in 1 CARTON	08/19/1992	
3		3000 mL in 1 BAG; Type 0: Not a Combination Product		
4	NDC:49230-206-50	2 in 1 CARTON	08/19/1992	
4		5000 mL in 1 BAG; Type 0: Not a Combination Product		
5	NDC:49230-206-60	2 in 1 CARTON	08/19/1992	08/31/2023
5		6000 mL in 1 BAG; Type 0: Not a Combination Product		
6	NDC:49230-206-62	2 in 1 CARTON	08/19/1992	
6		6000 mL in 1 BAG; Type 0: Not a Combination Product		
7	NDC:49230-206-52	2 in 1 CARTON	08/19/1992	
7		5000 mL in 1 BAG; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA020171	08/19/1992		

## DELFLEX

dextrose monohydrate, sodium chloride, sodium lactate, calcium chloride, magnesium chloride solution

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:49230-209
Route of Administration	INTRAPERITONEAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	538 mg in 100 mL
<b>SODIUM LACTATE</b> (UNII: TU7HW0W0QT) (SODIUM CATION - UNII:LYR4M0NH37, LACTIC ACID - UNII:33X04XA5AT)	SODIUM LACTATE	448 mg in 100 mL
<b>CALCIUM CHLORIDE</b> (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	18.4 mg in 100 mL
<b>MAGNESIUM CHLORIDE</b> (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	5.08 mg in 100 mL
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SLOG7R0OK)	DEXTROSE MONOHYDRATE	2.5 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0K00R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49230-209-23	6 in 1 CARTON	08/19/1992	
1		2000 mL in 1 BAG; Type 0: Not a Combination Product		
2	NDC:49230-209-30	4 in 1 CARTON	08/19/1992	
2		3000 mL in 1 BAG; Type 0: Not a Combination Product		
3	NDC:49230-209-32	4 in 1 CARTON	08/19/1992	
3		3000 mL in 1 BAG; Type 0: Not a Combination Product		
4	NDC:49230-209-50	2 in 1 CARTON	08/19/1992	
4		5000 mL in 1 BAG; Type 0: Not a Combination Product		
5	NDC:49230-209-60	2 in 1 CARTON	08/19/1992	08/31/2023
5		6000 mL in 1 BAG; Type 0: Not a Combination Product		
6	NDC:49230-209-62	2 in 1 CARTON	08/19/1992	
6		6000 mL in 1 BAG; Type 0: Not a Combination Product		
7	NDC:49230-209-52	2 in 1 CARTON	08/19/1992	
7		5000 mL in 1 BAG; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020171	08/19/1992	

## DELFLEX

dextrose monohydrate, sodium chloride, sodium lactate, calcium chloride, magnesium chloride solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:49230-212
<b>Route of Administration</b>	INTRAPERITONEAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROSE MONOHYDRATE</b> (UNII: LX22YL083G) (ANHYDROUS DEXTROSE - UNII:5SLOG7R0OK)	DEXTROSE MONOHYDRATE	4.25 g in 100 mL
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	538 mg in 100 mL

<b>SODIUM LACTATE</b> (UNII: TU7HW0W0QT) (SODIUM CATION - UNII:LYR4M0NH37, LACTIC ACID - UNII:33X04XA5AT)	SODIUM LACTATE	448 mg in 100 mL
<b>CALCIUM CHLORIDE</b> (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	18.4 mg in 100 mL
<b>MAGNESIUM CHLORIDE</b> (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	5.08 mg in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0K00R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49230-212-23	6 in 1 CARTON	08/19/1992	
1		2000 mL in 1 BAG; Type 0: Not a Combination Product		
2	NDC:49230-212-30	4 in 1 CARTON	08/19/1992	
2		3000 mL in 1 BAG; Type 0: Not a Combination Product		
3	NDC:49230-212-32	4 in 1 CARTON	08/19/1992	
3		3000 mL in 1 BAG; Type 0: Not a Combination Product		
4	NDC:49230-212-50	2 in 1 CARTON	08/19/1992	
4		5000 mL in 1 BAG; Type 0: Not a Combination Product		
5	NDC:49230-212-60	2 in 1 CARTON	08/19/1992	08/31/2023
5		6000 mL in 1 BAG; Type 0: Not a Combination Product		
6	NDC:49230-212-62	2 in 1 CARTON	08/19/1992	
6		6000 mL in 1 BAG; Type 0: Not a Combination Product		
7	NDC:49230-212-52	2 in 1 CARTON	08/19/1992	
7		5000 mL in 1 BAG; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020171	08/19/1992	

**Labeler** - Fresenius Medical Care North America (958291411)