

**HEADACHE RELIEF- acetaminophen, aspirin and caffeine tablet, coated
Kroger Company**

370R Headache Relief Caplets KROGER 41226 370

DRUG FACTS

ACETAMINOPHEN/ ASPIRIN (NSAID)/ CAFFEINE

Active ingredient (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Pain reliever/Pain reliever aid

Uses

temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- you experience any of the following signs of stomach bleeding:

feel faint

have bloody or black stools

vomit blood

have stomach pain that does not get better

- ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, pregelatinized starch, povidone, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

call 1-800-632-6900



NDC 41226-370-01

Extra Strength

Headache Relief

Acetaminophen 250 mg
Pain Reliever

Aspirin (NSAID) 250 mg
Pain Reliever

Caffeine 65 mg
Pain Reliever Aid

100 CAPLETS

capsule-shaped tablet

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: headache, a cold, arthritis, muscular aches, toothache ■ premenstrual and menstrual cramps

Warnings. See carton for complete warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives ■ facial swelling ■ shock ■ asthma (wheezing)
- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 8 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not use more than directed

- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Questions or comments? 1-800-632-6900

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

OUR BRANDS. OUR GUARANTEE. LOVE IT OR YOUR MONEY BACK. 370F 0321
WWW.KROGER.COM

Lot No.: _____

Exp. Date: _____

UNVARNISHED AREA



HEADACHE RELIEF

acetaminophen, aspirin and caffeine tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-370
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg

Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
MINERAL OIL (UNII: T5L8T28FGP)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE ((Capsule Shaped Tablet))	Size	18mm
Flavor		Imprint Code	TCL;370
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-370-42	1 in 1 CARTON	04/10/2021	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41226-370-01	1 in 1 CARTON	04/10/2021	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/10/2021	

Labeler - Kroger Company (006999528)

Registrant - Time Cap Laboratories, Inc. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
Time Cap Laboratories, Inc.		037052099	manufacture(41226-370)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(41226-370)

Revised: 7/2025

Kroger Company