HEADACHE RELIEF- acetaminophen, aspirin and caffeine tablet, coated Kroger Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

370R Headache Relief Caplets KROGER 41226 370

DRUG FACTS

ACETAMINOPHEN/ ASPIRIN (NSAID)/ CAFFEINE

Active ingredient (in each caplet)

Acetaminophen 250 mg Aspirin 250 mg (NSAID)* Caffeine 65 mg *nonsteroidal anti-inflammatory drug

Pain reliever/Pain reliever aid

Uses

temporarily relieves minor aches and pains due to:

- headache
- •a cold
- arthritis
- muscular aches
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- •shock
- asthma (wheezing)

Allergyalert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- •more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- •3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- •are age 60 or older
- •have had stomach ulcers or bleeding problems
- •take a blood thinning (anticoagulant) or steroid drug
- •take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- •have 3 or more alcoholic drinks every day while using this product
- •take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.

Do not use

- •with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- •for more than 10 days for pain unless directed by a doctor
- •for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- •you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- •taking a prescription drug for diabetes, gout, or arthritis
- •taking any other drug or are under a doctor's care for any serious condition

Stop using and ask a doctor if

- •symptoms do not improve
- •new symptoms occur
- •pain or fever persists or gets worse
- •you experience any of the following signs of stomach bleeding:

feel faint

have bloody or black stools

vomit blood

have stomach pain that does not get better

•ringing in the ears or loss of hearing occurs, consult a doctor before taking any more of this product.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- do not use more than directed
- drink a full glass of water with each dose
- •adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

Other information


- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, hypromellose, microcrystalline cellulose, mineral oil, pregelatinized starch, povidone, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

call 1-800-632-6900





Exp. Date: :.ON 10.1

41260 "01207 SO%"SIZE 12-digit UPC-A (non-suppressed) For Position Only

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF

Drug Facts (continued)

Stomach bleeding warning: This product contains an ISAID, which may cause severe stomach bleeding. The chance is higher if you. ■ are age of or older that war had stomach ulcers or oldering problems take a blood thinning (anticaquilant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ilbuprofen, naproxen, crathen). ■ had a convex alsolated drive the containing crackets.

nonprescription NSAIDs (aspirin, lbuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed Caffeline warning: The recommended dose of this product contains about as much caffeline as a cup of coffee. Limit the use of caffeline-containing medications, foods, or beverages while taking this product because too much caffeline may cause nervousness, irritability, steeplessness, and operationally candib eartheat sleeplessness, and, occasionally, rapid heartbeat

sleeplesshess, and, occasionally, rapid nearureax.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. — for more than 10 days for pain unless directed by a doctor — if or more than 3 days for fever unless directed by a doctor

agys for rever unless directed by a doctor
Ask a doctor before use if a stomach bleeding
warning applies to you you have a history of
stomach problems, such as hearburn you have
high blood pressure, heard disease, liver cirrhosis or
kidney disease you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout, or arthritis ■ taking any other drug or are under a doctor's care for any serious condition Drug Facts (continued)

Stop using and ask a doctor if symptoms do

If pregnant or breast-leeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental

weep and to least in children. In case of accident overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even i you do not notice any signs or symptoms.

Directions and not use more than directed

■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 2 caplets
every 6 hours; not more than 8 caplets in 24 hours
■ children under 12 years: ask a doctor

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients carnauba wax, colloidal silicor dioxide, hypromellose, microcrystalline cellulose,

Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ a cold ■ arthritis
■ muscular aches ■ toothache
■ premenstrual and menstrual cramps

OMPLETE

Purpose

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and yomiting occur,

Drug Facts

Active ingredient (in each caplet)

changes in behavior with nausea and vorniting occur. consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious lilness. Altergy alert. Aspirin may cause a severe allergic reaction, which may include: "

I whese # Table allergic may be a may cause a severe skin reactions. Symptoms may include: "

skin readering billsters in reaction scyring include: "

skin readering billsters in rash if a skin reaction occurs, stop use and seek medical help right away.

* Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take: "

more than

Severe liver damage may occur if you take: I more than 8 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen I or more alcoholic drinks every day while using this product

MADE IN USA FOR MORE PRODUCT INFORMATION, SCAN UPC USING YOUR KROGER APP OR CALL 800-632-6900 NATI, OHIO 45202 8

Questions or comments? 1-800-632-6900 d starch, sodium ım dioxide

Drug Facts (continued)

NDC 41226-370-42 Kroger **Extra Strength Headache Relief** Acetaminophen 250 mg Pain Reliever Aspirin (NSAID) 250 mg

Pain Reliever Caffeine 65 mg Pain Reliever Aid

> **24** CAPLETS capsule-shaped tablet

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION
Uses = temporarily relieves minor aches and pains due to: headache, a
ood, arthritis, muscular aches, toothache = premenstrual and menstrual
camps
Warnings See carton for complete warnings

Warnings See carton for complete warnings
Reye's syndrome. Children and renages who that on are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in betravior with nausea and vomiting occur, consult a doctor becauses these symptoms should not use this product. When using this product, if changes in betravior with nausea and vomiting occur, consult a doctor becauses these symptoms could be an early sign of Reye's syndrome, a rate but serious illness. Allergy alert. Aepirim may cause a severe allergic reaction, which may include: a livines a lactal swelling severe skin reactions occurs, stop use and seek medical help right away. Liver varning. This product conditians and serior in capacity in the maximum daily amount a with other drugs containing acetaminophen. Severe skin reaction occurs, stop use and seek medical help right away. Liver varning. This product contains and this sing this product. Stomach theeding warning. This product contains a mids. Wish. Which may cause severe stomach bleeding. The chance is higher if your. are age 60 or older. The about the maximum daily contained to the same other drugs containing acetaminophen. Severe stomach bleeding. The chance is higher if your. are age 60 or older. The about the maximum daily contained the distribution to the saving may brobuct a may be obtained the serior drugs of the separately important not to use aspiring the last 3 months of preparancy unless definitely directed to do so by a doctor because it may especially important not to use aspiring unting the last 3 months of programs. The may be approxed to the separately important not to use aspiring until gains of water proper the depotence any signs or symptoms. The program of one of the depotence of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. You do not notice any signs or symptoms.

with each dose ■adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours ■ children under 12

CHARLES AGRICATE THE SHOULD IN THE PRODUCT IF THE MPRINTED FOL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, THE IMPRINTED FOLL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROCKEN OR MISSING — store at 25°C (57°F); securions permitted between 15°-30°C (59°-86°F) — ares by expiration date on package Questions or comments? 1-800-632-6900

DIST, BY THE KROGER CO., CINCINNATI, OHIO 45202

370R 0321

Date: Lot No.:



HEADACHE RELIEF

acetaminophen, aspirin and caffeine tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-370
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	250 mg	

Inactive Ingredients				
Ingredient Name	Strength			
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
STARCH, CORN (UNII: O8232NY3SJ)				
POVIDONE (UNII: FZ989GH94E)				
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)				
CARNAUBA WAX (UNII: R12CBM0EIZ)				
MINERAL OIL (UNII: T5L8T28FGP)				

Product Characteristics			
Color	white	Score	no score
Shape	CAPSULE ((Capsule Shaped Tablet))	Size	18mm
Flavor		Imprint Code	TCL;370
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:41226-370- 42	1 in 1 CARTON	04/10/2021		
1		24 in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:41226-370- 01	1 in 1 CARTON	04/10/2021		
2		100 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/10/2021	

Labeler - Kroger Company (006999528)

Registrant - Time Cap Laboratories, Inc. (037052099)

Establishment				
Name	Address	ID/FEI	Business Operations	
Time Cap Laboratories, Inc.		037052099	manufacture(41226-370)	

Revised: 4/2021 Kroger Company