

**IBUPROFEN- ibuprofen tablet, coated**  
**Kroger Company**

-----  
**Kroger Ibuprofen Tablets 200 mg (Brown Round)**

**DRUG FACTS**

***Active ingredient (in each tablet)***

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/Fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:  
backache  
headache  
menstrual cramps  
minor pain of arthritis  
muscular aches  
the common cold  
toothache
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening.

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of the body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
  - the smallest effective dose should be used
- adults and children 12 years and over:**

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

**children under 12 years:**

- ask a doctor

**OTHER INFORMATION**

**Other information**

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

**Questions or comments? 1-800-632-6900**



NDC 41226-601-53

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS

# Ibuprofen

TABLETS USP, 200 mg

**PAIN RELIEVER/  
FEVER REDUCER  
(NSAID)\***

ACTUAL SIZE  
**500 TABLETS**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)**  
Ibuprofen 200 mg (NSAID)\* .....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued under label)**

\*Advil is a registered trademark of Haleson US Holdings LLC. Haleson US Holdings LLC is not affiliated with The Kroger Co. or this product.

**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**

**MADE IN INDIA**

For More Product Information,  
Scan UPC Using Your Kroger App  
or Call 800-632-6900  
**OUR PHARMACIST RECOMMENDED**  
601R 0284



0 41260 02171 8

**LOT:** \_\_\_\_\_ **EXP:** \_\_\_\_\_

Varnish Omit Area

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**  
■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist.
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

**Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments? Call 1-800-632-6900**

Adhesive Area



COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS\*

# Ibuprofen

Tablets USP, 200 mg

**PAIN RELIEVER/  
 FEVER REDUCER  
 (NSAID)\***

**250 COATED TABLETS**

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

### Drug Facts

**Active ingredient (in each tablet)**  
 Ibuprofen 200 mg (NSAID)\*  
 \*nonsteroidal anti-inflammatory drug

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
 ■ skin redness ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Drug Facts (continued under label)

\*Advil® is a registered trademark of Haleon US Holdings LLC.  
**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**  
**MADE IN INDIA**

### YOUR PHARMACIST RECOMMENDED

Our Brands. Our Guarantee. Love It or Your Money Back.  
 www.kroger.com



601R 0624

LOT:

EXP.:

**Varnish Omit Area**



PEEL HERE

### Drug Facts (continued)

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### Other Information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

**Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** Call 1-800-632-6900

Adhesive Area



COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS\*

# Ibuprofen

Tablets USP,  
200 mg

**PAIN RELIEVER/  
FEVER REDUCER  
(NSAID)\***

**200 COATED TABLETS**

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

### Drug Facts

**Active ingredient (in each tablet)** Ibuprofen 200 mg (NSAID)\*  
 \*nonsteroidal anti-inflammatory drug

**Uses**  
 Temporarily relieves minor aches and pains due to:  
 ■ headache ■ toothache ■ backache ■ menstrual cramps  
 ■ the common cold ■ muscular aches ■ minor pain of arthritis  
 ■ temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ stock  
 ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Drug Facts (continued under label)

\*Advil® is a registered trademark of Haleon US Holdings LLC, DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA

OUR PHARMACIST RECOMMENDED

Our Brands. Our Guarantee. Love It or Your Money Back.

www.kroger.com



LOT:  
EXP:

Varnish Omit Area



Adhesive Area

**Questions or comments?** Call 1-800-632-6900

**Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Other information** ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used

■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

■ children under 12 years: ask a doctor

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**any new symptoms appear.**

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present in the painful area

■ dizziness or lightheadedness

■ blurred vision

■ ringing in ears

■ trouble breathing

■ chest pain

■ weakness in one part or side of body

■ slurred speech

■ stomach pain that does not get better

■ you have symptoms of heart problems or stroke:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ take with food or milk if stomach upset occurs

**When using this product**

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

■ you experience any of the following signs of stomach bleeding:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ take with food or milk if stomach upset occurs

**Ask a doctor or pharmacist before use if you are**

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

**Do not use**

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ right before or after heart surgery

**Ask a doctor before use if**

■ stomach bleeding warning applies to you

■ you have problems or serious side effects from taking pain relievers or fever reducers

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

■ you are taking a diuretic

**Other information**

■ read all warnings and directions before use

■ store at 20-25°C (68-77°F)

**Questions or comments?** Call 1-800-632-6900

**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**

**MADE IN INDIA**

Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com

601R 0624

LOT:

EXP:

Varnish Omit Area

# Ibuprofen

Tablets USP, 200 mg

**PAIN RELIEVER/  
FEVER REDUCER  
(NSAID)**

**100 COATED TABLETS**

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses**  
 Temporarily relieves minor aches and pains due to:  
 ■ headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ stock  
 ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use, and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Other information** ■ read all warnings and directions before use, ■ store at 20-25°C (68-77°F)

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used

■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

■ children under 12 years: ask a doctor

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**any new symptoms appear.**

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present in the painful area

■ dizziness or lightheadedness

■ blurred vision

■ ringing in ears

■ trouble breathing

■ chest pain

■ weakness in one part or side of body

■ slurred speech

■ stomach pain that does not get better

■ you have symptoms of heart problems or stroke:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ take with food or milk if stomach upset occurs

**When using this product**

■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

■ you experience any of the following signs of stomach bleeding:

■ feel faint

■ vomit blood

■ have bloody or black stools

■ take with food or milk if stomach upset occurs

**Ask a doctor or pharmacist before use if you are**

■ under a doctor's care for any serious condition

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

■ taking any other drug

**Do not use**

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ right before or after heart surgery

**Ask a doctor before use if**

■ stomach bleeding warning applies to you

■ you have problems or serious side effects from taking pain relievers or fever reducers

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

■ you are taking a diuretic

**Other information**

■ read all warnings and directions before use, ■ store at 20-25°C (68-77°F)

**Questions or comments?** Call 1-800-632-6900

**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**

**MADE IN INDIA**

Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com

601R 0624

LOT:

EXP:

Varnish Omit Area





NDC 41226-601-42

# Ibuprofen

Tablets USP, 200 mg

**PAIN  
RELIEVER/  
FEVER  
REDUCER  
(NSAID)**

**24  
COATED  
TABLETS**

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses** ■ temporarily relieves minor aches and pains due to headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

**Directions** ■ do not take more than directed ■ the smallest effective use should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

**Other information** ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

**Questions or comments?** Call 1-800-632-6900

**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**

Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com

**MADE IN INDIA**

601R 0524 LOT:

EXP:

**Varnish  
Omit Area**

E826R

# Varnish/Print Omit Area

NDC 41226-601-42



COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS\*

# Ibuprofen

Tablets USP, 200 mg

**PAIN RELIEVER/  
FEVER REDUCER (NSAID)\***

**24 COATED  
TABLETS**

ACTUAL  
SIZE

Varnish  
Omit

Varnish  
Omit

Front of Package

Bottom of Package

### RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

#### Drug Facts

**Active ingredient (in each tablet)** Purpose  
Ibuprofen 200 mg (NSAID)\* Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Uses

- Temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

#### Warnings

- Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
  - hives
  - facial swelling
  - asthma (wheezing)
  - stomach
  - skin redness/itching
  - rash
  - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
  - are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 2 or more alcoholic drinks every day while using this product
  - take more or for a longer time than directed

#### Drug Facts (continued)

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk, if stomach upset occurs
- do not take more than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - stomach pain
  - vomit blood
  - black stools
  - stools that look like coffee grounds
  - you have symptoms of heart problems or stroke: chest pain, trouble breathing

**Drug Facts (continued)**

- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

**Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium

**Drug Facts (continued)**

- stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?**

Call 1-800-532-6900

LOT: EXP:

Varnish Omit

Back of Package

Top of Package

142818103

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

OTC drug facts and directions are provided as a service. Read the printed instructions. Keep this and the other printed instructions for later reference.

**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**

**MADE IN INDIA**

**OUR PHARMACIST RECOMMENDED Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com**

Side of Package

Side of Package



Varnish Omit

NDC 41226-601-53



COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS

# Ibuprofen

TABLETS USP, 200 mg

**PAIN RELIEVER/FEVER REDUCER (NSAID)\***

ACTUAL SIZE

**500 TABLETS**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Ibuprofen 200 mg (NSAID)\* ..... Pain reliever/Fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- stomach
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued under label)**

Advil is a registered trademark of Haleon US Holdings LLC. Haleon US Holdings LLC is not affiliated with The Kroger Co. or this product.

**DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202**

**MADE IN INDIA**

For More Product Information, Scan UPC Using Your Kroger App or Call 1-800-632-6900

**YOUR PHARMACIST RECOMMENDED**

LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

SEE HERE

Varnish Omit Area

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use:** if you have ever had an allergic reaction to any other pain reliever/fever reducer; right before or after heart surgery.

**Ask a doctor before use if:** stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers or fever reducers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; you are taking a diuretic.

**Ask a doctor or pharmacist before use if you are:** under a doctor's care for any serious condition; taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin; taking any other drug.

**When using this product:** take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if:** you experience any of the following signs of stomach bleeding:
 

- red or black stools
- stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

**Inactive ingredients:** colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** Call 1-800-632-6900

Adhesive Area

## IBUPROFEN

ibuprofen tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-601
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients	
Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6130)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND (biconvex film coated tablets)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL;114
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-601-42	1 in 1 CARTON	04/10/2021	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41226-601-05	1 in 1 CARTON	06/12/2021	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41226-601-01	1 in 1 CARTON	06/12/2021	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:41226-601-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2021	
5	NDC:41226-601-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2021	
6	NDC:41226-601-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	04/15/2024	
7	NDC:41226-601-10	2 in 1 PACKAGE, COMBINATION	04/15/2024	
7	NDC:41226-601-53	500 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/10/2021	

**Labeler** - Kroger Company (006999528)

**Registrant** - Time Cap Laboratories, Inc. (037052099)

## Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(41226-601)

Revised: 9/2025

Kroger Company