

ASPIRIN- aspirin tablet, coated
Kroger Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

481R Kroger 41226-481 Aspirin 81 mg Delayed-Release Tablets

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problem
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

feel faint

have bloody or black stools

vomit blood

have stomach pain that does not get better

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

OTHER INFORMATION**Other information**


- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80,

simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-632-6900

COMPARE TO the active ingredient in ASPIRIN
REGIMEN BAYER® LOW DOSE †See side panel



Aspirin 81 mg
Pain Reliever (NSAID)
Aspirin Regimen**

500 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts	Active ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID)*.....Pain reliever *nonsteroidal anti-inflammatory drug
Uses	<ul style="list-style-type: none"> ■ temporarily relieves minor aches and pains Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin
Warnings	<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction, which may include: <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) </p>
Drug Facts (continued under label)	
<p>†Bayer® Low Dose Aspirin is a registered trademark of Bayer AG, Pittsburgh, PA 15205. Bayer AG is not affiliated with The Kroger Co. or this product.</p> <p>DISTRIBUTED BY THE KROGER CO. MADE IN USA CINCINNATI, OHIO 45202</p> <p>FOR MORE PRODUCT INFORMATION, SCAN UPC USING YOUR KROGER APP OR CALL 800-632-6900</p> <p>Our Brands. Our Guarantee. Low it or Your Money Back. www.kroger.com 481R 0321</p>	

Lot No.: 0 41260 01256 3

Exp Date:

SEE THE FREE

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product.

■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have not been drinking fluids
- you have lost a lot of fluid due to vomiting or diarrhea

Ask a doctor/pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments?
1-800-632-6900

Exp Date:
Lot No.:
0 41260 01296 9
90% SIZE
121 FPO
For Position Only
Using Your Kroger App or Call 800-632-6900
For More Product Information, Scan UPC
DISTR. BY THE KROGER CO., CINCINNATI, OHIO 45202
With The Kroger Co. or this product. © 2015 Kroger Co.
Bayer® Low Dose Aspirin is a registered trademark of Bayer AG, Pittsburgh, PA 15205. Bayer AG is not affiliated with The Kroger Co. or this product. 4819 021

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient (in each tablet) Purpose
Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses
■ temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older

Drug Facts (continued)
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:

Drug Facts (continued)
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information
■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package
■ Avoid excessive heat above 40°C (104°F).

Drug Facts (continued)
Inactive ingredients anhydrous lactose, croscarmellose sodium, croscarmellose sodium, D&G yellow #10, silicon dioxide, iron oxide color, hydroxypropyl methylcellulose, polyethylene glycol, hydroxypropyl methylcellulose, polyethylene glycol, sodium hydroxide, sodium phosphate dibasic, sodium phosphate monobasic, titanium dioxide, triethyl citrate

WARNING: RIBBON DO NOT USE THIS PRODUCT IF THE AMBERED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

NDC 41226-481-12

COMPARE TO THE ACTIVE INGREDIENT IN BAYER® LOW DOSE ASPIRIN (See side panel)

OUR PHARMACISTS RECOMMEND

Safety Coated Aspirin Regimen

120 ENTERIC COATED TABLETS

actual size

81 mg

Low Dose Aspirin

81 mg Delayed-Release Tablets / Pain Reliever (NSAID)

Kroger

E826R



NDC 41226-481-12

Low Dose

81 mg

Aspirin

81 mg Delayed-Release Tablets

Pain Reliever (NSAID)

Safety Coated Aspirin Regimen

120 ENTERIC COATED TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Uses ■ Temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away, 1-800-222-1222

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ Avoid excessive heat above 40°C (104°F). **Questions? 1-800-632-6900**

DIST. BY THE KROGER CO., CINCINNATI, OHIO 45202
481R 0421

Lot No.:

Exp. Date:



Top of Package
READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THE BOX FOR IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each tablet) Purpose
 Aspirin 81 mg (NSAID*) Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses
 ■ temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
 ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
 ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Back of Package

Exp Date: _____
 Lot No.: 4 1260 01297 6
 TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOL SEAL IS CUT, TORN, BROKEN OR MISSING

Drug Facts (continued)

■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important

Bottom of Package

Drug Facts (continued)

not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: consult a doctor

Other information
 ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
 ■ use by expiration date on package
 ■ Avoid excessive heat above 40°C (104°F).

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-632-6900

†Bayer® Low Dose Aspirin is a registered trademark of Bayer AG, Pittsburgh, PA 15205. Bayer AG is not affiliated with The Kroger Co. or this product. 481R 0421

DISTRIBUTED BY THE KROGER CO.
 CINCINNATI, OHIO 45202
FOR MORE PRODUCT INFORMATION, SCAN UPC USING YOUR KROGER APP OR CALL 800-632-6900

Front of Package

NDC 41226-481-20

OUR PHARMACEUTICAL RECOMMENDED

Safety Coated Aspirin Regimen

actual size

200 ENTERIC COATED TABLETS

COMPARE TO the active ingredient in BAYER® LOW DOSE ASPIRIN (See top panel)

Kroger

Low Dose 81 mg

Aspirin

81 mg Delayed-Release Tablets / Pain Reliever (NSAID)

E827R1

NDC 41226-481-20



Low Dose **81 mg**

Aspirin

81 mg Delayed-Release Tablets
Pain Reliever (NSAID)

Safety Coated Aspirin Regimen

200 ENTERIC COATED TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION!

Uses ■ temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
 ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
 ■ 1-800-222-1222 **Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ Avoid excessive heat above 40°C (104°F). **Questions? 1-800-632-6900**

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

Lot No.: 481R
 Exp. Date: 0421

NDC 41226-481-31



Low Dose **81 mg**



Aspirin

81 mg Delayed-Release Tablets / Pain Reliever (NSAID)

Safety Coated Aspirin Regimen

300 ENTERIC COATED TABLETS

actual size

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts
Active ingredient (in each tablet) Purpose
 Aspirin 81 mg (NSAID) Pain reliever
 nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
 ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued under label)

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ Avoid excessive heat above 40°C (104°F). **Questions? 1-800-632-6900**

DIST. BY THE KROGER CO., CINCINNATI, OHIO 45202

FOR MORE PRODUCT INFORMATION, SCAN UPC USING YOUR KROGER APP OR CALL 800-632-6900

LOT #: 041260
 EXP. DATE: 01295

SEE HERE

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver/cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis ■ blood thinners

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ ever gets worse or lasts more than 3 days ■ fevers or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ Avoid excessive heat above 40°C (104°F).

Inactive ingredients anhydrous lactose, carnitine wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly/sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-632-6900

Drug Facts (continued)

Do not use
 ■ if you are allergic to aspirin or any other pain reliever/fever reducer
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver/cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis ■ blood thinners

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ ever gets worse or lasts more than 3 days ■ fevers or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ Avoid excessive heat above 40°C (104°F).

Inactive ingredients anhydrous lactose, carnitine wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly/sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-632-6900

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-481
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
BROWN IRON OXIDE (UNII: 1N032N7MFO)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	HEART
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-481-12	1 in 1 CARTON	06/12/2021	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41226-481-20	1 in 1 CARTON	06/12/2021	
3		200 in 1 BOTTLE; Type 0: Not a Combination		

4		Product		
3	NDC:41226-481-31	300 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2021	
4	NDC:41226-481-50	1 in 1 BOTTLE	04/10/2021	
4		500 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	04/10/2021	

Labeler - Kroger Company (006999528)

Registrant - Time Cap Laboratories, Inc. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
Time Cap Laboratories, Inc.		037052099	manufacture(41226-481)

Revised: 6/2021

Kroger Company