

ASPIRIN- aspirin tablet, coated
Kroger Company

481R Kroger 41226-481 Aspirin 81 mg Delayed-Release Tablets

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problem
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

OTHER INFORMATION

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-632-6900



NDC 41226-481-50

COMPARE TO THE ACTIVE INGREDIENT
IN ASPIRIN REGIMEN BAYER® LOW DOSE¹

LOW DOSE
Aspirin

81 mg Delayed-Release Tablets
Safety Coated
Aspirin Regimen[†]

ACTUAL
SIZE

PAIN RELIEVER (NSAID)*

500 ENTERIC COATED TABLETS

Talk to your doctor or other healthcare provider before using this product for your heart.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 81 mg (NSAID)[†] Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product

Drug Facts (continued under label)
¹Bayer® is a registered trademark of Bayer Aktiengesellschaft
DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202
481R 0624
YOUR PHARMACIST RECOMMENDED
Our Brands. Our Guarantee.
Love It or Your Money Back.
www.kroger.com

LOT: 41260101256
EXP: 03

Varnish Omit Area

PEEL HERE

Adhesive Area

Drug Facts (continued)

■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, low blood counts, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information
■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package
■ avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly sorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? Call 1-800-632-6900



NDC 41226-481-20

COMPARE TO THE ACTIVE INGREDIENT
IN ASPIRIN REGIMEN BAYER® LOW DOSE¹

LOW DOSE
Aspirin

81 mg DELAYED-RELEASE TABLETS
SAFETY COATED
ASPIRIN REGIMEN[†]

ACTUAL
SIZE

PAIN RELIEVER (NSAID)

200 ENTERIC COATED TABLETS

Talk to your doctor or other healthcare provider before using this product for your heart.

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package
■ avoid excessive heat above 40°C (104°F)


Questions or comments? Call 1-800-632-6900

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202
Our Brands. Our Guarantee.
Love It or Your Money Back. www.kroger.com

481R 0524

LOT: 0524
EXP: 03

Varnish Omit Area




NDC 41226-481-12

LOW DOSE

Aspirin

81 mg DELAYED-RELEASE TABLETS
SAFETY COATED
ASPIRIN REGIMEN[†]

120

ENTERIC COATED TABLETS

PAIN RELIEVER (NSAID)

Talk to your doctor or other healthcare provider before using this product for your heart.

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product

Directions ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

Other information ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ avoid excessive heat above 40°C (104°F)

Questions or comments? Call 1-800-632-6900

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com

481R 0524 LOT: EXP:

Varnish Omit Area

Coating/Print Omit Area



Coating/Print Omit Area

Top of Package

Bottom of Package

Coating Omit

Front of Package

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet) Purpose Aspirin 81 mg (NSAID)* Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Drug Facts (continued)

take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease you are taking a diuretic you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

gout diabetes arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away. you experience any of the following signs of stomach bleeding: feel faint vomit blood

Drug Facts (continued)

have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

drink a full glass of water with each dose adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor children under 12 years: consult a doctor

Other information

store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) use by expiration date on package avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide

NDC 41226-48-12



COMPARE TO THE ACTIVE INGREDIENT IN ASPIRIN REGIMEN BAYER® LOW DOSE

LOW DOSE Aspirin 81 mg Delayed-Release Tablets Safety Coated Aspirin Regimen†



ACTUAL SIZE

PAIN RELIEVER [NSAID] 120 ENTERIC COATED TABLETS

Coating/Print Omit Area

Coating Omit

Drug Facts (continued)

oic acid, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyvinylpyrrolidone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments?

Call 1-800-932-9300

IMPROVED SEAL: DO NOT USE THIS PRODUCT IF THE IMPROVED SEAL IS CRACKED, TORN, BROKEN OR MISSING. DO NOT USE THE PRODUCT IF THE SEAL IS CRACKED, TORN, BROKEN OR MISSING.

Take only your doctor or other healthcare provider's advice using the product for your heart.

COMPARE TO the active ingredient in BAYER® LOW DOSE ASPIRIN †See side panel

NDC 41226-481-31

Kroger

Low Dose **81 mg**

Aspirin
81 mg Delayed-Release Tablets / Pain Reliever (NSAID)

Safety Coated Aspirin Regimen

300 ENTERIC COATED TABLETS

actual size

Pharmacist Recommended

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 81 mg (NSAID)* Pain reliever
*nonsteroidal anti-inflammatory drug

Uses ■ Temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Drug Facts (continued under label)

†Bayer® Low Dose Aspirin is a registered trademark of Bayer AG, Pittsburgh, PA 15205. Bayer AG is not affiliated with The Kroger Co. or this product. 481R 0421

DIST. BY THE KROGER CO., CINCINNATI, OHIO 45202

FOR MORE PRODUCT INFORMATION, SCAN UPC USING YOUR KROGER APP OR CALL 800-632-6900

80% SIZE 12-digit UPC-A For Position Only

0 4 1260 01295 2

LOT #: EXP. DATE:

PEEL HERE

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- ever gets worse or lasts more than 3 days
- restlessness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away: 1-800-222-1222

Directions ■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ Avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose; carnauuba wax; colloidal silicon dioxide; croscarmellose sodium; D&C yellow #10 aluminum lake; iron oxide (red); methacrylic acid and ethyl acrylate copolymer; microcrystalline cellulose; polyсорate 80; simethicone sodium hydroxide; sodium lauryl sulfate; starch; talc; titanium dioxide; triethyl citrate

Questions or comments? 1-800-632-6900

Drug Facts (continued)

ASPIRIN aspirin tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-481
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	81 mg
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			

DIMETHICONE (UNII: 92RU3N3Y1O)
SODIUM HYDROXIDE (UNII: 55X04QC32I)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)
BROWN IRON OXIDE (UNII: 1N032N7MFO)
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)
POLYSORBATE 80 (UNII: 6OZP39ZG8H)
STARCH, CORN (UNII: O8232NY3SJ)
TALC (UNII: 7SEV7J4R1U)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	HEART
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-481-12	1 in 1 CARTON	06/12/2021	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41226-481-20	1 in 1 CARTON	06/12/2021	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41226-481-31	300 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2021	
4	NDC:41226-481-50	1 in 1 BOTTLE	04/10/2021	
4		500 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/10/2021	

Labeler - Kroger Company (006999528)

Registrant - Time Cap Laboratories, Inc. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
Time Cap Laboratories, Inc.		037052099	manufacture(41226-481)

Revised: 9/2025

Kroger Company