# ACETAMINOPHEN- acetaminophen tablet, coated Kroger Company

-----

#### Kroger Extra Strength Acetaminophen Rapid Release Tablets

# Drug Facts

Acetaminophen 500 mg

Acetaminophen 500 mg

Pain reliever/fever reducer

#### Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- •more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- •3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- •skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you are allergic to acetaminophen or any of the inactive ingredients in this product

### Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

### Stop use and ask a doctor if

pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days

### If pregnant or breast-feeding, ask a health professional before use.

# Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

# • do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

ask a doctor

### **OTHER INFORMATION**


• store between 20-25°C (68-77°F)


*Inactive ingredients* croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, mica based pearlescent pigment, polyethylene glycol 400, polyethylene glycol 6000, polysorbate 80, povidone, pregelatinized starch

### Questions or comments? Call 1-800-632-6900



Brug Facts (continued)         If You are a largic to acclaminophen or any of the inactive ingredients in this product         Ask a adoctor before use if you have liver disease         Ask a adoctor or pharmacist before use if you are taking the blood thining drug warfarin         Spouse and ask a doctor if speak or sould be signs of a serious condition.         If perguant to breast-feeding, ask a health professional before use. Neeroose warming: In case of overdose, get medical help or contact a peison Cartor O ciniter right away (1 - 600-222-1222), Ouick medical aretion is critical for adults are of the series of the series of the series of onto take more than directed. See overdose warming.         Interctions and over       Inter a large series and the other on take more than 0 captes in 24 hours, unless directed by doctor         Interctive ingredients cross-armelises solum, crospovidone, topodentive eignoid 400, polyethylere glycul 6000, polyesorbate 80, povidone, preparisine stearte, mica tased partices ent glyment, povidone, preparisine stearte, mica tased partices ent 90, opolyethylene glycul 6000, polyesorbate 80, povidone, preparisine stearte, mica tased partices ent 90, povidone, preparisine stearte, mica tased partises ent 90, povidone, preparisine stearte,		Using Your Kro	Call 1-800-632-6900	Inactive i FD&C red #40 hypromellose polyethylene i povidone, pre	Other info ■ store between the between	children under 12 years	adults and children 12 years and over	Do not take m	If pregnant o Keep out of r Overdose wa Poison Contro attention is cr notice any sig	Stop use and pain gets w fever gets v new sympt redness or These could t	Ask a doctor before u Ask a doctor or phari thinning drug warfarin	in this product	Drug Earte
	Adhesive Area	or More Product Information, Scan UPC Ising Your Kroger App or Call 800-632-6600	s or comments? 32-6900	Inactive ingredients croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, mica based pearlescent pigment, polyethylene glycol 400, polyethylene glycol 6000, polysorbate 80, povidone, pregelatinized starch	<b>Other information</b> I store between 20-25°C (68-77°F)	ask a doctor	take 2 caplets every 6 hours while symptoms unloss directed by a doctor to not take more than 6 caplets in 24 hours, and a stretced by a doctor to not use for more than 10 days unless direct by a doctor	Directions Do not take more than directed. See overdose warning.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Junck medical Poison Control Center right away (1-800-222-1222). Junck medical the statement of the statement of the statement of the statement of the statement attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Stop use and ask a doctor if pain gets worse or lasts more than 10 days lever gets worse or lasts more than 3 days new symptons occur express or swelling is present These could be signs of a serious condition.	use if you have liver disease macist before use if you are taking the	If you are allerigic to acetaminophen or any of the inactive ingredients in this product	(continued)







Directions at on ot take more than directed (see overdose warning) a adults and children 12 years and over: take 2 capties usery 6 hours, unless with sart, on on take more than 6 captes in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor con tuse for more than 10 years: ask a doctor If a skin reaction occurs, stop use and seek medical help right away. If pregnant or breast-freeding, ask a health professional before use. Reep out of reach of children. Overdose warning: In case of overdose, get medical help or orienta et Polson Control Centra (Palson 222: 1222). Outich macka attention is critical for autily as well as for children even if you do not notice any signs or symptoms. RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION Other information **—** Tamper Evident: do not use this product if the imprinted foll seal over the mouth of the bottle is cut, torn, broken or Missing **—** store between 20-25°C (89-77\*F) 697R 0321 USES m temporarity relieves minor aches and pains due to: m headache m the common cold m backache m muscular aches m minor pain of arthritis m toothache m muscular aches Warnings See carlon for complete warnings Liver warning: This product contains acetanimophen. Severe liver damage may occur if you take: a more than 4,000 mg of acetanimophen in 24 hours a with other drugs containing acetaminophen a 3 or more alcoholic drinks every day while DISTRIBUTED BY THE KROGER CO. CINCINNATI, OHIO 45202 MADE IN INDIA COMMON COMMONSIIS DUR Brands. OUR GUARTHED. COVE HT OT YOU'T MONEY Back. premenstrual and menstrual cramps temporarily Questions or comments? 1-800-632-6900 for adults Extra Strength Acetaminophen www.kroger.com Pain Reliever/ 24 RAPID RELEASE CAPLETS 500 mg EACH **Fever Reducer** Date: reduces fever Lot No.: Б.

500 mg

Fast Relief

A	CETAMINO	PHEN								
эс	etaminophen t	ablet, coatec	l							
Ρ	roduct Infor	mation								
Ρ	roduct Type		NDC:412	26-697						
R	oute of Admini	duct Type     HUMAN OTC DRUG     Item Code (Source)     NDC:41226-697       Ite of Administration     ORAL     ORAL     ORAL								
A	ctive Ingredi	ent/Active	Moiety							
Ingredient Name Basis of Stre								Strengt		
AC	ETAMINOPHEN	(UNII: 362O9ITL	9D) (ACETAMINOPHEN - UN	II:362O9IT	<sup>-</sup> L9D)	ACETAMINOP	HEN	500 mg		
Ir	active Ingre	dients								
Ingredient Name								Strength		
PC	OVIDONE K30 (U	NII: U725QWY32	X)							
S٦	ARCH, CORN (UI	NII: 08232NY35	)							
	ICA (UNII: V8A1AW									
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)										
	<b>D&amp;C RED NO. 40</b>									
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)										
			(UNII: 0WZ 8WG20P6)							
	AGNESIUM STEA	-	·							
	OLYSORBATE 80									
CROSPOVIDONE (UNII: 2S7830E561)										
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE) CROSCARMELLOSE SODIUM (UNII: M280L1HH48)										
CI	KUSCARMELLUS		1: M280L1HH48)							
Ρ	roduct Chara	acteristics								
	olor	red			Score		no	score		
	nape				Size		mm			
	avor							L;A71		
	ontains					couc		_,		
P	ackaging									
#	ltem Code	Pac	kage Description			ing Start ate		ting End Date		
1	NDC:41226-697- 42	1 in 1 CARTON		04	4/10/2021					
1		24 in 1 BOTTL Product	E; Type 0: Not a Combinati	on						
2	34	225 in 1 BOTT Product	LE; Type 0: Not a Combina	tion 04	4/15/2024					
3	NDC:41226-697- 01	1 in 1 CARTON			4/15/2024					
3		100 in 1 BOTT Product	LE; Type 0: Not a Combina	tion						

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC Monograph Drug	M013	04/10/2021					

Labeler - Kroger Company (006999528)

Registrant - Time Cap Laboratories, Inc. (037052099)

Establishment								
Name	Address	ID/FEI	<b>Business Operations</b>					
MARKSANS PHARMA LIMITED		925822975	manufacture(41226-697)					

Revised: 4/2024

Kroger Company