

**NIGHTTIME COLD AND FLU- dextromethorphan hydrobromide,
acetaminophen, doxylamine succinate capsule, liquid filled
Kroger Company**

**694T KROGER 41226 694 Acetaminophen Dextromethorphan Hydrobromide
Doxylamine Succinate CT 24**

DRUG FACTS

Acetaminophen 325 mg
Dextromethorphan Hydrobromide 15 mg
Doxylamine succinate 6.25 mg

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

- temporarily relieves common cold/flu symptoms:
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 softgels in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy Alert:

acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over: 2 softgels with water every 6 hrs
children 4 to under 12 yrs: ask a doctor

children under 4 yrs: do not use

OTHER INFORMATION

Other information

- store at room temperature

Inactive ingredients FD&C Blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide.

Questions ? 1-800-632-6900

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Our Brands. Our Guarantee.
Love It or Your Money Back.
www.kroger.com

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 15 mg Cough suppressant
Doxylamine succinate 6.25 mg Antihistamine

Purpose

Uses ■ temporarily relieves common cold/flu symptoms:
■ cough due to minor throat & bronchial irritation
■ sore throat ■ headache ■ minor aches & pains ■ fever ■ runny nose & sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 8 softgels in 24 hrs, which is the maximum daily amount for this product
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away

Some throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription nonanesthetic opioid analgesic (MOA) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MOA drug. If you do not know if your prescription drug contains an MOA, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Other information ■ store at room temperature

Inactive ingredients FD&C blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide.

Drug Facts (continued)

Directions

■ take only as directed	■ do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	2 softgels with water every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use



Cold & Flu

Nighttime

Multi-Symptom Relief

Acetaminophen
Dextromethorphan HBr
Doxylamine Succinate



actual size

COMPARE TO the active ingredients of VICKS™ NYQUIL™
COLD & FLU RELIEF LIQUICAPS™
**See side panel

NDC 41226-694-24

POWERFUL NIGHTTIME RELIEF

Pain Reliever
Fever Reducer
Cough Suppressant
Antihistamine
24 SOFTGELS

Questions? 1-800-632-6900
DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202
FOR MORE PRODUCT INFORMATION, SCAN UPC USING YOUR KROGER APP OR CALL 800-632-6900
**Vicks, NYQUIL and LiquiCaps are trademarks of The Procter & Gamble Company, Cincinnati, OH 45202. The Procter & Gamble Company is not affiliated with The Kroger Co. or this product.



PLEASE RECYCLE

100% SIZE
12-digit UPC-A
(non-suppressed)
For Position Only
0 41260 01241 9

694T 0321

SAFETY SEALED: DO NOT USE IF THE INDIVIDUAL BUSTER UNIT IS OPEN OR TORN

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Lot #:

Exp. Date:

NIGHTTIME COLD AND FLU

dextromethorphan hydrobromide, acetaminophen, doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-694
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GLYCERIN (UNII: PDC6A3C0OX)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
GELATIN (UNII: 2G86QN327L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL (Oblong shaped)	Size	21mm
Flavor		Imprint Code	71
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-694-24	2 in 1 CARTON	05/11/2021	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/11/2021	

Labeler - Kroger Company (006999528)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(41226-694)

Revised: 2/2025

Kroger Company