

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**MEIJER DISTRIBUTION INC.**

-----  
***Drug Facts***

***Active ingredient (in each capsule)***

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/Fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### ***Other information***

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### ***Inactive ingredients***

gelatin, glycerin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

For India Manufacturing Site:

FD&C Blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

### ***Questions or comments?***

**1-888-235-2466**

†This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademarks Motrin® .

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH “ SEALED for YOUR PROTECTION” IS BROKEN OR MISSING.**

DIST.

MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

PID 5506556

R0125

L0000944

**For India Manufacturing Site:**

**MADE IN INDIA**

Code: TN/DRUGS/TN/25/00074

L0000992

R0225

P3294/00/25

**20's carton**

**NDC 41250-286-11**

**†COMPARE TO THE ACTIVE**

**INGREDIENT IN MOTRIN®**

**IB LIQUID GELS**

**meijer**

**Dye-free**

**Ibuprofen**

**mini**

**capsules**

**capsules| 200 mg**

**PAIN RELIEVER/FEVER REDUCER (NSAID)**

**SMALLER CAPSULE**

**SAME STRENGTH**

**20 Capsules\*\***

**\*\*liquid-filled capsules**

**Drug Facts (continued)**

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing  
■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used  
■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

**meijer.**  
Dye-Free  
**Ibuprofen**

NDC 41250-286-11

COMPARE TO THE ACTIVE INGREDIENT IN MOTRIN® IB LIQUID GELS

MINI CAPSULES Capsules | 200 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)  
SMALLER CAPSULE  
SAME STRENGTH

20 Capsules\*\*

\*\*liquid-filled capsules



This product is not manufactured or distributed by Meijer Inc., owner of the registered trademark Meijer.

DIST. MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

PID 5506556

R0125 L0000944



Lot No.:  
Exp. Date:

**Drug Facts (continued)**  
**Other information** ■ each capsule contains:  
potassium 20 mg ■ read all warnings and directions before use. Keep carton. ■ store at 20° to 25°C (68° to 77°F) ■ avoid excessive heat above 40°C (104°F). Protect from light.  
**Inactive ingredients**  
gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol.  
**Questions or comments?**  
call toll-free 1-888-232-2466  
DO NOT USE IF TAMPER-EVIDENT SEAL UNDER CAP IS CRACKED OR MISSING. WHEN SEALED BY YOUR PHARMACEUTICAL MANUFACTURER, THIS PRODUCT IS PROTECTED AGAINST UNLAWFUL REPACKAGING AND COUNTERFEITING. READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

**Drug Facts**

**Active ingredient (in each capsule)** **Purpose**  
Sorbilized ibuprofen equal to 200 mg ibuprofen (NSAID)\* ..... Pain reliever/Fever reducer  
(present as the free acid and potassium salt) ..... \*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache  
■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis  
■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

**Drug Facts (continued)**

■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

Rev # 02/22  
T00087

120's label-Made in India

NDC 41250-286-22

meijer

dye-free

ibuprofen

Compare to the active ingredient

in Motrin® IB LIQUID GELS

capsules 200 mg

PAIN RELIEVER/FEVER REDUCER

(NSAID)

120 Softgels\*\*

\*\*liquid-filled capsules

**meijer.** NDC 41250-286-22

**Dye-Free ibuprofen**

Compare to the active ingredient in Motrin® IB LIQUID GELS†

**Capsules | 200 mg**

**PAIN RELIEVER/FEVER REDUCER (NSAID)**

120 Softgels\*\*  
\*\*Liquid-Filled Capsules Actual Size

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

**Drug Facts**

**Active ingredient (in each capsule)** Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*. **Purpose** Pain reliever/fever reducer (present as the free acid and potassium salt) \*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn

Manufactured for: **Bionpharma Inc.**, Princeton, NJ 08540  
**MADE IN INDIA** 7 60236 50427 6  
Code: TN / DRUGS / TN/25/00074 Lot No.: P-254/0025 Exp. Date: R0225 L0000992

**PEEL HERE**

**Drug Facts (continued)**

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools  
■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

**Other information** ■ each capsule contains: potassium 20 mg ■ read all warnings and directions before use ■ store at 20° to 25°C (68° to 77°F) ■ avoid excessive heat above 40°C (104°F). Protect from light.

**Inactive ingredients** gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

**Questions or comments?**  
call toll-free 1-888-235-2466

This product is not manufactured or distributed by the owners of Meijer® IB Liquid Gels.

T00074R0325

## IBUPROFEN

ibuprofen capsule, liquid filled

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:41250-286

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

### Product Characteristics

Color	blue	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	IB4
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-286-22	120 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2021	
2	NDC:41250-286-11	1 in 1 CARTON	02/24/2021	
2		20 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	02/24/2021	

Labeler - MEIJER DISTRIBUTION INC. (006959555)

Registrant - Bionpharma Inc. (079637826)

Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Patheon Softgels Inc.		002193829	manufacture(41250-286)

## **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
ESJAY PHARMA PRIVATE LIMITED		957309497	manufacture(41250-286)

Revised: 10/2025

MEIJER DISTRIBUTION INC.