

IBUPROFEN- ibuprofen capsule, liquid filled
Kroger Company

Kroger Ibuprofen Capsules 200mg

DRUG FACTS

Active ingredient(in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)'
(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:

headache

toothache

backache

menstrual cramps

the common cold

muscular aches

minor pain of arthritis

- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin,

ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or alter heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

- you have symptoms of heart problems or stroke:

chest pain

trouble breathing

weakness in one part or side of body

slurred speech

leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

OTHER INFORMATION

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F).

Questions or comments? 1-800-632-6900

Inactive ingredients

ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monooleate, sorbitol sorbitan

NDC 41226-610-80



COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®

Ibuprofen

Ibuprofen Capsules 200 mg

PAIN RELIEVER/
FEVER REDUCER
[NSAID]*

80 SOFTGELS



ACTUAL SIZE

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each capsule)
Solubilized ibuprofen (NSAID)*

200 mg ibuprofen (NSAID)*.....Pain reliever/
fever reducer
(present as the free acid and potassium salt) Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued under label)

Ibuprofen is a registered trademark of Heileman US Holdings, LLC. Heileman US Holdings, LLC is not affiliated with The Kroger Co. or this product. Liqui-Gels is a registered trademark of Cabot Pharmaceuticals, Inc. Cabot Pharmaceuticals, Inc. is not affiliated with The Kroger Co. or this product.

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA
For More Product Information,
Scan UPC Using Your Kroger App
or Call 800-632-6900

OUR PHARMACIST
RECOMMENDED



4 1260 02173 2

LOT: 6107 0324

EXP:

Vainish Ornit Area

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - teal faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

- keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monododecylate, sorbitol sorbitan

Questions? Call 1-800-632-6900

Adhesive Area

NDC 41226-610-20



COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®

Ibuprofen

Ibuprofen Capsules 200 mg

PAIN RELIEVER/
FEVER REDUCER
(NSAID)*

200
SOFTGELS



ACTUAL SIZE

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen (NSAID) 200 mg (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/
Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - tubercle
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- swelling
- asthma (wheezing)
- stomach
- skin reddening
- rash
- blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued under label)

*Advil is a registered trademark of Reliance US Holdings LLC. Reliance US Holdings LLC is not affiliated with The Kroger Co. or this product. Liqui-Gels is a registered trademark of Cabot Pharmaceuticals, Inc. Cabot Pharmaceuticals, Inc. is not affiliated with The Kroger Co. or this product.

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA

For More Product Information:
Scan UPC Using Your Kroger App
or Call 800-452-6900

YOUR PHARMACIST RECOMMENDED 0 41260-02172 5

LOT:

EXP.:

Varnish Omit Area



Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shellac, sorbitan monoleate, sorbitol sorbitan

Questions or comments? Call 1-800-632-6900

Adhesive Area

COMPARE TO the active ingredient of
ADVIL® LIQUI-GELS® †See inside panel



NDC 41226-610-66

EASY OPEN CAP

Ibuprofen
Capsules, 200 mg
Pain Reliever/Fever
Reducer (NSAID)



actual
size



THIS PACKAGE FOR
HOUSEHOLDS WITHOUT
YOUNG CHILDREN

160 SOFTGELS[†]
(†liquid filled capsules)

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL
SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

Drug Facts

Active ingredient (in each capsule) Purpose
Sulbutilized ibuprofen equal to.....Pain reliever/fever reducer
200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:
● headache ● toothache ● backache ● menstrual cramps
● the common cold ● muscular aches ● minor pain of arthritis
● temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
● hives ● facial swelling ● asthma (wheezing)
● shock ● skin reddening ● rash ● blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
● are age 60 or older
● have had stomach ulcers or bleeding problems
● take a blood thinning (anticoagulant) or steroid drug
● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
● have 3 or more alcoholic drinks every day while using this product
● take more or for a longer time than directed

Drug Facts (continued under label)

DIST. BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA

FOR MORE PRODUCT
INFORMATION, SCAN UPC
USING YOUR KROGER APP
OR CALL 800-632-6900
6111 44Z



Lot No.:

Exp. Date:

PEEL
HERE

Non Varnish Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. Use right before or after heart surgery.

Ask a doctor before use if ● stomach bleeding warning applies to you ● you have problems or serious side effects from taking pain relievers or fever reducers ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ● you are taking a diuretic

Ask a doctor or pharmacist before use if you are ● under a doctor's care for any serious condition ● taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ● taking any other drug

When using this product ● take with food or milk if stomach upset occurs

Stop use and ask a doctor if

● you experience any of the following signs of stomach bleeding:

- feel faint ● vomit blood ● have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling
- pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● redness or swelling is present in the painful area ● any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ● do not take more than directed ● the smallest effective dose should be used ● adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist ● if pain or fever does not respond to 1 capsule, 2 capsules may be used ● do not exceed 6 capsules in 24 hours, unless directed by a doctor ● children under 12 years: ask a doctor

Other information ● each capsule contains:

potassium 20 mg ● read all warnings and directions before use ● store at 20-25°C (68-77°F) ● avoid excessive heat above 40°C (104°F).

Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, shallic, sorbitan monooleate, sorbitol sorbitan

Questions or comments? Call 1-800-632-6900

†Advil is a registered trademark of GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, Wilmington, DE 19808. GlaxoSmithKline Consumer Healthcare Holdings (US) LLC is not affiliated with The Kroger Co. or this product. Liqui-Gels is a registered trademark of Catalent Pharma Solutions, Inc., Somerset, NJ 08873. Catalent Pharma Solutions, Inc., is not affiliated with The Kroger Co. or this product.

NDC 41226-610-23
Kroger health



Ibuprofen

Ibuprofen Capsules 200 mg

PAIN RELIEVER/
FEVER REDUCER
(NSAID)*

20 SOFTGELS

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION

Uses

- temporarily relieves minor aches and pains, due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs,

Drug Facts (continued under label)

LOT:
PER
HERE
EXP:

Varnish Omit Area

Drug Facts (continued)

except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

do not take more than directed
the smallest effective dose should be used

- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

each capsule contains: potassium 20 mg

Questions or comments?

Call 1-800-632-6900

DISTRIBUTED BY THE KROGER CO.
CINCINNATI, OHIO 45202

MADE IN INDIA

For More Product Information,
Scan UPC Using Your Kroger App or
Call 800-632-6900

YOUR PHARMACIST RECOMMENDED

610T 0324

Adhesive Area

C6011R2
Varnish/Print Omit Area

NDC 41226-610-23

Kroger⁺ health



COMPARE TO THE ACTIVE
INGREDIENT IN ADVIL® LIQUID-GELS®

Ibuprofen

Ibuprofen Capsules 200 mg

PAIN RELIEVER/FEVER
REDUCER (NSAID)*

20 SOFTGELS



Varnish Omit

Varnish Omit

Drug Facts

SEAL UNDER BOTTLE CAP
WARNING: READ PRODUCT INFORMATION

Active ingredient (in each capsule) Purpose
Sulfonized ibuprofen (NSAID)*
200 mg Ibuprofen (NSAID)*
Pain reliever/fever reducer (NSAID)*

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- snook
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Advil is a registered trademark of Heinen US Holdings, LLC. Heinen US Holdings, LLC is not affiliated with the Kroger Co. or this product. Liquid is a registered trademark of Cabot Pharmaceuticals, Inc. Cabot Pharmaceuticals, Inc. is not affiliated with the Kroger Co. or this product.

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OH 45202
MADE IN INDIA

OUR PHARMACIST RECOMMENDED
Love It or Your Money Back.
www.kroger.com

Drug Facts (continued)

Take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer
 right before or after heart surgery

Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers
 you have a history of stomach problems, such as heartburn
 you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 under a doctor's care for any serious condition
 taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 taking any other drug

When using this product:
 Take with food or milk if stomach upset occurs

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:
 red, black, or tarry stools
 vomit blood
 have bloody or black stools
 have stomach pain that does not

Drug Facts (continued)

get better
 chest pain
 trouble breathing
 weakness in one part or side of body
 slurred speech
 leg swelling
 pain gets worse or lasts more than 10 days
 fever gets worse or lasts more than 3 days
 redness or swelling is present in the painful area
 any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions Do not take more than directed
 the smallest effective dose should be used
 adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
 if pain or fever does not respond to 1 capsule, 2 capsules may be used
 do not exceed 6 capsules in 24 hours, unless directed by a doctor
 children under 12 years: ask a doctor

Other information each capsule contains: potassium 20 mg
 read all warnings and directions before use
 store at 20-25°C (68-77°F)
 avoid excessive heat above 40°C (104°F)

Drug Facts (continued)
Inactive ingredients ammonium hydroxide, FD&C green #3, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, potassium hydroxide, propylene glycol, purified water, stibelic, sorbitan monooleate, sorbitol sorbitan

Questions or comments?
 Call 1-800-632-6900

LOT:
EXP:
Varnish Omit

0 41260 02174 9

IBUPROFEN
 ibuprofen capsules, liquid filled

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:41226-610 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| GELATIN (UNII: 2G86QN327L) | |
| MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) | |
| WATER (UNII: 059QF0KO0R) | |
| SHELLAC (UNII: 46N107B71O) | |
| SORBITAN MONOOLEATE (UNII: 06XEA2VD56) | |
| AMMONIA (UNII: 5138Q19F1X) | |
| FERROUS OXIDE (UNII: G7036X8B5H) | |
| FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SORBITOL (UNII: 506T60A25R) | |

Product Characteristics

| | | | |
|-----------------|---------|---------------------|----------|
| Color | green | Score | no score |
| Shape | CAPSULE | Size | 19mm |
| Flavor | | Imprint Code | 133 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:41226-610-66 | 160 in 1 BOTTLE; Type 0: Not a Combination Product | 06/12/2021 | |
| 2 | NDC:41226-610-20 | 200 in 1 BOTTLE; Type 0: Not a Combination Product | 04/15/2024 | |
| 3 | NDC:41226-610-23 | 1 in 1 CARTON | 04/15/2024 | |
| 3 | | 20 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 4 | NDC:41226-610-80 | 80 in 1 BOTTLE; Type 0: Not a Combination Product | 04/15/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| | | | |

| | | | |
|------|------------|------------|--|
| ANDA | ANDA079205 | 06/12/2021 | |
|------|------------|------------|--|

Labeler - Kroger Company (006999528)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| MARKSANS PHARMA LIMITED | | 925822975 | manufacture(41226-610) |

Revised: 4/2024

Kroger Company