

MCKESSON ACETAMINOPHEN 325- acetaminophen tablet, coated
McKesson Medical-Surgical, Inc.

McKesson Acetaminophen 325 mg

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves of minor aches and pains due to:

- headache
- muscular aches
- minor arthritis pain
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a

drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse and lasts for more than 10 days

- fever gets worse or lasts for more than 3 days

- new symptoms occur

- redness or swelling is present

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**

Adults and children: (12 years and over)

- take 2 tablets every 4 to 6 hours while symptoms last.

- do not take more than 10 tablets in 24 hours, unless directed by a doctor

- do not use for more than 10 days unless directed by a doctor

Children under 12 years:

- Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)

- tamper-evident sealed packets

- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-800-777-4908

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	AZ;234
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68599-4679-9	250 in 1 BOX	04/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/01/2021	

Labeler - McKesson Medical-Surgical, Inc. (023904428)

Registrant - Unifirst Fiirst Aid Corporation (832947092)

Revised: 9/2025

McKesson Medical-Surgical, Inc.