


**HYDROXYZINE PAMOATE- hydroxyzine pamoate capsule**  
**Amneal Pharmaceuticals of New York LLC**

-----  
**Hydroxyzine Pamoate Capsules, USP**  
**(25 mg and 50 mg)**

**Rx only**

**DESCRIPTION**

Hydroxyzine pamoate, USP is a light yellow, practically odorless powder, practically insoluble in water and methanol and freely soluble in dimethylformamide. It is chemically designated as 1-(p-chlorobenzhydryl) 4-[2-(2-hydroxyethoxy) ethyl] diethylenediamine salt of 1,1'-methylene bis (2 hydroxy-3-naphthalene carboxylic acid) and can be structurally represented as follows:



Chemical Formula: C<sub>21</sub>H<sub>27</sub>ClN<sub>2</sub>O<sub>2</sub>•C<sub>23</sub>H<sub>16</sub>O<sub>6</sub>  
Molecular Weight: 763.29

Each capsule, for oral administration, contains hydroxyzine pamoate, USP equivalent to 25 mg or 50 mg of hydroxyzine hydrochloride. In addition, each capsule contains the following inactive ingredients: colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, gelatin, magnesium stearate, pregelatinized starch, sodium lauryl sulfate, and titanium dioxide. The imprinting ink on the capsules contains synthetic black iron oxide.

**CLINICAL PHARMACOLOGY**

Hydroxyzine pamoate is unrelated chemically to the phenothiazines, reserpine, meprobamate, or the benzodiazepines. Hydroxyzine pamoate is not a cortical depressant, but its action may be due to a suppression of activity in certain key regions of the subcortical area of the central nervous system. Primary skeletal muscle relaxation has been demonstrated experimentally. Bronchodilator activity and antihistaminic and analgesic effects have been demonstrated experimentally and confirmed clinically. An antiemetic effect, both by the apomorphine test and the veriloid test, has been demonstrated. Pharmacological and clinical studies indicate that hydroxyzine in therapeutic dosage does not increase gastric secretion or acidity and in most cases has mild antisecretory activity. Hydroxyzine is rapidly absorbed from the gastrointestinal tract and hydroxyzine pamoate's clinical effects are usually noted within 15 to 30 minutes after oral administration.

**INDICATIONS**

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested.

Useful in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses, and in histamine-mediated pruritus.

As a sedative when used as premedication and following general anesthesia, **Hydroxyzine may potentiate meperidine (Demerol®) and barbiturates**, so their use in pre-anesthetic adjunctive therapy should be modified on an individual basis. Atropine and other belladonna alkaloids are not affected by the drug. Hydroxyzine is not known to interfere with the action of digitalis in any way and it may be used concurrently with this agent.

The effectiveness of hydroxyzine as an antianxiety agent for long-term use, that is, more than 4 months, has not been assessed by systematic clinical studies. The physician should reassess periodically the usefulness of the drug for the individual patient.

## CONTRAINDICATIONS

Hydroxyzine, when administered to the pregnant mouse, rat, and rabbit, induced fetal abnormalities in the rat and mouse at doses substantially above the human therapeutic range. Clinical data in human beings are inadequate to establish safety in early pregnancy. Until such data are available, hydroxyzine is contraindicated in early pregnancy.

Hydroxyzine is contraindicated in patients with a prolonged QT interval.

Hydroxyzine pamoate is contraindicated for patients who have shown a previous hypersensitivity to any component of this medication.

Hydroxyzine is contraindicated in patients with known hypersensitivity to hydroxyzine products, and in patients with known hypersensitivity to cetirizine hydrochloride or levocetirizine hydrochloride.

## WARNINGS

### Nursing Mothers:

It is not known whether this drug is excreted in human milk. Since many drugs are so excreted, hydroxyzine should not be given to nursing mothers.

## PRECAUTIONS

THE POTENTIATING ACTION OF HYDROXYZINE MUST BE CONSIDERED WHEN THE DRUG IS USED IN CONJUNCTION WITH CENTRAL NERVOUS SYSTEM DEPRESSANTS SUCH AS NARCOTICS, NON-NARCOTIC ANALGESICS AND BARBITURATES. Therefore, when central nervous system depressants are administered concomitantly with hydroxyzine, their dosage should be reduced. Since drowsiness may occur with use of the drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery while taking hydroxyzine pamoate. Patients should be advised against the simultaneous use of other CNS depressant drugs, and cautioned that the effect of alcohol may be increased.

**QT Prolongation/Torsade de Pointes (TdP):** Cases of QT prolongation and Torsade de Pointes have been reported during post-marketing use of hydroxyzine. The majority of reports occurred in patients with other risk factors for QT prolongation/TdP (pre-existing heart disease, electrolyte imbalances or concomitant arrhythmogenic drug use). Therefore, hydroxyzine should be used with caution in patients with risk factors for QT prolongation, congenital long QT syndrome, a family history of long QT syndrome, other conditions that predispose to QT prolongation and ventricular arrhythmia, as well as recent myocardial infarction, uncompensated heart failure, and bradyarrhythmias.

Caution is recommended during the concomitant use of drugs known to prolong the QT interval. These include Class 1A (e.g., quinidine, procainamide) or Class III (e.g., amiodarone, sotalol) antiarrhythmics, certain antipsychotics (e.g., ziprasidone, iloperidone, clozapine, quetiapine, chlorpromazine), certain antidepressants (e.g., citalopram, fluoxetine), certain antibiotics (e.g., azithromycin, erythromycin, clarithromycin, gatifloxacin, moxifloxacin); and others (e.g., pentamidine, methadone, ondansetron, droperidol).

**Acute Generalized Exanthematous Pustulosis (AGEP):** Hydroxyzine may rarely cause acute generalized exanthematous pustulosis (AGEP), a serious skin reaction characterized by fever and numerous small, superficial, non-follicular, sterile pustules, arising within large areas of edematous erythema. Inform patients about the signs of AGEP, and discontinue hydroxyzine at the first appearance of a skin rash, worsening of pre-existing skin reactions which hydroxyzine may be used to treat, or any other sign of hypersensitivity. If signs or symptoms suggest AGEP, use of hydroxyzine should not be resumed and alternative therapy should be considered. Avoid cetirizine or levocetirizine in patients who have experienced AGEP or other hypersensitivity reactions with hydroxyzine, due to the risk of cross-sensitivity.

**Geriatric Use:** A determination has not been made whether controlled clinical studies of hydroxyzine pamoate included sufficient numbers of subjects aged 65 and over to define a difference in response from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function and of concomitant disease or other drug therapy.

The extent of renal excretion of hydroxyzine pamoate has not been determined. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selections. Sedating drugs may cause confusion and over sedation in the elderly; elderly patients generally should be started on low doses of hydroxyzine pamoate and observed closely.

## ADVERSE REACTIONS

Side effects reported with the administration of hydroxyzine pamoate are usually mild and transitory in nature.

**Skin and Appendages:** Oral hydroxyzine hydrochloride is associated with Acute Generalized Exanthematous Pustulosis (AGEP) and fixed drug eruptions in post-marketing reports.

**Anticholinergic:** Dry mouth.

**Central Nervous System:** Drowsiness is usually transitory and may disappear in a few days of continued therapy or upon reduction of the dose. Involuntary motor activity, including rare instances of tremor and convulsions, has been reported, usually with doses considerably higher than those recommended. Clinically significant respiratory depression has not been reported at recommended doses.

**Cardiac System:** QT prolongation, Torsade de Pointes.

In post-marketing experience, the following additional undesirable effects have been reported:

**Body as a Whole:** allergic reaction,

**Nervous System:** headache,

**Psychiatric:** hallucination,

**Skin and Appendages:** pruritus, rash, urticaria.

**To report SUSPECTED ADVERSE REACTIONS, contact Amneal Pharmaceuticals at 1-877-835-5472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## OVERDOSAGE

The most common manifestation of overdosage of hydroxyzine pamoate is hypersedation. Other reported signs and symptoms were convulsions, stupor, nausea and vomiting. As in the management of overdosage with any drug, it should be borne in mind that multiple agents may have been taken.

If vomiting has not occurred spontaneously, it should be induced. Immediate gastric lavage is also recommended. General supportive care, including frequent monitoring of the vital signs and close observation of the patient, is indicated. Hypotension, though unlikely, may be controlled with intravenous fluids and vasopressors (**do not use epinephrine as hydroxyzine counteracts its pressor action**). Caffeine and Sodium Benzoate Injection, USP, may be used to counteract central nervous system depressant effects.

Hydroxyzine overdose may cause QT prolongation and Torsade de Pointes. ECG monitoring is recommended in cases of hydroxyzine overdose.

There is no specific antidote. It is doubtful that hemodialysis would be of any value in the treatment of overdosage with hydroxyzine. However, if other agents such as barbiturates have been ingested concomitantly, hemodialysis may be indicated. There is no practical method to quantitate hydroxyzine in body fluids or tissue after its ingestion or administration.

## DOSAGE

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested: in adults, 50 mg to 100 mg q.i.d.; children under 6 years, 50 mg daily in divided doses; and over 6 years, 50 mg to 100 mg daily in divided doses.

For use in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses, and in histamine-mediated pruritus: in adults, 25 mg t.i.d. or q.i.d.; children under 6 years, 50 mg daily in divided doses; and over 6 years, 50 mg to 100 mg daily in divided doses.

As a sedative when used as a premedication and following general anesthesia: 50 mg to 100 mg in adults, and 0.6 mg/kg in children. When treatment is initiated by the intramuscular route of administration, subsequent doses may be administered orally.

As with all medications, the dosage should be adjusted according to the patient's response to therapy.

## HOW SUPPLIED

Hydroxyzine pamoate capsules, USP (hydroxyzine pamoate equivalent to hydroxyzine hydrochloride) are supplied as follows:

**25 mg capsules:** Dark green opaque cap/light green opaque body filled with yellow powder and imprinted in black "IX" on the capsule cap and "657" on the capsule body.

They are available as follows:

Bottles of 100:	NDC 0115-1803-01
Bottles of 500:	NDC 0115-1803-02

**50 mg capsules:** Dark green opaque cap/white opaque body filled with yellow powder and imprinted in black "IX" on the capsule cap and "658" on the capsule body.

They are available as follows:

Bottles of 100:	NDC 0115-1804-01
Bottles of 500:	NDC 0115-1804-02

Store below 30°C (86°F) [see USP].

Dispense in a tight, light resistant container as defined in USP/NF.

## BIBLIOGRAPHY

Available on request.

Brands listed are the trademarks of their respective owners.

Manufactured by:

**Patheon Pharmaceuticals Inc.**

Cincinnati, OH 45237 USA

OR

Manufactured by:

**Amneal Pharmaceuticals Pvt. Ltd.**

Ahmedabad 382220, INDIA

Distributed by:

**Amneal Pharmaceuticals LLC**

Bridgewater, NJ 08807

Revised: 12-2019-00

## PRINCIPAL DISPLAY PANEL - 25 mg

NDC 0115-1803-01

**HydroXYzine**  
**Pamoate**  
**Capsules, USP**

**25 mg\***  
Tartrazine Dye-Free

  
**Rx only**  
100 Capsules



\*EACH CAPSULE CONTAINS:  
HydroXYzine Pamoate, USP equivalent to 25 mg  
Hydroxyzine Hydrochloride

**USUAL DOSAGE:** ADULTS: 25 mg tid to 100 mg qid.  
CHILDREN: Under 6 years - 50 mg daily in divided doses.  
Over 6 years - 50 mg to 100 mg daily in divided doses.

Dispense in a tight, light-resistant container as defined in USP/NF.

**Store below 30°C (86°F). [see USP].**

**Keep out of reach of children.**

Manufactured by: **Patheon Pharmaceuticals Inc.**  
Cincinnati, OH 45237 USA

Manufactured for: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Rev. 12-2018-00

70037754

  
01151803015

NDC 0115-1803-01

# hydrOXYzine Pamoate Capsules, USP

25 mg\*

Tartrazine Dye-Free



Rx only  
100 Capsules



\*EACH CAPSULE CONTAINS:  
HydrOXYzine Pamoate, USP equivalent to 25 mg  
Hydroxyzine Hydrochloride

**USUAL DOSAGE:** ADULTS: 25 mg t.i.d. to 100 mg q.i.d.

CHILDREN: Under 6 years - 50 mg daily in divided doses.  
Over 6 years - 50 mg to 100 mg daily in divided doses.  
See accompanying prescribing information.

Dispense in a tight, light-resistant container as defined in USP.

**Store below 30°C (86°F) [see USP].**

**Keep out of reach of children.**

Manufactured by: **Amneal Pharmaceuticals Pvt. Ltd.**  
Ahmedabad, 382220 INDIA  
Distributed by: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Mfg. Lic. No. G/25/1941

Rev. 12-2019-00



Non-Varnish Area  
(For Lot And Exp. Date)  
(22 X 45 mm)

## PRINCIPAL DISPLAY PANEL - 50 mg

NDC 0115-1804-01

# HydrOXYzine Pamoate Capsules, USP

50 mg\*

Tartrazine Dye-Free



Rx only  
100 Capsules



\*EACH CAPSULE CONTAINS:  
HydrOXYzine Pamoate, USP equivalent to 50 mg  
Hydroxyzine Hydrochloride

**USUAL DOSAGE:** ADULTS: 25 mg tid to 100 mg qid.

CHILDREN: Under 6 years - 50 mg daily in divided doses.  
Over 6 years - 50 mg to 100 mg daily in divided doses.

Dispense in a tight, light-resistant container as defined in USP/NF.

**Store below 30°C (86°F). [see USP].**

**Keep out of reach of children.**

Manufactured by: **Patheon Pharmaceuticals Inc.**  
Cincinnati, OH 45237 USA

Manufactured for: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Rev. 12-2018-00

70037758



Non-Varnish Area  
(For Lot And Exp. Date)  
(22 X 51 mm)

NDC 0115-1804-01

# hydrOXYzine Pamoate Capsules, USP

50 mg\*

Tartrazine Dye-Free



Rx only  
100 Capsules



\*EACH CAPSULE CONTAINS:  
HydrOXYzine Pamoate, USP equivalent to 50 mg  
Hydroxyzine Hydrochloride

**USUAL DOSAGE:** ADULTS: 25 mg t.i.d. to 100 mg q.i.d.

CHILDREN: Under 6 years - 50 mg daily in divided doses.  
Over 6 years - 50 mg to 100 mg daily in divided doses.  
See accompanying prescribing information.

Dispense in a tight, light-resistant container as defined in USP.

**Store below 30°C (86°F) [see USP].**

**Keep out of reach of children.**

Manufactured by: **Amneal Pharmaceuticals Pvt. Ltd.**  
Ahmedabad, 382220 INDIA  
Distributed by: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

Mfg. Lic. No. G/25/1941

Rev. 12-2019-00



Non-Varnish Area  
(For Lot And Exp. Date)  
(22 X 51 mm)

## HYDROXYZINE PAMOATE

hydroxyzine pamoate capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0115-1803	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
HYDROXYZINE PAMOATE (UNII: M20215MUFR) (HYDROXYZINE - UNII:30S50YM8OG)		HYDROXYZINE HYDROCHLORIDE	25 mg	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)				
GELATIN (UNII: 2G86QN327L)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
FERROSFERRIC OXIDE (UNII: XM0M87F357)				
STARCH, CORN (UNII: O8232NY3SJ)				
<b>Product Characteristics</b>				
<b>Color</b>	GREEN (dark green opaque/light green opaque)	<b>Score</b>	no score	
<b>Shape</b>	CAPSULE	<b>Size</b>	16mm	
<b>Flavor</b>		<b>Imprint Code</b>	IX;657	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0115-1803-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/15/1996	
2	NDC:0115-1803-02	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/15/1996	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA040156	07/15/1996		

## HYDROXYZINE PAMOATE

hydroxyzine pamoate capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0115-1804	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
HYDROXYZINE PAMOATE (UNII: M20215MUFR) (HYDROXYZINE - UNII:30S50YM8OG)		HYDROXYZINE HYDROCHLORIDE	50 mg	
<b>Inactive Ingredients</b>				
<b>Ingredient Name</b>			<b>Strength</b>	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)				
GELATIN (UNII: 2G86QN327L)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
FERROSO FERRIC OXIDE (UNII: XM0M87F357)				
<b>Product Characteristics</b>				
<b>Color</b>	GREEN (opaque dark green) , WHITE (opaque white)		<b>Score</b>	no score
<b>Shape</b>	CAPSULE		<b>Size</b>	19mm
<b>Flavor</b>			<b>Imprint Code</b>	IX;658
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0115-1804-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/15/1996	
2	NDC:0115-1804-02	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/15/1996	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA040156	06/15/1996		

**Labeler** - Amneal Pharmaceuticals of New York LLC (123797875)

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
-------------	----------------	---------------	----------------------------



Amneal Pharmaceuticals Private Limited Oral Solid Dosage Unit		650762060	ANALYSIS(0115-1803, 0115-1804) , LABEL(0115-1803, 0115-1804) , MANUFACTURE(0115-1803, 0115-1804) , PACK(0115-1803, 0115-1804)
---	--	-----------	--

## Establishment

Name	Address	ID/FEI	Business Operations
Patheon Pharmaceuticals Inc.		005286822	ANALYSIS(0115-1803, 0115-1804) , LABEL(0115-1803, 0115-1804) , MANUFACTURE(0115-1803, 0115-1804) , PACK(0115-1803, 0115-1804)

Revised: 12/2019

Amneal Pharmaceuticals of New York LLC