

IBUPROFEN- ibuprofen tablet, film coated
Walgreen Company

Walgreens 44-392

Active ingredient (in each orange tablet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - toothache
 - menstrual cramps
 - headache
 - the common cold
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
- facial swelling
- blisters
- rash
- hives
- skin reddening
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or Comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Walgreens

Compare to Motrin® IB Tablets active ingredient^{††}

NDC 0363-0392-12

Ibuprofen 200

Ibuprofen USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

TABLETS

100 TABLETS

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

^{††}This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Tablets.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2018 Walgreen Co.

50844 ORG061839212

ORG0618-F

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
 ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor

Other information

■ store between 20°-25° C (68°-77°F)
 ■ avoid excessive heat 40° C (104°F)
 ■ see end flap for expiration date and lot number

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

(Questions or comments?)

Monday-Friday
 Call 1-800-426-9991 8:30 AM-4:00 PM ET,
 Walgreens.com

No Print Area
 Lot no. & Exp. Date



ITEM 500729 W00000-0000-0
 SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts (continued)

and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if

■ you have a history of stomach problems, such as heartburn, ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ you have symptoms of heart problems or stroke:
 ■ chest pain ■ slurred speech
 ■ irregular heartbeat ■ trouble breathing
 ■ weakness in one part or side of body
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

50844 ORG061839212
 Walgreens.com ©2018 Walgreen Co.
 200 WILLMOT RD, DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 DISTRIBUTED BY: WALGREEN CO.
 Trademark Motrin® IB Tablets.
 Corporation, owner of the registered
 distributed by Johnson & Johnson
 Walgreens Pharmacist Recommended
 ! This product is not manufactured or

B-2201-392-12-H
 ORG061839212

Drug Facts

Active ingredient
 (in each orange tablet)
 Ibuprofen USP, 200 mg (NSAID)*
 *nonsteroidal anti-inflammatory drug

Purpose
 Pain reliever/fever reducer

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ toothache ■ headache ■ the common cold
 ■ minor pain of arthritis
 ■ temporarily reduces fever
 ■ temporarily relieves menstrual cramps

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:
 ■ rash ■ facial swelling ■ hives
 ■ shock ■ skin redness ■ asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

■ take more or for a longer time than directed
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure,

No Print
 Glue Area

Walgreens

Compare to Motrin® IB
 Tablets active ingredient!

Ibuprofen 200

NDC 0363-0392-12

IBUPROFEN USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

TABLETS

100 TABLETS



No Print
 Glue Area

Walgreens 44-392

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:0363-0392

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;392
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0392-15	1 in 1 CARTON	03/01/1999	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0392-12	1 in 1 CARTON	03/01/1999	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-0392-37	1 in 1 CARTON	03/01/1999	02/07/2021
3		75 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0392-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/1999	
5	NDC:0363-0392-16	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/1999	
6	NDC:0363-0392-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/1999	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

Labeler - Walgreen Company (008965063)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(0363-0392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(0363-0392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(0363-0392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(0363-0392)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(0363-0392)

Revised: 3/2020

Walgreen Company