CAREONE CHILDRENS IBUPROFEN- ibuprofen tablet, chewable American Sales Company

American Sales Company Children's Ibuprofen Drug Facts

Active ingredient (in each chewable tablet)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- · child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- · child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- chew or crush tablets completely before swallowing
- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than 4 times a day

Dosing Chart			
Weight (lb)	Age (yr)	Tablets	
under 24	under 2	ask a doctor	
24-35	2-3	1	
36-47	4-5	1 ½	
48-59	6-8	2	
60-71	9-10	2 ½	
72-95	11	3	

Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between 20-25°C (68-77°F)
- do not use if printed seal under cap is broken or missing

Inactive ingredients

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

Questions or comments?

1-800-719-9260

Principal Display Panel

CAREone[®]

For Ages 2 to 11

Children's IBUPROFEN Chewable Tablets, 100 mg

Pain Reliever/Fever Reducer (NSAID)

Chew or crush tablets completely before swallowing

Actual Size

OUR PHARMACISTS RECOMMEND

Lasts up to 8hrs

24 TABLETS

Grape-Flavored Chewables







Important: Read all product information before using. Keep this boxfor important information. This product is for children only.

Active ing redient (in each chewable tablet) Purposes buprofen 100 mg (NSAID)*......Pah reliever fever reducer *nonsteroidal anti-inflammatorydrug

Uses temporarily: ■ reduces fever
■ relieves minor aches and pains due to the common cold, flu, sore
throat, headaches and toothaches

Warnings

Warnings

Warnings

A lergy a lert i buprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

In hives Infacial swelling Instrima (wheeting)

In shock Inskin eddening Instrima (wheeting)

In an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAD, which may cause severe stomach bleeding. The chance is higher if your child:

In takes a blood thinning (anficcegularn) or steroid drug

Itakes other drugs containing pescription or nonprescription NSADe(saspirin, bupporen, nappowen, or others)

NS AIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed

Heart a ttack and stroke warming: NSA Ds, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Sore throat warning: Severe or persistent sore throat or sole throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Drug Facts (continued)

Do notuse

■ if the child has ever had an allergic reaction to ibuprofenor any other pain reliever/fever reduces ■right before or after heart surgery

Ask a doctor be fore use if

- stomach bleeding warning applies to your child child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- or rever reducers

 child has not been drinking fluids

 child has lost a bt of fluid due to vomiting or diarrhea
- child has high b bod pressure, heart disease, liver cirrhosis, kidney ase, or had a stroke

■child has asthma ■child is taking a diuretic

Ask a doctor or pharmacist before use if the child is ■under a doctor's care for any serious condition ■taking any other drug

When using this product

mouth or throat burning may occur; give with food or water
take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■child experiences any of the following signs of stomach bleeding: ■ feels faint ■ vormits blood ■ has bloody or black stools ■ has stomach poin that does not get better

■ child has symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling

Drug Facts (continued)

- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area.
- ■any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contacta Poison Control Center right away. (1-800-222-1222)

Directions

- chewor crush tablets completely before swallowing
 this product does not contain directions or complete warnings for adult use
- do not give more than directed ■find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ if needed, repeat dose every 6-8 hours
- do notuse more than 4 times a day

	Dooling Oliait	
Weight (lb)	Age (yr)	Tablets
under 24	under 2	aska doctor
24-35	2-3	-1
36-47	4-5	11/2
48 - 59	6-8	2
60-71	9-10	21/2
72-95	11	3

Drug Fads (continued)
Other information
phenykebounds:contains phenylatanine 6 mg per tablet
Estate between 20:25 °C (6 8-77 F)
on of use if printed seal under cap is broken ormissing hacthie ingredients aceatame potasium, annorium glysyribin, agariame, carriaba war, o ocamellesescium, post jet in. 27 duminum hke, FD8C bue no 1 duminum hke, hpromelloes, magnesium da rata, mamid, naturalandar fitidal flavors, slicondiaribe, accilum laur yis utate, sopeanol, succintracti Questions or comments ?1-800-719-9260 ute n Free ST RBU TEDBY NDOVER, MD2 019 S & Brands 7 FOO DHOLDU.S.A., LLC 2 078 5 · 1-877 - 846-9949 IS, LLC

CAREONE CHILDRENS IBUPROFEN

ibuprofen tablet, chewable

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:41520-633

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
W W W W W W W W W W W W W W W W W W W	IDLIBBOTEN	100

IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) IBUPROFEN 100 mg

Inactive Ingredients

mactive ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
ASPARTAME (UNII: Z0H242BBR1)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 30WL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SOYBEAN OIL (UNII: 241ATL177A)	
SUCCINIC ACID (UNII: AB6MNQ6J6L)	

Product Characteristics			
Color	PURPLE (Lavender)	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor	GRAPE	Imprint Code	L521
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-633- 62	1 in 1 CARTON	07/08/2019	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		

pplication Number or Monograph Citation	Marketing Start Date	Marketing End Date
076359	07/08/2019	
	Citation	Citation Date

Labeler - American Sales Company (809183973)

Revised: 10/2023 American Sales Company