

CLOBETASOL PROPIONATE CREAM USP, 0.05% CLOBETASOL PROPIONATE CREAM USP, 0.05%- clobetasol propionate cream usp, 0.05% cream
Encube Ethicals Private Limited

Rx only


FOR TOPICAL DERMATOLOGIC USE ONLY.

NOT FOR OPHTHALMIC, ORAL, OR INTRAVAGINAL USE.

DESCRIPTION:

Clobetasol propionate cream USP contains the active compound clobetasol propionate, a synthetic corticosteroid, for topical dermatologic use. Clobetasol, an analog of prednisolone, has a high degree of glucocorticoid activity and a slight degree of mineralocorticoid activity.

Chemically, clobetasol propionate is (11 β , 16 β)-21-chloro-9-fluoro-11-hydroxy-16-methyl-17-(1-oxopropoxy)-pregna-1,4-diene-3,20-dione, and it has the following structural formula:



Clobetasol propionate has the empirical formula $C_{25}H_{32}ClFO_5$ and a molecular weight of 467. It is a white to cream-colored crystalline powder insoluble in water.

Clobetasol propionate cream USP, 0.05% contains clobetasol propionate 0.5 mg/g in a cream base of propylene glycol, mono- and di-glycerides, cetostearyl alcohol, stearyl macroglycerides, white wax, chlorocresol, sodium citrate, anhydrous citric acid, and purified water.

CLINICAL PHARMACOLOGY:

Like other topical corticosteroids, clobetasol propionate has anti-inflammatory, antipruritic, and vasoconstrictive properties. The mechanism of the anti-inflammatory activity of the topical steroids, in general, is unclear. However, corticosteroids are thought to act by the induction of phospholipase A2 inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid.

Arachidonic acid is released from membrane phospholipids by phospholipase A2.

Pharmacokinetics:

The extent of percutaneous absorption of topical corticosteroids is determined by

many factors, including the vehicle and the integrity of the epidermal barrier. Occlusive dressings with hydrocortisone for up to 24 hours has not been demonstrated to increase penetration; however, occlusion of hydrocortisone for 96 hours markedly enhances penetration. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin may increase percutaneous absorption.

Studies performed with clobetasol propionate cream indicate that it is in the super-high range of potency as compared with other topical corticosteroids.

INDICATIONS AND USAGE:

Clobetasol propionate cream USP, 0.05% is super-high potency corticosteroid formulation indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses. Treatment beyond 2 consecutive weeks is not recommended, and the total dosage should not exceed 50 g /week because of the potential for the drug to suppress the hypothalamic-pituitary- adrenal (HPA) axis. Use in pediatric patients under 12 years of age is not recommended. As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of the diagnosis may be necessary.

CONTRAINDICATIONS:

Clobetasol propionate cream USP, 0.05% is contraindicated in those patients with a history of hypersensitivity to any of the components of the preparations.

PRECAUTIONS

General:

Clobetasol propionate cream USP, 0.05% should not be used in the treatment of rosacea or perioral dermatitis, and should not be used on the face, groin, or axillae.

Systemic absorption of topical corticosteroids can produce reversible HPA axis suppression with the potential for glucocorticosteroid insufficiency after withdrawal from treatment. Manifestations of Cushing's syndrome, hyperglycemia, and glucosuria can also be produced in some patients by systemic absorption of topical corticosteroids while on therapy.

Patients applying a topical steroid to a large surface area or to areas under occlusion should be evaluated periodically for evidence of HPA axis suppression. This may be done by using the ACTH stimulation, A.M. plasma cortisol, and urinary free cortisol tests. Patients receiving super-potent corticosteroids should not be treated for more than 2 weeks at a time, and only small areas should be treated at any one time due to the increased risk of HPA suppression.

Clobetasol propionate cream produced HPA axis suppression when used at doses as low as 2 g/day for 1 week in patients with eczema.

If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent corticosteroid. Recovery of HPA axis function is generally prompt upon discontinuation of topical corticosteroids. Infrequently, signs and symptoms of glucocorticosteroid insufficiency may occur that require supplemental systemic corticosteroids. For information on systemic supplementation, see prescribing information for those products.

Pediatric patients may be more susceptible to systemic toxicity from equivalent doses due to their larger skin surface to body mass ratios (see PRECAUTIONS: Pediatric Use).

If irritation develops, clobetasol propionate cream should be discontinued and appropriate therapy instituted. Allergic contact dermatitis with corticosteroids is usually diagnosed by observing a failure to heal rather than noting a clinical exacerbation as with most topical products not containing corticosteroids. Such an observation should be corroborated with appropriate diagnostic patch testing.

If concomitant skin infections are present or develop, an appropriate antifungal or antibacterial agent should be used. If a favorable response does not occur promptly, use of clobetasol propionate cream should be discontinued until the infection has been adequately controlled.

Information for Patients:

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. The treated skin area should not be bandaged, otherwise covered, or wrapped so as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions to the physician.

Laboratory Tests:

The following tests may be helpful in evaluating patients for HPA axis suppression:

ACTH stimulation test
A.M. plasma cortisol test
Urinary free cortisol test

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Long-term animal studies have not been performed to evaluate the carcinogenic potential of clobetasol propionate.

Studies in the rat following subcutaneous administration at dosage levels up to 50 mcg/kg per day revealed that the females exhibited an increase in the number of resorbed embryos and a decrease in the number of living fetuses at the highest dose.

Clobetasol propionate was nonmutagenic in 3 different test systems: the Ames test, the *Saccharomyces cerevisiae* gene conversion assay, and the *E. coli* B WP2 fluctuation test.

Pregnancy: Teratogenic Effects:

Corticosteroids have been shown to be teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Some corticosteroids have been shown to be teratogenic after dermal application to laboratory animals.

Clobetasol propionate has not been tested for teratogenicity when applied topically; however, it is absorbed percutaneously, and when administered subcutaneously it was a significant teratogen in both the rabbit and mouse. Clobetasol propionate has greater teratogenic potential than steroids that are less potent.

Teratogenicity studies in mice using the subcutaneous route resulted in fetotoxicity at the highest dose tested (1 mg/kg) and teratogenicity at all dose levels tested down to 0.03 mg/kg. These doses are approximately 1.4 and 0.04 times, respectively, the human topical dose of clobetasol propionate cream. Abnormalities seen included cleft palate and skeletal abnormalities.

In rabbits, clobetasol propionate was teratogenic at doses of 3 and 10 mcg/kg. These doses are approximately 0.02 and 0.05 times, respectively, the human topical dose of clobetasol propionate cream. Abnormalities seen included cleft palate, cranioschisis, and other skeletal abnormalities.

There are no adequate and well-controlled studies of the teratogenic potential of clobetasol propionate in pregnant women. Clobetasol propionate cream should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers:

Systemically administered corticosteroids appear in human milk and could suppress growth, interfere with endogenous corticosteroid production, or cause other untoward effects. It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in human milk. Because many drugs are excreted in human milk, caution should be exercised when clobetasol propionate cream is administered to a nursing woman.

Pediatric Use:

Safety and effectiveness of clobetasol propionate cream in pediatric patients have not been established. Use in pediatric patients under 12 years of age is not recommended. Because of a higher ratio of skin surface area to body mass, pediatric patients are at a greater risk than adults of HPA axis suppression and Cushing's syndrome when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency during or after withdrawal of treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children.

HPA axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain, and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations of adrenal suppression in children include low plasma cortisol levels and an absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Geriatric Use:

A limited number of patients at or above 65 years of age have been treated with clobetasol propionate cream (n = 231) in US and non-US clinical trials. While the number of patients is too small to permit separate analysis of efficacy and safety, the adverse reactions reported in this population were similar to those reported by younger patients. Based on available data, no adjustment of dosage of clobetasol propionate cream in geriatric patients is warranted.

ADVERSE REACTIONS:

In controlled clinical trials, the most frequent adverse reactions reported for clobetasol propionate cream were burning and stinging sensation in 1% of treated patients. Less frequent adverse reactions were itching, skin atrophy, and cracking and fissuring of the skin.

Cushing's syndrome has been reported in infants and adults as a result of prolonged use of topical clobetasol propionate formulations.

The following additional local adverse reactions have been reported with topical corticosteroids, and they may occur more frequently with the use of occlusive dressings and higher potency corticosteroids. These reactions are listed in an approximately decreasing order of occurrence: dryness, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, irritation, striae, and miliaria. To report SUSPECTED ADVERSE REACTIONS, contact Encube Ethicals Private Limited at 1-833-285-4151 or FDA at 1-800-FDA-1088

or www.fda.gov/medwatch.

OVERDOSAGE:

Topically applied clobetasol propionate cream can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

DOSAGE AND ADMINISTRATION:

Apply a thin layer of clobetasol propionate cream to the affected skin areas twice daily and rub in gently and completely. (See INDICATIONS AND USAGE).

Clobetasol propionate cream is super- high potency topical corticosteroids; therefore, **treatment should be limited to 2 consecutive weeks and amounts greater than 50 g / week should not be used.**

As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of diagnosis may be necessary.

Clobetasol propionate cream should not be used with occlusive dressings.

Geriatric Use: In studies where geriatric patients (65 years of age or older, see PRECAUTIONS) have been treated with clobetasol propionate cream, safety did not differ from that in younger patients; therefore, no dosage adjustment is recommended.

HOW SUPPLIED:

Clobetasol Propionate Cream USP, 0.05% is supplied in:

15 g tube (NDC 21922-016-04)

30 g tube (NDC 21922-016-05)

45 g tube (NDC 21922-016-06)

60 g tube (NDC 21922-016-07)

Store cream between 15° and 30°C (59°-86°F). Clobetasol propionate cream should not refrigerated.

Manufactured by:

Encube Ethicals Pvt. Ltd.

Plot No. C-1, Madkaim Industrial Estate,

Madkaim, Post: Mardol,

Ponda, Goa - 403 404,

India.

Distributed by:

Encube Ethicals Inc.

200 Meredith Avenue,

Suite 101A

Durham, NC 27713

USA

Rev: 00

Sep 2020

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

PRINCIPAL DISPLAY PANEL-Carton


NDC 21922-016-04

Clobetasol Propionate Cream USP, 0.05%

For dermatologic use only.

Not for ophthalmic use.

Rx only 15 grams



PRINCIPAL DISPLAY PANEL-Label

NDC 21922-016-04

Clobetasol Propionate Cream USP, 0.05%

For dermatologic use only.

Not for ophthalmic use.

Rx only 15 grams

NDC 21922-016-04

Clobetasol Propionate Cream USP, 0.05%

Encube

15 grams

**For dermatologic use only.
Not for ophthalmic use.**

EACH GRAM CONTAINS: 0.5 mg clobetasol propionate in a cream base composed of propylene glycol, mono- and di-glycerides, cetostearyl alcohol, stearyl macroglycerides, white wax, chlorocresol, sodium citrate, citric acid, and purified water;

USUAL DOSAGE: See package insert for full prescribing information.


WARNING : Keep out of reach of children.

**Store cream between 15°C and 30°C (59° and 86°F).
Clobetasol propionate cream should not be refrigerated.
See crimp for lot no. and expiration date.**

IMPORTANT: Do not use if seal has been punctured or is not visible.

TO OPEN: Use cap to puncture seal

ECC1A/00



N
3 2 1 9 2 2 1 0 1 6 0 4 1 8


Mfg. Lic. No.: 361

Manufactured by: **Encube Ethicals Pvt, Ltd,**
Plot No, C-1, Madkaim Industrial Estate, Madkaim,
Post: Mardol, Ponda, Goa - 403 404, India.

Distributed by: **Encube Ethicals, Inc.**
200 Meredith Avenue, Suite 101A
Durham, NC 27713 USA



PRINCIPAL DISPLAY PANEL-Carton NDC 21922-016-05 Clobetasol Propionate Cream USP, 0.05% For dermatologic use only. Not for ophthalmic use. Rx only 30 grams



PRINCIPAL DISPLAY PANEL-Label
NDC 21922-016-05
Clobetasol Propionate Cream USP, 0.05%
For dermatologic use only.
Not for ophthalmic use.
Rx only 30 grams

NDC 21922-016-05

Clobetasol Propionate
Cream USP, 0.05%

30 grams

For dermatologic use only.
Not for ophthalmic use.

EACH GRAM CONTAINS: 0.5 mg clobetasol propionate in a cream base composed of propylene glycol, mono- and di-glycerides, cetostearyl alcohol, stearyl macroglycerides, white wax, chlorocresol, sodium citrate, citric acid, and purified water.

USUAL DOSAGE: See package insert for full prescribing information.

WARNING: Keep out of reach of children.

Store cream between 15°C and 30°C (59° and 86°F).
Clobetasol propionate cream should not be refrigerated.

See crimp for lot no, and expiration date.


IMPORTANT: Do not use if seal has been punctured or is not visible.

TO OPEN: Use cap to puncture seal.

Mfg. Lic. No.: 361 ECC1B/00

Encube


Rx only




N
3 21922 01605 5

Manufactured by:
Encube Ethicals Pvt, Ltd.
Plot No. C-1, Madkaim Industrial Estate,
Madkaim, Post: Mardol, Ponda, Goa - 403 404, India.

Distributed by:
Encube Ethicals, Inc.
200 Meredith Avenue, Suite 101A
Durham, NC 27713 USA



PRINCIPAL DISPLAY PANEL-Carton
NDC 21922-016-06
Clobetasol Propionate Cream USP, 0.05%
For dermatologic use only.
Not for ophthalmic use.
Rx only 45 grams



PRINCIPAL DISPLAY PANEL-Label
NDC 21922-016-06
Clobetasol Propionate Cream USP, 0.05%
For dermatologic use only.
Not for ophthalmic use.
Rx only 45 grams

NDC 21922-016-06

Encube

Clobetasol Propionate

Cream USP, 0.05%

45 grams

**For dermatologic use only.
Not for ophthalmic use.**

EACH GRAM CONTAINS: 0.5 mg clobetasol propionate in a cream base composed of propylene glycol, mono- and di-glycerides, cetostearyl alcohol, stearyl macroglycerides, white wax, chlorocresol, sodium citrate, citric acid, and purified water.

USUAL DOSAGE: See package insert for full prescribing information.

WARNING: Keep out of reach of children.

**Store cream between 15°C and 30°C (59° and 86°F).
Clobetasol propionate cream should not be refrigerated.**

See crimp for lot no. and expiration date.

IMPORTANT: Do not use if seal has been punctured or is not visible.

TO OPEN: Use cap to puncture seal.

Mfg. Lic. No.: 361

ECC1C/00



Manufactured by:
Encube Ethicals Pvt. Ltd.
Plot No. C-1, Madkaim Industrial Estate,
Madkaim, Post: Mardol, Ponda,
Goa - 403 404, India.

Distributed by:
Encube Ethicals, Inc.
200 Meredith Avenue, Suite 101A
Durham, NC 27713 USA

PRINCIPAL DISPLAY PANEL-Carton


NDC 21922-016-07

Clobetasol Propionate Cream USP, 0.05%

For dermatologic use only.

Not for ophthalmic use.

Rx only 60 grams



PRINCIPAL DISPLAY PANEL-Label
NDC 21922-016-07
Clobetasol Propionate Cream USP, 0.05%
For dermatologic use only.
Not for ophthalmic use.
Rx only 60 grams

NDC 21922-016-07


Encube

Clobetasol Propionate

Cream USP, 0.05%

60 grams

**For dermatologic use only.
Not for ophthalmic use.**



EACH GRAM CONTAINS: 0.5 mg clobetasol propionate in a cream base composed of propylene glycol, mono- and di-glycerides, cetostearyl alcohol, stearyl macrogolglycerides, white wax, chlorocresol, sodium citrate, citric acid, and purified water.

USUAL DOSAGE: See package insert for full prescribing information.

WARNING: Keep out of reach of children.

**Store cream between 15°C and 30°C (59° and 86°F).
Clobetasol propionate cream should not be refrigerated.**

See crimp for lot no. and expiration date.

IMPORTANT: Do not use if seal has been punctured or is not visible.

TO OPEN: Use cap to puncture seal.

ECC1D/00



Mfg. Lic. No.: 361

Manufactured by:
Encube Ethicals Pvt. Ltd.
Plot No. C-1, Madkaim Industrial Estate,
Madkaim, Post: Mardol, Ponda,
Goa - 403 404, India.

Distributed by:
Encube Ethicals, Inc.
200 Meredith Avenue, Suite 101A
Durham, NC 27713 USA



CLOBETASOL PROPIONATE CREAM USP, 0.05% CLOBETASOL PROPIONATE CREAM USP, 0.05%

clobetasol propionate cream usp, 0.05% cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:21922-016
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLOBETASOL PROPIONATE (UNII: 779619577M) (CLOBETASOL - UNII:ADN79D536H)	CLOBETASOL PROPIONATE	0.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)	
GLYCERYL STEARATE/PEG-100 STEARATE (UNII: RD25J5V947)	
WHITE WAX (UNII: 7G1J5DA97F)	

CHLOROCRESOL (UNII: 36W53O7109)	
WATER (UNII: 059QF0K00R)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
ANHYDROUS TRISODIUM CITRATE (UNII: RS7A450LGA)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21922-016-04	15 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2020	
2	NDC:21922-016-05	30 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2020	
3	NDC:21922-016-06	45 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2020	
4	NDC:21922-016-07	60 g in 1 TUBE; Type 0: Not a Combination Product	10/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212982	10/01/2020	

Labeler - Encube Ethicals Private Limited (915834105)

Registrant - Encube Ethicals Private Limited (915834105)

Establishment

Name	Address	ID/FEI	Business Operations
Encube Ethicals Private Limited		725076298	ANALYSIS(21922-016) , LABEL(21922-016) , MANUFACTURE(21922-016) , PACK(21922-016) , PARTICLE SIZE REDUCTION(21922-016)

Revised: 10/2020

Encube Ethicals Private Limited