

AUTO-CHLOR BASE- sodium chlorite liquid IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Auto-Chlor Base

Active Ingredient

Sodium Chlorite 80%...0.5%

Inert Ingredients

Stabilizers & Water

Mixing Directions

1. It is recommended to wear nitrile or rubber gloves when handling a concentrate to avoid staining hands.
2. Always mix solutions into a clean, well labeled, non-metallic, empty container.
3. Fill empty container with proper amount of cold, soft water.
4. Add proper amount of Duet Base to container as directed.
5. Add proper amount of Duet Activator to container as directed.
6. Mix in a well ventilated area and avoid breathing fumes.
7. Use mixed solution within 12 hours.
8. When Duet Base and Duet Activator are mixed properly, according to the label directions, a solution of acidified sodium chlorite is achieved in the ready-to-use product.

Recommended Procedures for Teat Dip Sanitation

A. Pre-Milking (Pre-Dipping)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with Ready-to-Use Duet Teat Dip.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, test orifices & udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. Post-Milking (Post-Dipping)

Immediately after milking, dip or spray all teats full length with Ready-to-Use Duet Teat Dip. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

Warning

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.
5. Protect this product from freezing.

Danger

Causes severe skin burns & eye damage.

Harmful if swallowed.

May be corrosive to metals.

First Aid

If in Eyes: Rinse cautiously with water for several minutes. Remove contact lenses if present & easy to do. Continue rinsing for 15 minutes. Get immediate medical advice/attention.

If Swallowed: Rinse mouth. Do not induce vomiting. Get immediate medical advice/attention.

If on Skin: Immediately remove all contaminated clothing. Wash with plenty of water for 15 minutes. If skin irritation occurs, get medical advice/attention. Wash contaminated clothing before use.

If Inhaled: If breathing is difficult, remove victim to fresh air & keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms get medical advice/attention immediately.

Caution

Keep out of reach of children.

Auto-Chlor Base

NDC 29983-1096-8



**NPE
FREE**

AUTO-CHLOR

**NPE
FREE**

A CONCENTRATED SANITIZING PRE-MILKING AND POST-MILKING TEAT DIP A HIGHLY EFFECTIVE GERMICIDAL NON-IODINE FORMULATION

ACTIVE INGREDIENT:

Contains 7.05% Sodium Chlorite 78%
(Provides A Full 5.5% Available Sodium Chlorite).....5.5%

INERT INGREDIENTS: Stabilizers and Water

MIXING DIRECTIONS

1. It is recommended to wear nitrile or rubber gloves when handling this concentrate to avoid staining hands.
2. Always mix solutions into a clean, well labeled, non-metallic, empty container.
3. Fill empty container with proper amount of cold, soft water.
4. Add proper amount of AUTO-CHLOR Base to container as directed.
5. Add proper amount of AUTO-CHLOR Activator to container as directed. Mix until the color is uniform throughout.
6. Mix in a well ventilated area and avoid breathing fumes.
7. Use mixed solution within 12 hours.
8. When AUTO-CHLOR BASE and AUTO-CHLOR ACTIVATOR are mixed properly according to label directions a solution of acidified sodium chlorite is achieved in the ready-to-use product.

AUTO-CHLOR MIXING GUIDE					
READY-TO-USE VOLUME DESIRED	TARGET PPM ACIDIFIED SODIUM CHLORITE	WATER	AUTO-CHLOR ACTIVATOR	AUTO-CHLOR BASE	
		OZ.	OZ.	OZ.	
1 GALLON / 3.8 LITERS	1500 3000	120 112	4 8	4 8	
1 QUART / .95 LITERS	1500 3000	30 28	1 2	1 2	

RECOMMENDED PROCEDURES FOR TEAT SANITATION


A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip, foam or spray the cow's teats full length with READY-TO-USE AUTO-CHLOR TEAT DIP.
4. Wait at least 15-30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

Immediately after milking, dip, foam or spray all teats full length with READY-TO-USE AUTO-CHLOR TEAT DIP. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

BASE FOR USE ONLY WITH AUTO-CHLOR ACTIVATOR



WARNING

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.
5. Protect this product from freezing.

DANGER

AUTO-CHLOR Base **CONCENTRATE** causes severe skin burns and eye damage.

Harmful if swallowed.

May be corrosive to metals.



Avoid breathing dust/fume/gas/mist/vapors/spray. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Do not eat, drink or smoke when using this product.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing for 15 minutes. Get immediate medical advice/attention.

IF SWALLOWED: Rinse mouth. Do not induce vomiting. Get immediate medical advice/attention.

IF ON SKIN: Immediately remove all contaminated clothing. Wash skin with plenty of water for 15 minutes. If skin irritation occurs, get medical advice/attention. Wash contaminated clothing before reuse.

IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms get medical advice/attention immediately.

MEDICAL and TRANSPORTATION EMERGENCIES contact CHEMTREC 1-800-424-9300

A PART OF THE COMPLETE IBA SANITATION PROGRAM

UN 3266

Corrosive liquid, basic, inorganic, n.o.s. (Sodium Chlorite), 8, PG II

EXP. DATE:

LOT NO.:

NET CONTENTS: 55 GALLONS

IBA STOCK #261007

1224W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

AUTO-CHLOR BASE

sodium chlorite liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1096
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H3745B6)	SODIUM CHLORITE	0.5 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1096-9	1125 kg in 1 CONTAINER		

2	NDC:29983-1096-8	222.5 kg in 1 DRUM		
3	NDC:29983-1096-6	60.68 kg in 1 DRUM		
4	NDC:29983-1096-4	20.23 kg in 1 PAIL		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/24/2024	

Labeler - IBA (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA in Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 11/2025

IBA