

SOLBAR SHIELD SPF40- solbar shield spf40 cream

Person and Covey

Solbar Shield SP40

Indications and use

Helps prevent sunburn. If used as directed with other sun protection measures, decreases the risk of skin cancer and early skin aging caused by the sun. Dosage and Administration: Apply liberally and evenly to all sun exposed areas of DRY skin 15 minutes before sun exposure. Reapply after 80 minutes of swimming or sweating and immediately after towel drying. Apply at least every 2 hours. For children under 6 months, ask a physician.

Purpose

Sunscreen

Keep out of the reach of children

Yes. If swallowed, get medical help or contact a Poison Control Center right away.

Dosage and Administration

Apply liberally and evenly to all sun exposed areas of DRY skin 15 minutes before sun exposure. Reapply after 80 minutes of swimming or sweating and immediately after towel drying. Apply at least every 2 hours. For children under 6 months, ask a physician.

Warnings

For external use only. Do not use on damaged or broken skin. Keep out of eyes. Rinse eyes thoroughly with water to remove. Stop use and ask a physician if rash or irritation develops and lasts. Store away from excessive heat and direct sun.

OTC - ACTIVE INGREDIENT SECTION

Homosalate

Octinoxate

Oxybenzone

Avobenzone

INACTIVE INGREDIENT SECTION

Water

Isobutyl Stearate
 Glycerin
 Benzyl Alcohol
 Dimethicone
 Cetyl Phosphate
 Triethanolamine
 Stearic Acid
 Silica
 Carbomer 1342
 Acrylates/C10-30 Alkyl Acrylate Crosspolymer
 Disodium EDTA

SOLBAR[®]
SUNSCREEN

40

SHIELD

BROAD SPECTRUM SPF 40 WITH
 ZINC OXIDE & TITANIUM DIOXIDE

Water Resistant (80 Mins.)

MINERAL BASED

Enriched with aloe and Vitamin E

PC Person & Covey, Inc. Glendale, California 91201, U.S.A. MADE IN USA personandcovey.com 8T

Drug Facts

Active Ingredient	Purpose
Zinc Oxide (Z-Cote [®] HPI) 7.5%	Sunscreen
Titanium Dioxide 5.0%	Sunscreen

Use SOLBAR[®] Shield Sunscreen helps prevent sunburn if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings For external use only

Do not use on damaged or broken skin

When using this product
 keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions apply liberally and evenly 15 minutes before sun exposure. Apply to all skin exposed to the sun. **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m.-2 p.m. wear long-sleeved shirts, pants, hats, and sunglasses reapply after 80 minutes of swimming or sweating immediately after towel drying at least every 2 hours children under 6 months: Ask a doctor.

Other Information store away from excessive heat or cold.

Inactive Ingredients
 Water (Aqua), Glycerin, Stearate, Oleth-3 Phosphate, PVP/Hexadecene Copolymer, Dimethicone, Isobutyl Stearate, C12-15 Alkyl Benzoate, Cetearyl Ethylhexanoate (and) Isopropyl Myristate, Glycerin, Benzyl Alcohol, Glyceryl Stearate (and) PEG-100 Stearate, Tocopheryl Acetate, Xanthan Gum, Aminomethyl Propanol, Disodium EDTA, Aloe Barbadensis Leaf Extract, Citric Acid

Questions? Please call 1 (800) 423-2341

SOLBAR SHIELD SPF40			
solbar shield spf40 cream			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0096-0682
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.05 g in 1 g
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	0.075 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
GLYCERYL STEARATE SE (UNII: FCZ5MH785I)	
PEG-100 STEARATE (UNII: YD01N1999R)	
.ALPHA.-TOCOPHEROL, DL- (UNII: 7QWA1RIO01)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
XANTHAN GUM (UNII: TTV12P4NEE)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
ISOBUTYL STEARATE (UNII: V8DPR6HNX3)	
CETEARYL ETHYLHEXANOATE (UNII: 9M64UO4C25)	
ISOPROPYL MYRISTATE (UNII: ORE8K4LNJS)	
C12-20 ALKYL BENZOATE (UNII: Y15I6XI14C)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
WATER (UNII: 059QF0KO0R)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
DIETHANOLAMINE OLETH-3 PHOSPHATE (UNII: Y67NX5905E)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
GLYCERIN (UNII: PDC6A3C0OX)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0096-0682-04	129 g in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2000	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	06/01/1996	

Labeler - Person and Covey (008482473)

Establishment

Name	Address	ID/FEI	Business Operations
Person and Covey		008482473	manufacture(0096-0682)

