

**ANTACID- aluminum hydroxide, magnesium hydroxide, simethicone liquid**  
**Rugby Laboratories**

-----

**Rugby Regular Strength Antacid Liquid**

**ACTIVE INGREDIENTS (in each 10 mL )**

Aluminum hydroxide (equiv. to dried gel, USP) 400 mg  
Magnesium hydroxide 400 mg  
Simethicone 40 mg

**PURPOSE**

Antacid  
Antacid  
Antigas

**USE(S)**

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

**WARNINGS**

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

**ASK A DOCTOR BEFORE USE IF YOU HAVE**

- kidney disease
- a magnesium-restricted diet

**ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**STOP USE AND ASK DOCTOR**

if symptoms last more than two weeks

## **KEEP OUT OF REACH OF CHILDREN**

### **DIRECTIONS**

- **shake well before use**
- mL = milliliter
- adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor
- children under 12 years: consult a doctor

### **OTHER INFORMATION**

- **each 10 mL contains:** magnesium 170 mg, sodium 5 mg
- store at controlled room temperature 20°C-25°C (68°-77°F)
- do not freeze

### **INACTIVE INGREDIENTS**

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

### **PRINCIPAL DISPLAY PANEL**

**ugby**

NDC 0536-1293-83

#### **Regular Strength**

#### **Antacid Liquid**

Aluminum hydroxide, 400 mg

Magnesium hydroxide, 400 mg

Simethicone, 40 mg

#### **Antacid and Antigas**

Original Flavor

#### **Relief of:**

Heartburn

Acid Indigestion

Sour Stomach

Pressure & Bloating

**12 FL OZ (355 mL)** Alcohol content 0.2% v/v



NDC 0536-1293-83

Regular Strength

# Antacid Liquid

Aluminum Hydroxide, 400 mg  
Magnesium Hydroxide, 400 mg  
Simethicone, 40 mgAntacid and Antigas  
Original FlavorRelief of:  
Heartburn  
Acid Indigestion  
Sour Stomach  
Pressure and Bloating**12 FL OZ (355 mL)**

Alcohol Content 0.2% v/v

**Drug Facts** **DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

Active ingredients (in each 10 mL)	Purpose
Aluminum hydroxide (equiv. to dried gel, USP) 400 mg	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

**Uses** relieves: ■ acid indigestion ■ heartburn ■ sour stomach  
■ upset stomach and gas associated with these symptoms

**Warnings**

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Ask a doctor before use if you have ■ kidney disease  
■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than two weeks.

Keep out of reach of children.

**Directions**■ shake well before use ■ mL = milliliter  
■ adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor ■ children under 12 years: consult a doctor

**Other information** ■ each 10 mL contains: magnesium 170 mg, sodium 5 mg ■ store at controlled room temperature 20°C-25°C (68°F-77°F) ■ do not freeze

**Inactive ingredients** ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

Distributed by:  
RUGBY® LABORATORIES  
Livonia, MI 48152Questions or comments?  
Call 1-800-645-2158  
www.rugbylaboratories.com

Rev. 05/21 8-30 Re-order No. 271027 MFR# 53041



3 05361 29383 3

## ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0536-1293
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 10 mL
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 10 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYDROXYETHYL CELLULOSE, UNSPECIFIED</b> (UNII: T4V6TWG28D)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	

<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1293-83	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/02/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	06/02/2021	

**Labeler** - Rugby Laboratories (079246066)

### Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(0536-1293)

Revised: 12/2024

Rugby Laboratories