**ACETAMINOPHEN- acetaminophen tablet**
**Rising Pharma Holdings, Inc.**

---

**Compare to active ingredient in Tylenol® Regular Strength†**

**Acetaminophen**
**325 mg**

*** Pain Reliever**
*** Fever Reducer**
*** Contains No Aspirin**

**100 Film Coated Tablets**

**DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

## *Drug Facts*

***Active Ingredient (in each tablet)*** ***Purpose***
Acetaminophen USP, 325 mg.....Pain Reliever/Fever Reducer

## *Uses*

To reduce fever and for the temporary relief of minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain from arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever.

## *Warnings*

**Liver warning**: This product contains acetaminophen.

Severe liver damage may occur if

- adult takes more than 10 tablets in 24 hours, which is the maximum daily amount
- child takes more than 5 tablets in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have** liver disease.

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

• pain gets worse or lasts more than 10 days in adults and children.
• pain gets worse or lasts more than 5 days in children under 12 years.
• fever gets worse or lasts more than 3 days.
• new symptoms occur.
• redness or swelling is present.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

**Do not take more than directed**

**Adults and Children 12 years and over:**

- take 2 tablets every 4 to 6 hours while symptoms last.
- do not take more than 10 tablets in 24 hours.
- do not take for more than 10 days unless directed by a doctor

**Children 6 years to under 12 years:**

- take 1 tablet every 4 to 6 hours while symptoms last.
- do not take more than 5 tablets in 24 hours.
- do not use for more than 5 days unless directed by a doctor

**Children under 6 years:**

ask a doctor

**Other Information**

- SODIUM FREE
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- see end panel for lot number and expiration date.

**Inactive Ingredients**

Hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide.

**Questions and Comments?** Call **1-844-474-7464**

**Distributed by:**

Rising Pharma Holdings, Inc.

East Brunswick, NJ 08816

Made in India

Mfg. Lic. No.: G/25/2258

**Issued:** 06/2025

S-065

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Tylenol® Regular strength.

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**Acetaminophen**

**325 mg**

**100 Film Coated Tablets**







# ACETAMINOPHEN

acetaminophen tablet

| Product Information | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:57237-346 |
| **Route of Administration** | ORAL | | |

| Active Ingredient/Active Moiety | | |
|---|---|---|
| **Ingredient Name** | **Basis of Strength** | **Strength** |
| **ACETAMINOPHEN** (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |

| Inactive Ingredients | |
|---|---|
| **Ingredient Name** | **Strength** |
| **HYPROMELLOSE, UNSPECIFIED** (UNII: 3NXW29V3WO) | |
| **MAGNESIUM STEARATE** (UNII: 70097M6I30) | |

| | |
|---|---|
| **MICROCRYSTALLINE CELLULOSE** (UNII: OP1R32D61U) | |
| **POLYETHYLENE GLYCOL, UNSPECIFIED** (UNII: 3WQ0SDW1A) | |
| **POVIDONE** (UNII: FZ989GH94E) | |
| **STARCH, CORN** (UNII: O8232NY3SJ) | |
| **STEARIC ACID** (UNII: 4ELV7Z65AP) | |
| **TALC** (UNII: 7SEV7J4R1U) | |
| **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP) | |

## Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | white (Off White) | **Score** | no score |
| **Shape** | ROUND | **Size** | 10mm |
| **Flavor** | | **Imprint Code** | S99 |
| **Contains** | | | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:57237-346-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 07/09/2025 | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC Monograph Drug | M013 | 07/09/2025 | |

## Labeler - Rising Pharma Holdings, Inc. (116880195)

## Registrant - Elysium Pharmaceuticals Ltd (863182240)

## Establishment

| Name | Address | ID/FEI | Business Operations |
|---|---|---|---|
| Elysium Pharmaceuticals Ltd | | 863182240 | analysis(57237-346) , label(57237-346) , manufacture(57237-346) , pack(57237-346) |

Revised: 7/2025 Rising Pharma Holdings, Inc.