ACETAMINOPHEN- acetaminophen tablet, film coated, extended release CVS PHARMACY INC

Arthritis Pain Reliever

ACETAMINOPHEN EXTENDED-RELEASE TABLETS USP, 650 mg

PAIN RELIEVER / FEVER REDUCER

ARTHRITIS 8 HOURCAPLETS

- For the temporary relief of minor arthritis pain
- Lasts up to 8 hours

Active ingredient(in each caplet)

Acetaminophen USP, 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children

Keep out of reach of children

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ d o not take more than directed (see overdose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor under 18 years of age
- ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **■** do not use if foil inner seal is broken or missing

Inactive ingredients

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?

Contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST.

PDP



Inside (adhesive side)

"This product is not manufactured or distributed by Notked Consumer Healthcare, distributor of Tylend" SHR Arthritis Pain.

Stap use and ask a dooter if

pain gets worse or leafs more than 10 days
there gets worse or leafs more than 3 days Ask a doctor or pharmacist before use if you are taking the idlood thinning Drug Facts (continued) 2015c11-477-470-4183 Mon-Fit 8:00 AM EST to 5:00 PM PST collulare, polyethylane glyral, povidare, progréditized starch, actium starch lisik a doctor before use if you have liver disease hactive ingredients yeby etyl cellus, hyporelise, magresim sterck, nicocystellor new symptoms occur Byou are not sure whether a drug portains acetaminophen, ask a doctor or ther information store between 20-25°C (68-77°F) redness or swelling is present if you are allergic to exeterninghen or any of the insother ingredients in this with any other drug containing acetaminophen (prescription or norperscription endiose merming in case of overclose, get medical help or contact a Polson nimil Center right away. (1–303–272–1272) Cusick medical ablention is critical adults as well as for children awan in you do not notice any signs or symptoms. e could be signs of a serious condition, sgeant or breast-leading, ask a health professional before use, out of reach of children. stions or comme n, stears and ■ take 2 capales every 8 hours with water
■ swellow whole; do not crush, chew, spift or disordie
■ do not take more than 6 capales in 24 hours
■ do not use for more than 10 days unless directed by a doctor ask a doctor



ACETAMINOPHEN

acetaminophen tablet, film coated, extended release

		_
Product	+ Inform	ation
FIUUUC		alivii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-322

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D) ACETAMINOPHEN 650 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROXYETHYL CELLULOSE (140 MPA.S AT 5%) (UNII: 8136Y38GY5)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
STARCH, CORN (UNII: O8232NY3SJ)

SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)

STEARIC ACID (UNII: 4ELV7Z65AP)

MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	

Product Characteristics			
Color	white (White to off white colored)	Score	no score
Shape	OVAL (Capsule shaped, biconvex intact film coated tablets)	Size	19mm
Flavor		Imprint Code	G;650
Contains			

ı	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:69842-322- 24	24 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2021	
	NDC:69842-322-	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	02/01/2021	

Labeler - CVS PHARMACY INC (062312574)

Revised: 12/2023 CVS PHARMACY INC