

**CHLORHEXIDINE GLUCONATE- chlorhexidine gluconate cloth**  
**Sage Products LLC**

-----  
**Chlorhexidine Gluconate 2% Solution**

**2% CHLORHEXIDINE GLUCONATE CLOTH**

---

<b>Active Ingredient</b>	<b>Purpose</b>
Chlorhexidine Gluconate 2% Solution	Antiseptic

---

**Uses**

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

**Warnings**

**For external use only**

**Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar punctures or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product**

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- do not microwave
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.

## **To open package**

- holding top of package in one hand, lift flap on backside of package with other hand
- grasp flap at top and pull down to tear flap away and expose foam
- hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination

## **Or**

- using sterile scissors, cut off end seal of package
- transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- **dry surgical sites** (such as abdomen or arm): use one cloth to cleanse each 161 cm<sup>2</sup> area (approximately 5 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- **moist surgical sites** (such as inguinal fold): use one cloth to cleanse each 65 cm<sup>2</sup> area (approximately 2 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- discard each cloth after a single use
- after package has been opened discard any unused cloths

## **Other information**

store product flat

store between 20-25°C (68-77°F)

avoid excessive heat above 40°C (104°F)

## **□ Inactive Ingredients**

aloe vera, dimethicone, fragrance, glucono-delta-lactone, glycerin, nonoxynol-9, polysorbate 20, propylene glycol, USP purified water

cloth: polyester

## **Questions or comments?**

call toll-free 800-323-2220

(Monday to Friday 8 AM - 5 PM CST)

www.sageproducts.com

DO NOT FLUSH CLOTHS

Not made with natural rubber latex

Patents:

www.sageproducts.com/patents

2 pack

**Drug Facts (continued)**

**Directions**

- Using sterile scissors, cut off and seal of package
- transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- dry surgical sites (such as abdomen or arm): use one cloth to cleanse each 161 cm<sup>2</sup> area (approximately 5 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- moist surgical sites (such as inguinal fold): use one cloth to cleanse each 60 cm<sup>2</sup> area (approximately 2 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- discard each cloth after a single use
- if the package has been opened discard any unused cloths

**Other information**

- store product flat
- store between 20–25°C (66–77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

aloe vera, dimethicone, fragrance, glucono-delta-lactone, glycerin, nonoxonyl-8, polyisobutylene 20, propylene glycol, USP purified water, cloth: polyester

**Questions or comments?**  
call toll-free 800-323-2220 (Monday to Friday 8:00 am – 5:00 pm CST) stryker.com

**Do not flush cloths**

Not made with natural rubber latex

Sage Products LLC  
3909 Stone Oaks Road • Cary, IL  
60013 • 800 323 2220 • stryker.com  
U.S. Patents: stryker.com/patents

**Reorder #9705**

4 18029 85002 7  
IL52013F

NDC 53462-705-20

**stryker**

Sage  
**2% Chlorhexidine  
Gluconate\* Cloth**  
Patient preoperative  
skin preparation  
**Non-sterile**

\*equivalent to 500 mg chlorhexidine gluconate per cloth

**Rinse-free – alcohol-free  
for external use only – single use**

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application



**2**  
Disposable cloths  
7.5 in x 7.5 in (19.1 cm x 19.1 cm)  
Do not microwave

**Drug Facts**

Active ingredient	Purpose
Chlorhexidine gluconate	Antiseptic
2% solution	Antiseptic

**Uses**

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

**Warnings**

For external use only

**Allergy alert:**  
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing / difficulty breathing
- choking
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbal punctures or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product**

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- do not microwave
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.

**To open package**

- holding top of package in one hand, lift flap on backside of package with other hand
- grasp flap at top and pull down to tear flap away and expose foam
- hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination

6 pack



**Drug Facts (continued)**

**On:**

- Using sterile scissors, cut off and use 1 package.
- Transfer cloths onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination.
- Use a cloth to prepare the skin area indicated on the dry area. Keep the remaining cloths where they will not be contaminated. Use additional cloths to prepare for larger areas.
- Dry surgical sites (such as incisions or arms), use one cloth to cleanse each 100 cm<sup>2</sup> area approximately 1.5-2 inches of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- Model surgical sites (such as inguinal fold) use one cloth to cleanse each 100 cm<sup>2</sup> area approximately 2 x 3 inches of skin to be prepared. Vigorously scrub skin back and forth for 2 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- Discard each cloth after a single use after package has been opened. Discard any unused cloths.

**Other information**

- Store product flat.
- Store between 22°C (72°F) and 30°C (86°F).
- Avoid excessive heat above 40°C (104°F).

**Inactive ingredients**

sterile water, dimethylamine, hydroxyglucuronate-beta-lactone, glycerin, monoglycol diacetate, polypropylene glycol, USP grade preservative.

**Questions or comments?**  
 call toll-free 800-525-2200.  
 Monday to Friday 8:00 am - 5:00 pm CST  
 stryker.com

**stryker**

Sage Products LLC  
 3609 Three Caves Road  
 Cary, Illinois 60013  
 800 525 2200  
 stryker.com  
 U.S. Import: stryker.com/betas

**Reorder #9717**

6 18029 85060 7

NDC 53462-705-26

**stryker**

Sage  
**2% Chlorhexidine Gluconate\* Cloth**  
 Patient preoperative skin preparation  
**Non-sterile**

\*equivalent to 500 mg chlorhexidine gluconate per cloth

**Rinse-free – alcohol-free  
 for external use only – single use**

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

**6**  
 Disposable cloths  
 7.5 in x 7.5 in (19.1 cm x 19.1 cm)  
 Do not microwave

**Drug Facts**

**Active ingredient**  
 Chlorhexidine gluconate  
 2% solution

**Purpose**  
 Antiseptic

**Uses**

- Helps reduce bacteria that can potentially cause skin infection or preparation of skin prior to surgery.

**Warnings**  
 For external use only

**Allergy alert**  
 This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product.
- on further punctures or in contact with the mucous membranes.
- on open skin wounds or as a general skin cleanser.

**When using this product**

- Avoid use of hands, eyes and mouth. May cause irritation or permanent injury if chlorhexidine gluconate is used and washed. If contact occurs, wash with cold water right away and contact a doctor.

Stop use and seek a doctor if irritation, numbness, or allergic reaction occurs. There may be signs of a serious condition.

**Keep out of reach of children**  
 If misused, get medical help or contact a Poison Control Center right away.

**Directions**

- Use with care in premature infants or infants under 2 months of age. These products may cause infection or chemical burns.
- Do not microwave.
- Product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.

**To open package**

- Holding top of package in one hand, lift flap on backside of package with other hand.
- Grasp flap at top and pull down to tear flap away and expose foam.
- Hold outside of package to prevent foam and cloth to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination.

**Do not flush cloths**

**Not made with natural rubber latex.**

# CHLORHEXIDINE GLUCONATE

chlorhexidine gluconate cloth

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53462-705
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	500 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>NONOXYNOL-9</b> (UNII: 48Q180SH9T)	
<b>POLYSORBATE 20</b> (UNII: 7T1F30V5YH)	
<b>DIMETHICONE 350</b> (UNII: 2Y53S6ATLU)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	

**GLYCERIN** (UNII: PDC6A3C0OX)

**GLUCONOLACTONE** (UNII: WQ29KQ9POT)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53462-705-20	2 in 1 PACKAGE; Type 0: Not a Combination Product	02/01/2006	
2	NDC:53462-705-23	6 in 1 PACKAGE; Type 0: Not a Combination Product	02/01/2006	04/01/2024
3	NDC:53462-705-26	6 in 1 PACKAGE; Type 0: Not a Combination Product	02/01/2006	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021669	02/01/2006	

**Labeler** - Sage Products LLC (054326178)

**Registrant** - Sage Products LLC (054326178)

Revised: 12/2025

Sage Products LLC