CHILDRENS PAIN AND FEVER- acetaminophen suspension Walgreen Company

Walgreen Co. Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

1	-	İ
72-95	11 years	15 mL

^{*}or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- · do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Principal Display Panel

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredient in Children's Tylenol® Oral Suspension

Children's Pain & Fever

ACETAMINOPHEN 160 mg PER 5 mL

PAIN RELIEVER / FEVER REDUCER

ORAL SUSPENSION

Aspirin Free

Alcohol Free


Ibuprofen Free

Paraben Free

AGES 2-11 YEARS

4 FL OZ (118 mL)

Cherry flavor





Drug Facts

Active ingredient (in each 5 mL)

Purpose

cetaminophen 160 mg

...Pain reliever/fever reduce

Uses temporarily:

- reduces fever
 relieves minor aches and pains due to:
 the common cold thu headache
 sore throat toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum

- daily amount

 with other drugs containing acetaminophen
 Allergy alert: Acetaminophen may cause severe skin
- reactions. Symptoms may include:

 skin reddening blisters rash
 If a skin reaction occurs, stop use and seek medical help

right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- Do not use

 with any other drug containing acetaminophen
 (prescription or nonprescription). If you are not sure
 whether a drug contains acetaminophen, ask a doctor
 or pharmacist.

 if your child has ever had an allergic reaction to this
 product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

- Stop use and ask a doctor if

 pain gets worse or lasts more than 5 days
 fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present These could be signs of a serious condition

Keep out of reach of children.

verdose warning: In case of overdose, get medical alp or contact a Poison Control Center right away (1-900-222-1222). Quick medical attention is critical ven if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use do not give more than directed (see overdose). warning)
 ■ shake well before using
 ■ mL = milliliter

- mIL = milliliter
 infor light case on chart below. If possible, use weight to dose; otherwise, use age.
 remove the child protective cap and squeeze your child's dose into the dosing cup
 repeat dose every 4 hours while symptoms last
 do not give more than 5 times in 24 hours

Г	Weight (lb)	Age (yr)	Dose (mL)*
Г	under 24	under 2 years	ask a doctor
Г	24-35	2-3 years	5 mL
Γ	36-47	4-5 years	7.5 mL
Г	48-59	6-8 years	10 mL
Г	60-71	9-10 years	12.5 mL
Г	72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL contains: sodium 3 mg
 store at 20-25°C (68-77°F)
 do not use if printed neckband is broken or missing

Inactive ingredients anhydrous citric acid, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

ur pharmacists recommend the Walgreens brand e invite you to compare to national brands. nis product is not manufactured or distributed by phnson & Johnson Consumer Inc., distributor of Children's Tylenol⁸ Oral Suspension



DOSAGE CUP PROVIDED

Gluten Free

NDC 0363-0800-26

w



Compare to the active ingredient in Children's Tylenol® Oral Suspension[®]

Pain &

Fever

ORAL SUSPENSION

Aspirin Free

Alcohol Free

Ibuprofen Free

Paraben Free

AGES 2-11

YEARS

4 FL OZ (118 mL)

ACETAMINOPHEN 160 mg PER 5 mL

PAIN RELIEVER / FEVER REDUCER



DIRECTIONS: STEP 1. Shake well before using

STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.







Cherry

17526 94 C17



CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:0363-0800

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
BUTYLPARABEN (UNII: 3QPI1U3FV8)			
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)			
CARRAGEENAN (UNII: 5C69YCD2YJ)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL (UNII: 506T60A25R)			
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)			

Product Characteristics			
Color	RED (opaque)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0363-0800- 26	1 in 1 CARTON	10/08/1993		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:0363-0800- 28	1 in 1 CARTON	08/23/2013	02/28/2021	
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product			
3	NDC:0363-0800- 34	2 in 1 CARTON	08/19/2013	02/28/2021	
3		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/08/1993	

Labeler - Walgreen Company (008965063)

Revised: 5/2024 Walgreen Company