

**IBUPROFEN- ibuprofen capsule, liquid filled  
Family Dollar (FAMILY WELLNESS)**

-----  
**DRUG FACTS**

**Active ingredient (in each capsule)**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\***

**(present as the free acid and potassium salt)**

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

## Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

## Inactive ingredients

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

## Principal Display Panel

**COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®†**

### Ibuprofen

Liquid Filled Capsules, 200 mg

Pain Reliever/Fever Reducer (**NSAID**)

SOFTGELS\*\*

(\*\*LIQUID FILLED CAPSULES)

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

**TAMPER-EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

DISTRIBUTED BY: MIDWOOD BRANDS, LLC

10611 MONROE RD., MATTHEWS, NC 28105

## Product Label

Lot No.:  
Exp. Date:

Rev# 03/12  
PDU-B7E  
FC004302



**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® LIQUI-GELS®†

NDC 55319-199-80

FAMILY **wellness**™

# Ibuprofen

## Liquid Filled Capsules, 200 mg

### Pain Reliever/Fever Reducer (NSAID)

80 SOFTGELS\*\*  
(\*\*LIQUID FILLED CAPSULES)



ACTUAL SIZE



#### Drug Facts

**Active ingredient (in each capsule)** Purpose  
Sulbized ibuprofen equal to Pain reliever/  
200 mg ibuprofen (NSAID)\* ..... fever reducer  
(present as the free acid and potassium salt) ..... nonsteroidal  
anti-inflammatory drug

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
- menstrual cramps
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Drug Facts (continued)

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Drug Facts (continued)

#### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
  - Take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

**Inactive ingredients** FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

#### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

DISTRIBUTED BY: MIDWOOD BRANDS, LLC  
10611 MONROE RD., MATTHEWS, NC 28105

NOT 100% SATISFIED?

Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

# FAMILY WELLNESS Ibuprofen 200 mg Softgels

## IBUPROFEN

ibuprofen capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-199
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	IB200
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-199-80	1 in 1 BOX	06/30/2017	06/30/2025
1		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55319-199-30	1 in 1 BOX	06/30/2017	06/30/2025
2		30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA078682	06/30/2017	06/30/2025

**Labeler** - Family Dollar (FAMILY WELLNESS) (024472631)

Revised: 10/2022

Family Dollar (FAMILY WELLNESS)