

STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid Hy-Vee

Drug Facts

Active ingredient (in each 30 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
 - 30 mL (1 dose) every 1/2 to 1 hour as needed
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** sodium 14 mg
- **each 30 mL contains:** salicylate 248 mg
- sugar free
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid, sucralose

Questions or comments?

Call toll free **1-800-289-8343 Monday-Friday 9AM-4PM CST**

Principal Display Panel

*COMPARE TO the active ingredient in Pepto-Bismol®

Regular Strength

Stomach Relief

Bismuth Subsalicylate 525mg -

Upset Stomach Reliever / Antidiarrheal

CHERRY Flavor

Relieves:

- **heartburn**
- **Indigestion**
- **Nausea**
- **Upset Stomach**
- **Diarrhea**

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company, Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY HY-VEE, INC.

5820 WESTOWN PARKWAY

WEST DES MOINES, IA 50266

www.hy-vee.com

Package Label

Drug Facts (continued)

Directions ■ mL = milliliter ■ shake well before using
 ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 30 mL (1 dose) every ½ to 1 hour as needed ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years of age: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: sodium 14 mg
 ■ each 30 mL contains: salicylate 248 mg
 ■ sugar free ■ keep tightly closed ■ protect from freezing
 ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid, sucralose

Questions or comments?

Call toll free 1-800-289-8343 Monday-Friday 9AM-4PM CST

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

DISTRIBUTED BY HY-VEE, INC. SATISFACTION GUARANTEED
 5820 WESTOWN PARKWAY OR PURCHASE PRICE REFUNDED
 WEST DES MOINES, IA 50266 CALL 1-800-289-8343 www.hy-vee.com



HyVee
health

***COMPARE TO**
the Active Ingredient in
Pepto-Bismol®

NDC 42507-825-08

Regular Strength Stomach Relief

Bismuth subsalicylate 525 mg -
Upset Stomach Reliever/Antidiarrheal

Cherry Flavor

Relieves:

- Heartburn
- Indigestion
- Nausea
- Upset Stomach
- Diarrhea



8 FL OZ (237 mL)

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL
AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredient (in each 30 mL) Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

Uses relieves ■ diarrhea ■ travelers' diarrhea
 ■ upset stomach due to overindulgence in food and drink, including
 ■ heartburn ■ gas ■ indigestion
 ■ nausea ■ fullness ■ belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have

■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs
 ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) ▶

Hy-Vee Regular Strength Stomach Relief Cherry

STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:42507-825 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-----------------------|-----------------|
| BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ) | BISMUTH SUBSALICYLATE | 525 mg in 30 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| BENZOIC ACID (UNII: 8SKN0B0MIM) | |
| D&C RED NO. 22 (UNII: 1678RKX8RT) | |
| D&C RED NO. 28 (UNII: 767IP0Y5NH) | |
| MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC) | |
| WATER (UNII: 059QF0KO0R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SALICYLIC ACID (UNII: O414PZ4LPZ) | |
| SODIUM SALICYLATE (UNII: WQ1H85SYP) | |
| SORBIC ACID (UNII: X045WJ989B) | |
| METHYLCELLULOSE (1500 CPS) (UNII: PONTE48364) | |
| SUCRALOSE (UNII: 96K6UQ3Z D4) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | CHERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:42507-825-08 | 237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 10/31/2015 | 06/30/2024 |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M008 | 10/31/2015 | 06/30/2024 |

Labeler - Hy-Vee (006925671)

Revised: 4/2024

Hy-Vee