

HEB DAYTIME FLU AND SEVERE COLD AND COUGH- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride powder, for solution

HEB

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HEB® DayTime Flu & Severe Cold & Cough

Drug Facts

<i>Active ingredients (in each packet)</i>	<i>Purposes</i>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant
Phenylephrine hydrochloride 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 4 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor

Age	Dose
children under 4 years of age	do not use
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-

15 minutes.

- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 4 mg, sodium 27 mg
- **phenylketonurics:** contains phenylalanine 34 mg per packet
- store at controlled room temperature 20°-25°C (68°-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, aspartame, citric acid anhydrous, FD&C blue no. 1, FD&C red no. 40, flavors, maltodextrin, sodium citrate anhydrous, sucrose, and pregelatinized starch.

Questions or Comments?

Call 1-866-923-4914

DISTRIBUTED BY H-E-B® , SAN ANTONIO, TX 78204

PRINCIPAL DISPLAY PANEL - 6 Packet Carton

Compare to the active ingredients in Theraflu®
Daytime Severe Cold & Cough*

NDC 37808-539-07

H-E-B®

FLU & SEVERE COLD & COUGH

Daytime

Acetaminophen - Pain reliever/fever reducer

Dextromethorphan HBr - Cough suppressant

Phenylephrine HCl - Nasal decongestant

- Nasal & Sinus Congestion
- Cough
- Body Ache
- Sore Throat Pain
- Headache
- Fever

**Berry Infused with
Menthol & Green Tea Flavors**

**See New Warnings
Information & Directions**

6 Packets



FLU & SEVERE COLD & COUGH

Daytime

Compare to the active ingredients in Theraflu®
Daytime Severe Cold & Cough*

NDC 37808-539-07



FLU & SEVERE COLD & COUGH

Daytime

Acetaminophen - Pain reliever/fever reducer
Dextromethorphan HBr - Cough suppressant
Phenylephrine HCl - Nasal decongestant

- Nasal & Sinus Congestion
- Cough
- Body Ache
- Sore Throat Pain
- Headache
- Fever



Berry Infused with
Menthol & Green Tea Flavors

See New Warnings
Information & Directions **6** Packets



Compare to the active ingredients
in Theraflu®
Daytime Severe Cold & Cough*



FLU & SEVERE COLD & COUGH

Daytime

**TAMPER EVIDENT INNER PACKET
DO NOT USE IF SEALED PACKET
IS TORN OR BROKEN**

*This product is not manufactured or
distributed by Novartis Consumer Health, Inc.
or their affiliates, owner of the registered
trademark Theraflu®.

DISTRIBUTED BY H-E-B®, SAN ANTONIO, TX 78204
MADE IN ISRAEL



20708-1114-0
784

**READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.**

Drug Facts

Active ingredients (in each packet)

Active ingredients (in each packet)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg.....	Cough suppressant
Phenylephrine hydrochloride 10 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 4 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are • taking the blood thinning drug warfarin

When using this product • do not exceed recommended dosage

Stop use and ask a doctor if

- fever gets worse or lasts more than 3 days
- nervousness, dizziness, or sleeplessness occurs
- new symptoms occur
- redness or swelling is present
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not use more than directed
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor

Age	Dose
children under 4 years of age	do not use
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 4 mg, sodium 27 mg
- phenylketonurics: contains phenylalanine 34 mg per packet
- store at controlled room temperature 20°-25°C (68°-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, aspartame, citric acid anhydrous, FD&C blue no. 1, FD&C red no. 40, flavors, maltodextrin, sodium citrate anhydrous, sucrose, and pregelatinized starch.

Questions or Comments?

Call 1-866-923-4914

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

HEB DAYTIME FLU AND SEVERE COLD AND COUGH

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-539
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	650 mg
Dextromethorphan hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan hydrobromide	20 mg
Phenylephrine hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine hydrochloride	10 mg

Inactive Ingredients

Ingredient Name	Strength
acesulfame potassium (UNII: 23OV73Q5G9)	
aspartame (UNII: Z0H242BBR1)	
anhydrous citric acid (UNII: XF417D3PSL)	
FD&C blue no. 1 (UNII: H3R47K3TBD)	
FD&C red no. 40 (UNII: WZB9127XOA)	
maltodextrin (UNII: 7CVR7L4A2D)	
anhydrous trisodium citrate (UNII: RS7A450LGA)	
sucrose (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-539-07	6 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/28/2013	

Labeler - HEB (007924756)