

**ONELAX- bisacodyl suppository**  
**AKRON PHARMA INC**

-----  
**OneLAX**  
**BISACODYL 10 mg Suppositories**  
**Fast Acting Stimulant Laxative**

**Active Ingredients (per suppository)**

Bisacodyl USP, 10 mg

**Purpose**

Laxative

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Uses**

- for relief of occasional constipation
- this product usually produces bowel movement in 1/4 to 1 hour

**Warnings**

For rectal use only.

**Do not use laxative products**

- when abdominal pain, nausea or vomiting are present
- for a period longer than one week

**Ask a doctor before use if you have**

noticed a sudden change in bowel habits that persist over a period of 2 weeks

**When using this product**

it may cause abdominal discomfort, faintness, rectal burning and mild cramps

**Stop use and ask a doctor if**

rectal bleeding occurs, or you fail to have a bowel movement after using a laxative. This may indicate a serious condition.

**If pregnant or breast-feeding**

ask a health professional before use.

**Directions**

- adults and children 12 years of age and older
- detach one suppository from the strip
- remove from wrapper before inserting into the rectum
- the rectal suppository dose is one suppository per day or as directed by a doctor
- children under 12 years of age consult a doctor

**Other information**

store below 30° C (86° F)

**Inactive ingredient**

hydrogenated vegetable oil

**Manufactured for:**

Akron Pharma Inc.,

Fairfield, NJ-07004

[www.akronpharma.com](http://www.akronpharma.com)

**Inactive ingredient** hydrogenated vegetable oil

**Other information** ■ store at temperatures below 30°C (86°F)

Children under 12 years  
 Adults and children 12 years and over ■ 1 suppository once daily  
 Ask a doctor

**Directions**  
 ■ tear off one suppository from the strip  
 ■ lean and peel the wrapper in a downward motion  
 ■ continue leaning downward to almost the full length of the suppository (see illustration at the right)  
 ■ gently remove the suppository from the wrapper  
 ■ lie on your side and push suppository, with pointed end first, high into the rectum so it will not slip out  
 ■ remain in for 15 to 20 minutes  
 ■ Do not exceed recommended dose ■ If suppository seems soft, place in refrigerator for a short time before use

**Warnings** For rectal use only  
 Do not use when abdominal pain, nausea, or vomiting are present.  
 Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.  
 When using this product you may experience ■ rectal burning ■ mild cramps ■ abdominal discomfort ■ flatness  
 Stop use and ask a doctor if ■ you have rectal bleeding ■ you fail to have a bowel movement after using a laxative. These may be signs of a serious condition.  
 ■ you need to use a laxative for more than one week  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Drug Facts** (continued)

**Active ingredient (in each suppository)** Bisacodyl USP 10 mg  
 Stimulant laxative

**Purpose** ■ relieves occasional constipation ■ this product generally produces a bowel movement in 15 minutes to 1 hour

NDC 71399-8460-2  
 Compare to the active ingredient in Dulcolax® Suppositories\*

# OneLAX™

## Bisacodyl 10mg Suppositories

Fast Acting Stimulant Laxative

**12**  
**SUPPOSITORIES**

TAMPER EVIDENT: SUPPOSITORIES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED.

**Bisacodyl 10mg Suppositories**

**Akron Pharma**  
 Manufactured for: Akron Pharma Inc., Fairfield, NJ-07004  
 www.akronpharma.com  
 Questions or comments? toll-free (877)225-6999

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, owner of the registered trademark Dulcolax®





# ONELAX

bisacodyl suppository

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71399-8460
<b>Route of Administration</b>	RECTAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg

**Inactive Ingredients**

Ingredient Name	Strength
FAT, HARD (UNII: 8334LX7S21)	

**Product Characteristics**

Color	white (WHITE TO OFF WHITE)	Score	
Shape	OVAL (TORPEDO SHAPE)	Size	
Flavor		Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399-8460-2	12 in 1 BOX	09/13/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:71399-8460-5	50 in 1 BOX	09/13/2024	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	09/13/2024	

**Labeler** - AKRON PHARMA INC (067878881)

Revised: 11/2025

AKRON PHARMA INC